



# **Influence of Different Planting Dates and Mulches on Growth, Yield and Economics of Potato**

**Gopal Swaroop Pathak<sup>a++\*</sup>, Shrish Kumar Singh<sup>a#</sup> and C. P. Singh<sup>at</sup>**

<sup>a</sup> Department of Agronomy, Tilak Dhari Post Graduate College, Jaunpur, Uttar Pradesh, India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/JEAI/2024/v46i42336

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/113581>

**Original Research Article**

**Received: 16/12/2023**

**Accepted: 22/02/2024**

**Published: 29/02/2024**

## **ABSTRACT**

The present research work was conducted at Pili kothi farm, Department of Agronomy, Tilak Dhari Post Graduate College, Jaunpur, Uttar Pradesh during winter season of 2018 and 2019. The experiment was laid out in Factorial Randomized Block Design. The treatments comprised of different planting dates viz. 20 days early planting on 17<sup>th</sup> October, 10 days early planting on 27<sup>th</sup> October, Optimum planting time on 7<sup>th</sup> November, 10 days late planting on 17<sup>th</sup> November and 20 days late planting on 27<sup>th</sup> November and different mulching practices viz. Organic mulch (Paddy straw at 5 t/ha), Plastic mulch (Black colour) and Control (Without mulch) of potato. There were 15 treatments and each were replicated to four folds. These treatments were randomly arranged in each replication and divided into sixty plots. The results revealed that the significantly maximum number of haulms/hill at 45 DAP (4.50, 4.70 and 4.60), plant height (52.30, 54.00 and 53.10 cm), dry weight/ (27.90, 28.20 and 28.00) plant at 60 DAP, weight of tubers/plant (496.40, 504.10 and 500.30), weight of tubers/plant (>75 g) (207.70, 209.00 and 208.30) and tuber yield (369.30, 370.70

<sup>++</sup> Research Scholar;

<sup>#</sup> Professor and Head;

<sup>†</sup> Assistant Professor;

\*Corresponding author: E-mail: gswaroopathak@gmail.com;

and 370.00 q/ha) were recorded under 20 Days Early Planting On 17<sup>th</sup> October among different planting dates during first, second year and mean data also. Whereas among the mulching practices significant variation were recorded among the all parameter. However maximum number of haulms/hill (3.60, 3.70 and 3.70) at 45 DAP, plant height (50.20, 52.0 and 51.10 cm), dry weight/plant (25.5, 25.9 and 25.7) at 60 DAP, weight of tubers/plant (454.80, 460.70 and 457.70), weight of tubers/plant (>75 g) (191.90, 193.30 and 192.60) and tuber yield (333.80, 334.90 and 334.30 q/ha) under the organic mulch (paddy straw at 5 t/ha) respectively, during first, second year as well as in mean data also.

**Keywords:** Date of planting; mulching; potato; growth attributes; yield attributes; economics.

## 1. INTRODUCTION

“Potato (*Solanum tuberosum* L.) is the third most important food crop in consumption in the world after rice and wheat” (Hancock et al., 2014). “The higher profit from the potato crop under rice fallows not only depends on the management practices, but it also depends on soil moisture content and temperature conditions. Because potato crop is very sensitive to lack of soil moisture and temperature which affects the growth and developmental process, and finally leads to higher and lower tuber yield. The optimum date of planting for potato is highly location-specific and every production region has an ‘optimum’ planting period during which conditions are most favourable for giving the higher yield” [1]. “Mulching improved seedling emergence, plant height and numbers of stems” (Singh and Ahmed, 2008). “Potato is grown during the winter season when rainfall is scarce and irrigation become essential for providing sufficient moisture to the growing crop. To minimize the cultivation cost mulching could be effectively used instead of irrigation. Mulching is one of the important agronomic practice beneficial in conserving the soil moisture, suppressing the weeds, improving soil fertility” [2]. “Organic mulch are efficient in reduction of nitrates leaching, improve soil physical properties, prevent erosion, supply organic matter, regulate temperature and water retention, improve nitrogen balance, take part in nutrient cycle as well as increase the biological activity” [3]. Mulching had significant influence on potato growth and yield. Considering this, an attempt was made to find out the suitable planting date of potato with mulching for higher productivity.

## 2. MATERIALS AND METHODS

A field experiment entitled “Influence of different planting dates and mulches on growth,

yield and economics of potato” was conducted at crop research farm (Pili kothi) of Agronomy Department, Tilak Dhari Post Graduate College, Jaunpur (U.P.) during *rabi* season of 2018-19 and 2019-20. There were 15 treatments and each were replicated to four folds. These treatments were randomly arranged in each replication and divided into sixty plots. The gross plot size was 5 m × 3 m (15 m<sup>2</sup>). The pre-sowing irrigation was done after the harvest of the preceding crop. The whole experimental field was irrigated and left for four days. So that the rhizosphere attain moisture and good soil structure. For getting good condition of field ploughing was done along with planking in each ploughing, which helps in breaking the clods and levelling the land for proper seedbed. Previous crop residues or stubbles and weeds were removed with the help of rake and also partial work done manually by hand. The row to row and plant to plant seed potatoes were planted in each unit plot maintaining a spacing of 60 cm × 20 cm. The fertilizer inputs used are urea as a source of nitrogen, diammonium phosphate (DAP) as a source of phosphorus, muriate of potash (MOP) as a source of potassium. 120 kg N, 60 kg P<sub>2</sub>O<sub>5</sub> and 80 kg K<sub>2</sub>O/ha are used as recommended dose of fertilizers during the experiment and fertilizer requirement calculated individually on the basis of each plot. Organic mulching was done after top dressing and polythene sheet (Plastic mulch) was placed on the ridge by making holes at the planting time. Height of the five randomly selected plants was measured with the help of meter scale from soil surface to the apical bud at 60 days after planting and the mean values were taken in cm. Five selected plants were taken for fresh weight at the time of haulm cutting. For dry weight they were firstly dried in an oven at 60°C temperature. After complete drying, average weight of plant (g) was weighed by electronic balance.

### 3. RESULTS AND DISCUSSION

#### 3.1 Number of Haulms/hill at 45 DAP

The data pertaining to number of haulms/hill as influenced by different dates of planting and mulching were presented in Table 1. The data on the number of haulms at 45 DAP as influenced by different planting dates and mulching showed significant difference. However, potato planted on 20 days early planting on 17 October recorded significantly higher (4.50, 4.70 and 4.60) number of haulms/hill at 45 DAP and none of the treatments found statistically at par value during first, second as well as in mean data also. The data on mulches shown that significantly higher number of haulms at 45 DAP was recorded in Organic mulch (Paddy straw at 5 t/ha) (3.60, 3.70 and 3.70) during all experimental years which was followed by Plastic mulch (Black colour) (3.40, 3.50 and 3.40) during first year and second year except mean data. That may be attributed to rising soil temperature and conservation of more moisture in the black polythene covered soil that accelerated the germination and emergence of potato. Besides, mulching provides favourable physio-chemical conditions of soil like good soil aeration, porosity, water holding capacity, better availability of nutrients. Similar research finding also reported by Zahed et al. [4].

#### 3.2 Plant Height (cm) at 60 DAP

At 60 DAP, 20 days early planting on 17<sup>th</sup> October recorded significantly higher plant height (52.30, 54.00 and 53.10 cm) in 2018, 2019 and in mean data, respectively. None of the at par findings was reported at this stage. At 60 DAP, same trend was followed in all of the treatments following mulching practices in all the years. Organic mulch (Paddy straw at 5 t/ha) performed significantly maximum plant height (50.20, 52.0 and 51.10 cm) in all of the experimental years. Similarly, at par values lead to the Plastic mulch (Black colour) plots (49.60, 51.40 and 50.50 cm), respectively. This might be high moisture regime due to irrigation under this treatment which facilitated better water and nutrient uptake by the plants. Application of mulch also significantly produced better plant growth as compared to non-mulch condition [5]. Early planting resulted in tallest plant as during this stage, air temperature was maximum as compared to rest of the sowing dates Ilyas and Ayub [6].

#### 3.3 Dry Weight/Plant at 60 DAP

At 60 DAP, data analysed shown significant difference among all treatments of different dates of planting. In first year (2018), second year (2019) and in mean analysis, 20 days early planting on 17<sup>th</sup> October recorded significantly higher dry weight/plant (27.90, 28.20 and 28.00) and there was no at par values reported. Whereas, significantly higher dry weight/plant was obtained in Organic mulch (Paddy straw at 5 t/ha) treated plot (25.5, 25.9 and 25.7) and at par values are reported in Plastic mulch (Black colour) treated plot (25.0, 25.4 and 25.2), respectively. The growth duration was shortened due to delay in planting date, hence, dry matter accumulation, tuber fresh weight and final yield were all decreased because of insufficient cumulative temperature over the shorter growing periods Similar findings also observed by Gogoi and Ray (2019). The increase in dry matter can be attributed to the fact that at earlier stages, most of the dry matter was concentrated in leaf and stem and from tuber initiation to tuber development it accumulated in stolon. Higher interception of solar radiation in earlier sowing may be responsible for more dry matter accumulation per plant when soil moisture and light are non-limiting, higher temperatures accelerate growth process [7].

#### 3.4 Weight of Tubers/Plant (g)

The data regarding weight of tubers per plant represented significant difference among all the treatments of different dates of planting. In first year, second year and in mean analysis, 20 days early planting on 17<sup>th</sup> October recorded significantly higher weight of tubers/plant (496.40, 504.10 and 500.30) and there was no at par values recorded among the treatments. Among various mulching practices, significantly higher weight of tubers/plant was recorded in Organic mulch (Paddy straw at 5 t/ha) treated plot (454.80, 460.70 and 457.70) and none of treatments found at par values. The enhanced nutrient availability and microbial activity in mulch conditions especially in early growing stage helps in more number of stolon formation and the increased nutrient uptake lead to the better development that leads to improved size and weight of tubers. The above results are in accordance with the findings Zahed et al. [4].

**Table 1. Effect of different dates of planting and different mulches on growth attributes of potato**

| Treatments   | Number of haulms/hill at 45 DAP |      |      | Plant height (cm) at 60 DAP |       |       | Dry weight/plant at 60 DAP |       |       |
|--|---------------------------------|------|------|-----------------------------|-------|-------|----------------------------|-------|-------|
|  | 2018                            | 2019 | Mean | 2018                        | 2019  | Mean  | 2018                       | 2019  | Mean  |
| <b>Date of Planting (Factor : A)</b>               |                                 |      |      |                             |       |       |                            |       |       |
| 20 Days Early Planting On 17 <sup>th</sup> October | 4.50                            | 4.70 | 4.60 | 52.30                       | 54.00 | 53.10 | 27.90                      | 28.20 | 28.00 |
| 10 Days Early Planting On 27 <sup>th</sup> October | 3.80                            | 4.10 | 3.90 | 50.60                       | 51.80 | 51.20 | 26.70                      | 27.00 | 26.80 |
| Optimum Planting Time On 7 <sup>th</sup> November  | 3.20                            | 3.30 | 3.30 | 49.70                       | 51.40 | 50.50 | 24.50                      | 24.90 | 24.70 |
| 10 Days Late Planting On 17 <sup>th</sup> November | 2.90                            | 3.00 | 2.90 | 48.80                       | 50.70 | 49.80 | 23.60                      | 23.80 | 23.70 |
| 20 Days Late Planting On 27 <sup>th</sup> November | 2.40                            | 2.50 | 2.50 | 47.00                       | 48.60 | 47.80 | 22.90                      | 23.20 | 23.10 |
| F test   | S                               | S    | S    | S                           | S     | S     | S                          | S     | S     |
| SE(m)±   | 0.14                            | 0.15 | 0.14 | 0.38                        | 0.42  | 0.36  | 0.22                       | 0.27  | 0.24  |
| CD (p=0.05)  | 0.41                            | 0.43 | 0.40 | 1.10                        | 1.21  | 1.03  | 0.64                       | 0.76  | 0.68  |
| <b>Different Mulches (Factor : B)</b>              |                                 |      |      |                             |       |       |                            |       |       |
| Organic Mulch (Paddy Straw at 5 t/ha)              | 3.60                            | 3.70 | 3.70 | 50.20                       | 52.0  | 51.10 | 25.5                       | 25.9  | 25.7  |
| Plastic Mulch (Black Colour)                       | 3.40                            | 3.50 | 3.40 | 49.60                       | 51.40 | 50.50 | 25.0                       | 25.4  | 25.2  |
| Control (Without Mulch)                            | 3.20                            | 3.30 | 3.20 | 49.30                       | 50.40 | 49.80 | 24.8                       | 25.1  | 24.9  |
| F test   | S                               | S    | S    | S                           | S     | S     | S                          | S     | S     |
| SE(m)±   | 0.11                            | 0.12 | 0.11 | 0.73                        | 0.32  | 0.26  | 0.17                       | 0.21  | 0.18  |
| CD (p=0.05)  | 0.32                            | 0.33 | 0.31 | 0.23                        | 0.93  | 0.76  | 0.50                       | 0.59  | 0.53  |

**Table 2. Effect of different dates of planting and different mulches on yield attributes of potato**

| Treatments   | Weight of tubers/plant (g) |        |        | Weight of tubers/plant (>75 g) |        |        | Total tuber yield (q/ha) |        |        |
|--|----------------------------|--------|--------|--------------------------------|--------|--------|--------------------------|--------|--------|
|  | 2018                       | 2019   | Mean   | 2018                           | 2019   | Mean   | 2018                     | 2019   | Mean   |
| <b>Date of planting (Factor: A)</b>                |                            |        |        |                                |        |        |                          |        |        |
| 20 Days Early Planting On 17 <sup>th</sup> October | 496.40                     | 504.10 | 500.30 | 207.70                         | 209.00 | 208.30 | 369.30                   | 370.70 | 370.00 |
| 10 Days Early Planting On 27 <sup>th</sup> October | 475.30                     | 483.30 | 479.30 | 200.30                         | 201.90 | 201.10 | 355.90                   | 357.60 | 356.80 |
| Optimum Planting Time On 7 <sup>th</sup> November  | 449.40                     | 456.10 | 452.80 | 189.70                         | 191.10 | 190.40 | 333.10                   | 334.10 | 333.60 |
| 10 Days Late Planting On 17 <sup>th</sup> November | 420.00                     | 427.80 | 423.90 | 178.80                         | 180.60 | 179.70 | 305.30                   | 307.40 | 306.40 |
| 20 Days Late Planting On 27 <sup>th</sup> November | 385.60                     | 390.80 | 388.20 | 166.30                         | 167.70 | 167.00 | 280.50                   | 282.40 | 281.50 |
| F test   | S                          | S      | S      | S                              | S      | S      | S                        | S      | S      |
| SE(m)±   | 2.35                       | 2.07   | 2.11   | 1.39                           | 1.20   | 1.27   | 2.51                     | 2.31   | 2.06   |
| CD (p=0.05)  | 6.73                       | 5.94   | 6.05   | 3.98                           | 3.44   | 3.65   | 7.19                     | 6.62   | 5.91   |
| <b>Different mulches (Factor: B)</b>               |                            |        |        |                                |        |        |                          |        |        |
| Organic Mulch (Paddy Straw at 5 t/ha)              | 454.80                     | 460.70 | 457.70 | 191.90                         | 193.30 | 192.60 | 333.80                   | 334.90 | 334.30 |
| Plastic Mulch (Black Colour)                       | 445.80                     | 453.60 | 449.70 | 189.60                         | 191.00 | 190.30 | 328.80                   | 330.50 | 329.60 |
| Control (Without Mulch)                            | 435.60                     | 442.90 | 439.20 | 184.30                         | 185.90 | 185.10 | 324.00                   | 326.00 | 325.00 |
| F test   | S                          | S      | S      | S                              | S      | S      | S                        | S      | S      |
| SE(m)±   | 1.82                       | 1.61   | 1.64   | 1.08                           | 0.93   | 0.99   | 1.94                     | 1.79   | 1.60   |
| CD (p=0.05)  | 5.21                       | 4.60   | 4.69   | 3.08                           | 2.66   | 2.82   | 5.57                     | 5.13   | 4.58   |

**Table 3. Effect of dates of planting and different mulching on economics in potato (Mean data of two year)**

| Treatments   | Economics                    |                        |                      |           |
|--|------------------------------|------------------------|----------------------|-----------|
|  | Cost of cultivation (Rs./ha) | Gross returns (Rs./ha) | Net returns (Rs./ha) | B:C ratio |
| 20 Days Early Planting on 17 <sup>th</sup> October + Organic mulch (Paddy straw at 5 t/ha) | 75,046.00                    | 2,98,700.00            | 2,23,654.00          | 2.98      |
| 20 Days Early Planting on 17 <sup>th</sup> October + Plastic mulch (Black colour)          | 1,02,546.00                  | 2,95,900.00            | 1,93,354.00          | 1.89      |
| 20 Days Early Planting on 17 <sup>th</sup> October + Control (Without mulch)               | 73,596.00                    | 2,91,800.00            | 2,18,204.00          | 2.96      |
| 10 Days Early Planting on 27 <sup>th</sup> October + Organic mulch (Paddy straw at 5 t/ha) | 75,046.00                    | 2,88,600.00            | 2,13,554.00          | 2.85      |
| 10 Days Early Planting on 27 <sup>th</sup> October + Plastic mulch (Black colour)          | 1,02,546.00                  | 2,86,200.00            | 1,83,654.00          | 1.79      |
| 10 Days Early Planting on 27 <sup>th</sup> October + Control (Without mulch)               | 73,596.00                    | 2,81,400.00            | 2,07,804.00          | 2.83      |
| Optimum Planting time on 7 <sup>th</sup> November + Organic mulch (Paddy straw at 5 t/ha)  | 75,046.00                    | 2,70,900.00            | 1,95,854.00          | 2.61      |
| Optimum Planting time on 7 <sup>th</sup> November + Plastic mulch (Black colour)           | 1,02,546.00                  | 2,67,400.00            | 1,64,854.00          | 1.61      |
| Optimum Planting time on 7 <sup>th</sup> November + Control (Without mulch)                | 73,596.00                    | 2,62,300.00            | 1,88,704.00          | 2.57      |
| 10 Days Late Planting on 17 <sup>th</sup> November + Organic mulch (Paddy straw at 5 t/ha) | 75,046.00                    | 2,50,500.00            | 1,75,454.00          | 2.34      |
| 10 Days Late Planting on 17 <sup>th</sup> November + Plastic mulch (Black colour)          | 1,02,546.00                  | 2,44,600.00            | 1,42,054.00          | 1.39      |
| 10 Days Late Planting on 17 <sup>th</sup> November + Control (Without mulch)               | 73,596.00                    | 2,40,200.00            | 1,66,604.00          | 2.27      |
| 20 Days Late Planting on 27 <sup>th</sup> November + Organic mulch (Paddy straw at 5 t/ha) | 75,046.00                    | 2,28,600.00            | 1,53,554.00          | 2.05      |
| 20 Days Late Planting on 27 <sup>th</sup> November + Plastic mulch (Black colour)          | 1,02,546.00                  | 2,24,400.00            | 1,21,854.00          | 1.19      |
| 20 Days Late Planting on 27 <sup>th</sup> November + Control (Without mulch)               | 73,596.00                    | 2,22,500.00            | 1,48,904.00          | 2.03      |

### 3.5 Weight of Tubers/Plant (>75 g)

Weight of tubers/plant (>75 g) of potato significantly influenced due to different dates of planting and different mulching practices. Among different dates of planting significantly higher weight of tubers/plant (>75 g) (207.70, 209.00 and 208.30) was found in 20 days early planting on 17<sup>th</sup> October and none of the treatments found statistically at par value during all the years. Whereas regarding mulches, significantly maximum weight of tubers/plant (>75 g) (191.90, 193.30 and 192.60) was recorded under Organic mulch (Paddy straw at 5 t/ha) which was statistically at par with Plastic mulch (Black colour) (189.60, 191.00 and 190.30) during both the years of experimentation and also among mean data.

### 3.6 Total Tuber Yield (q/ha)

The data pertaining to number of haulms/hill as influenced by different dates of planting and mulching were presented in Table 2. Dates of planting represented statistically significant difference among various treatments. Significantly higher tuber yield (369.30, 370.70 and 370.00 q/ha) of potato was recorded under 20 days early planting on 17<sup>th</sup> October treated plot during the experiment in all of the years, which was significantly superior over rest of the treatments. Among various mulching practices, significantly maximum tuber yield was recorded in Organic mulch (Paddy straw at 5 t/ha) (333.80, 334.90 and 334.30 q/ha) which was statistically at par Plastic mulch (Black colour) (328.80, 330.50 and 329.60 q/ha) during both years and also mean data.

The early planted crop get warmer temperature during its vegetative growth and cooler temperature during its tuber development stage. This climatic suitability increases the tuber yield of potato. Subsequently late planted potato caught under unfavourable climatic conditions hence the tuber yield decreases. Similar results also reported by Malunekar and Solanke [8]. The straw mulch used after planting was the best mulching material for reaching the highest yield [9]. The maximum tuber yield in 10 October sown crop due to optimum moisture and temperature condition along with good growth of plant which were not prevailing during late sown crop also observed by (Haile et al., [10]), Thongam et al., [11] and Navneet et al., (2020). This might be due to difference in physiological age of the seed

tuber used for planting [1]. Mulching also resulted in higher tuber yield in all the yield categories over no mulching. Such beneficial effects of mulching have been demonstrated in the results achieved by Bijeta and Raghav [12]. Potato is reported to increase tuber yield when plants are mulched with plastics and straw mulch compared with plants grown in bare soil [13]. Total number of tubers decreased with delayed planting as compared to the optimum date of planting. The results confirm the findings of Singh et al. [14].

### 3.7 Cost of Cultivation (Rs./ha)

Among the different treatment combinations, cost of cultivation was recorded highest in the treatment of 20 days early planting on 17<sup>th</sup> October with Plastic mulch (Black colour) (Rs. 1,02,546.00/ha) found similar under mean data. While, least was accounted in the treatment where 20 days early planting was done on 17<sup>th</sup> October + Control (Without mulch) (Rs.73,596.00/ha), under mean data of study.

Mulching with paddy straw recorded increment in net returns, benefit: cost ratio, economic efficiency and production efficiency over no mulch. Similar results also reported by Satapathy et al. [1]

### 3.8 Gross Returns (Rs./ha)

The data pertaining to gross returns revealed that mulching with paddy straw and early date of planting (17<sup>th</sup> October) fetched maximum gross returns (2,98,700.00/ha) in mean data as compared to rest of the treatments. However, 20 days late planting of potato on 27<sup>th</sup> November without mulch obtained lowest readings (Rs. 2,28,600.00/ha).

### 3.9 Net Returns (Rs./ha)

A cursory glance showed that net returns recorded maximum of Rs. 2,23,654.00/ha) were accrued with the application of Organic mulch (Paddy straw at 5 t/ha) and 20 days early planting on 17<sup>th</sup> October during mean data. Where treatment combination of 20 days late planting on 27<sup>th</sup> November + Plastic mulch (Black colour) obtained least net returns of Rs. 1,21,854.00/ha mean data of two years, [15,16].

### 3.10 B: C ratio

As the gross returns and net returns were high benefit cost analysis are also high. However, treatment combination of 20 days early planting on 17<sup>th</sup> October + Organic mulch (Paddy straw at 5 t/ha) recorded highest B:C ratio (2.98) similar under mean data. Whereas, 20 days late planting on 27<sup>th</sup> November + Plastic mulch (Black colour) treatment received less benefit cost ratio (1.19).

### 4. CONCLUSION

From the present study it can be concluded that planting of potato on 17<sup>th</sup> October gave superior performance than other date of planting along with mulching with 5 ton/ha paddy straw was useful in enhancing the growth attributes, yield attributes, yield as well as economics of potato crop.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

- Satapathy BS, Singh T, PUN KB, Saika K, Rautaray SK. Influence of dates of planting and mulching on growth and tuber yield of potato (*Solanum tuberosum*) in rice (*Oryza sativa*) fallows of shallow lowland of Asom. Indian Journal of Agronomy. 2016;61(4): 455-459.
- Pulok Md A, Roy TS, Bhuiyan Md SR, Haque Md N. Effect of potassium and mulches on growth, yield and economics of potato. Potato J. 2016; 43(2):200-210.
- Bhardwaj RL. Effect of mulching on crop production under rainfed condition - A Review. Agri. Reviews. 2013;34(3):188-197.
- Zahed Z, Mufti S, Mushtaq F, Narayan S. Impact of organic and inorganic synthetic mulches on growth and yield of potato under temperate conditions. Biological Forum – An International Journal. 2021; 13(3a):731-737.
- Begum M, Saikia M. Effect of irrigation and mulching on growth and yield attributes of potato. Agric. Sci. Digest. 2014;34(1):76-78.
- Ilyas M, Ayub G. Role of planting depth and mulching on growth and yield components of autumn potato crop sown at different dates. Pure Appl. Biol. 2017; 6(4):1436-1449.
- Vishwas U, Rathiya PS, Sinha AK, Verma C, Gupta A. Response of different date of planting on growth, yield and economics of potato (*Solanum tuberosum* L.) genotypes under Northern hill region of Chhattisgarh. Journal of Pharmacognosy and Phytochemistry. 2020;9(3): 1203-1205.
- Malunjkar BD, Solanke AV. Response of type of tubers used with different sowing dates on growth and yield of potato (*Solanum tuberosum* L.). Internat. J. Agric. Sci. 2007;3(2): 186-188.
- Petr D, Tomasek I, Hamouz K. Cultivation of organic potatoes with the use of mulching materials. Zeszyty Problemowe Postqpow Nauk Rolniczych. 2010;557:95-102.
- Haile B, Mohammed A, Woldegiorgis G. Effect of planting date on growth and tuber yield of potato varieties at Anderacha District, Southwestern Ethiopia. International Journal of Research in Agricultural Sciences. 2012;2:2348-3997.
- Thongam B, Kadam AS, Singh AA, Singh YH. Influence of planting dates on growth and yield of potato (*Solanum tuberosum* L.). Journal of Pharmacognosy and Phytochemistry. 2017;6(6): 1243-1246.
- Bijeta, Raghav M. Comparison of potato yield, quality, weed control and economics obtained with different mulch materials. The Pharma Innovation Journal. 2019; 8(6):512-518.
- Bharat, Kumar R. Effect of various types of mulching on growth, yield and quality of different processable cultivars of potato (*Solanum tuberosum* L.). International Journal of Chemical Studies. 2021;9(3): 166-169.
- Singh SP, Kumar S, Tomar SK, Rao AP. Effect of different sowing dates and varieties on yield, size and number of tubers of potato (*Solanum tuberosum* L.). J. Agric. Res. Technol. 2022;1:137-142.



15. Jaysawal N, Singh DG, Kaniija A, Debbarma B. Effect of different mulches on growth and yield of carrot (*Daucus carota* L.). International Journal of Chemical Studies. 2018;6(4):381-384.
16. Singh N, Singh A, Singh K. Effect of time of planting on growth and yield parameters of potato crop. Int. J. Curr. Microbiol. App. Sci. 2020;9(5): 2847-2851.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

*<https://www.sdiarticle5.com/review-history/113581>*