



Analysis of Norm-Attainability and Convergence Properties of Orthogonal Polynomials in Weighted Sobolev Spaces

Mogoi N. Evans^{a*}, Amos Otieno Wanjara^a
and Samuel B. Apima^a

^a Department of Mathematics and Statistics, Kaimosi Friends University, Kenya.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ARJOM/2024/v20i4792

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/113722>

Received: 03/02/2024

Accepted: 06/04/2024

Published: 09/04/2024

Original Research Article

Abstract

This paper explores norm-attainability of orthogonal polynomials in Sobolev spaces, investigating properties like existence, uniqueness, and convergence. It establishes the convergence of these polynomials in Sobolev spaces, addressing orthogonality preservation and derivative behaviors. Spectral properties, including Sturm-Liouville eigenvalue problems, are analyzed, enhancing the understanding of these polynomials. The study incorporates fundamental concepts like reproducing kernels, Riesz representations, and Bessel's inequality. Results contribute to the theoretical understanding of orthogonal polynomials, with potential applications in diverse mathematical and computational contexts.

Keywords: Orthogonal polynomials; Sobolev spaces; norm-attainability; Hilbert space; weighted Sobolev spaces; Sturm-Liouville eigenvalue problem; convergence of orthogonal polynomials; reproducing kernel; Bessel's inequality; Sobolev embedding; derivative properties of orthogonal polynomials; uniqueness of orthogonal polynomials; compactness of embeddings; pointwise convergence; Riesz representation.

*Corresponding author: E-mail: mogoievans4020@gmail.com;

Asian Res. J. Math., vol. 20, no. 4, pp. 1-7, 2024

2010 Mathematics Subject Classification: 53C25, 83C05, 57N16.

1 Introduction

This research paper investigates the norm-attainability of orthogonal polynomials within Sobolev spaces, motivated by their fundamental role in approximation theory and numerical analysis [1-9]. The introduction highlights the significance of understanding the conditions under which the norm of orthogonal polynomials can be attained, emphasizing the broader implications for computational algorithms and mathematical analysis [10,11,5,12,13,14,15]. The research objectives focus on establishing foundational theorems, propositions, lemmas, and corollaries to characterize the norm-attainability phenomenon. The methodology comprises a thorough literature review, the formulation of a theoretical framework, rigorous proof construction, and, optionally, numerical experiments for validation. The approach aims to contribute to functional analysis, providing insights that bridge theoretical developments with practical applications in various mathematical domains.

2 Preliminaries

In this section, we establish the foundational concepts and results that form the basis of our investigation into the norm-attainability of orthogonal polynomials in Sobolev spaces.

2.1 Hilbert spaces and orthogonal polynomials

We begin by introducing the notion of Hilbert spaces, which are essential mathematical structures for our analysis. A Hilbert space \mathcal{H} is a complete inner product space. Our investigation will be conducted in such spaces, emphasizing their role in the study of orthogonal polynomials.

Theorem 2.1 (Existence of orthogonal Polynomials). *Let \mathcal{H} be a Hilbert space, and consider a set of orthogonal polynomials $\{P_n\}_{n=0}^{\infty}$ defined on a compact interval $[a, b]$. Then, there exists a unique sequence of polynomials that are orthogonal with respect to a weight function $w(x)$ in \mathcal{H} .*

2.2 Sobolev spaces and embeddings

Our analysis extends to Sobolev spaces, which provide a framework for studying functions with certain smoothness properties. Understanding the embedding of these spaces is crucial for our exploration.

Theorem 2.2 (Sobolev Embedding). *For any $p \geq 1$, there exists a constant C_p such that the Sobolev space $W^{m,p}([a, b])$ is continuously embedded in $C^k([a, b])$ for $k < m - \frac{n}{p}$.*

2.3 Norm-attainability of orthogonal polynomials

Our main focus lies in the norm-attainability of orthogonal polynomials within Sobolev spaces. We introduce key results that form the backbone of our investigation.

Theorem 2.3 (Norm-Attainability of Orthogonal Polynomials). *Suppose $\{P_n\}_{n=0}^{\infty}$ is a sequence of orthogonal polynomials in $L^2([a, b], w(x)dx)$. Then, the norm of each polynomial P_n is attained, i.e., there exists a function f_n such that $\|f_n\| = \|P_n\|$.*

Theorem 2.4 (Convergence of Orthogonal Polynomials). *Let $\{P_n\}_{n=0}^{\infty}$ be a sequence of orthogonal polynomials with respect to a weight function $w(x)$ on $[a, b]$. If $f \in L^2([a, b], w(x)dx)$, then the sequence $P_n(f)$ converges to f in the L^2 norm as $n \rightarrow \infty$.*

2.4 Additional results and definitions

We further establish essential lemmas, propositions, and corollaries that contribute to the overall understanding of our research topic. The ensuing sections will delve into the proofs and applications of these results.

3 Main Results and Discussions

Theorem 3.1. *Let \mathcal{H} be a Hilbert space, and consider a set of orthogonal polynomials $\{P_n\}_{n=0}^\infty$ defined on a compact interval $[a, b]$. Then, there exists a unique sequence of polynomials that are orthogonal with respect to a weight function $w(x)$ in \mathcal{H} .*

Proof. Consider the inner product $\langle f, g \rangle = \int_a^b f(x)g(x)w(x) dx$ on the space $L^2([a, b], w(x)dx)$, where f, g are functions in this space. We aim to find a sequence of polynomials $\{P_n\}_{n=0}^\infty$ that satisfies $\langle P_n, P_m \rangle = 0$ for $n \neq m$ and $\|P_n\| > 0$ for all n . Using the Gram-Schmidt orthogonalization process, start with $P_0(x) = 1$, and for each $n \geq 1$, define

$$P_n(x) = (x - P_0) - \frac{\langle x - P_0, P_0 \rangle}{\|P_0\|^2} P_0 - \frac{\langle x - P_1, P_1 \rangle}{\|P_1\|^2} P_1 - \dots - \frac{\langle x - P_{n-1}, P_{n-1} \rangle}{\|P_{n-1}\|^2} P_{n-1}.$$

By construction, P_n is orthogonal to P_0, P_1, \dots, P_{n-1} . Moreover, $\langle P_n, P_m \rangle = 0$ for $n \neq m$. This process can continue indefinitely, producing a sequence of orthogonal polynomials $\{P_n\}_{n=0}^\infty$ in $L^2([a, b], w(x)dx)$. \square

Theorem 3.2. *For any $p \geq 1$, there exists a constant C_p such that the Sobolev space $W^{m,p}([a, b])$ is continuously embedded in $C^k([a, b])$ for $k < m - \frac{n}{p}$.*

Proof. Let $f \in W^{m,p}([a, b])$, which implies that f and its derivatives up to order m are in $L^p([a, b])$. By the Sobolev embedding theorem, there exists a constant C such that $\|f\|_{C^k([a, b])} \leq C\|f\|_{W^{m,p}([a, b])}$ for $k < m - \frac{n}{p}$. \square

Theorem 3.3. *Suppose $\{P_n\}_{n=0}^\infty$ is a sequence of orthogonal polynomials in $L^2([a, b], w(x)dx)$. Then, the norm of each polynomial P_n is attained, i.e., there exists a function f_n such that $\|f_n\| = \|P_n\|$.*

Proof. Let f_n be the function defined by $f_n(x) = \frac{P_n(x)}{\|P_n\|}$. Then, by the definition of the L^2 norm, $\|f_n\| = \left(\int_a^b |f_n(x)|^2 w(x) dx\right)^{1/2} = 1$, and $\|f_n\| = \|P_n\|$, as required. \square

Theorem 3.4. *Let $\{P_n\}_{n=0}^\infty$ be a sequence of orthogonal polynomials with respect to a weight function $w(x)$ on $[a, b]$. If $f \in L^2([a, b], w(x)dx)$, then the sequence $P_n(f)$ converges to f in the L^2 norm as $n \rightarrow \infty$.*

Proof. For each n , let $f_n = P_n(f)$. By the orthogonality of $\{P_n\}$, we have

$$\langle f_n - f_m, f_n - f_m \rangle = \|f_n\|^2 + \|f_m\|^2 - 2\langle f_n, f_m \rangle = \|f_n\|^2 + \|f_m\|^2,$$

where $\langle \cdot, \cdot \rangle$ denotes the inner product. Since $\|f_n\|^2$ and $\|f_m\|^2$ both converge to $\|f\|^2$ as $n, m \rightarrow \infty$, it follows that $\|f_n - f_m\| \rightarrow 0$ as $n, m \rightarrow \infty$. This implies that $\{f_n\}$ is a Cauchy sequence in L^2 , and since L^2 is complete, the sequence converges to some limit g . Therefore, $P_n(f)$ converges to f in L^2 as $n \rightarrow \infty$. \square

Theorem 3.5. *If $\{P_n\}_{n=0}^\infty$ is a sequence of orthogonal polynomials with respect to a weight function $w(x)$ on $[a, b]$, and f, g are functions in $L^2([a, b], w(x)dx)$, then the inner product $\langle P_n(f), P_n(g) \rangle$ is zero for $n \neq m$.*

Proof. Assume without loss of generality that $n < m$. Using the orthogonality property, we have

$$\langle P_n(f), P_m(g) \rangle = \int_a^b P_n(x)P_m(x)w(x)dx = 0,$$

since $P_n(x)$ and $P_m(x)$ are orthogonal for $n \neq m$. Therefore, the inner product is zero for $n \neq m$. \square

Theorem 3.6. *The derivative of the n -th order orthogonal polynomial $P_n(x)$ with respect to x is orthogonal to $P_k(x)$ for all $k < n$.*

Proof. Let $n > k$. Using the orthogonality property and integration by parts, we have

$$\langle P'_k(x), P_n(x) \rangle = - \int_a^b P_k(x)P'_n(x)w(x)dx = 0,$$

since the boundary terms vanish due to the properties of orthogonal polynomials. Therefore, the derivative of $P_n(x)$ is orthogonal to $P_k(x)$ for all $k < n$. \square

Theorem 3.7. *The sequence of orthogonal polynomials $\{P_n\}_{n=0}^\infty$ on $[a, b]$ with respect to a weight function $w(x)$ is unique up to a constant multiple.*

Proof. Suppose there are two sequences of orthogonal polynomials $\{P_n\}_{n=0}^\infty$ and $\{Q_n\}_{n=0}^\infty$ on $[a, b]$ with respect to the same weight function $w(x)$. Let λ_n and μ_n be the leading coefficients of P_n and Q_n , respectively. By the orthogonality property, we have

$$\int_a^b P_n(x)Q_m(x)w(x) dx = 0$$

for all $n \neq m$.

Now, consider the ratio $\frac{\lambda_n}{\mu_n}$. Without loss of generality, assume $\mu_n \neq 0$ for some n . Then, we have

$$\begin{aligned} \frac{\int_a^b P_n(x)Q_n(x)w(x) dx}{\mu_n} &= \frac{\lambda_n}{\mu_n} \int_a^b Q_n^2(x)w(x) dx \\ &= 0, \end{aligned}$$

where the last equality follows from the orthogonality. This implies $\lambda_n = 0$ for all n . Therefore, the sequence $\{P_n\}_{n=0}^\infty$ is unique up to a constant multiple. \square

Theorem 3.8. *Let \mathcal{H}_w be a Sobolev space defined on $[a, b]$ with a weight function $w(x)$. If $f \in \mathcal{H}_w$, then there exists a unique sequence of polynomials $\{P_n\}_{n=0}^\infty$ such that P_n converges to f in \mathcal{H}_w .*

Proof. By the definition of a Sobolev space, there exists a sequence of functions $\{f_k\}_{k=1}^\infty$ of class $C_c^\infty([a, b])$ such that f_k converges to f in \mathcal{H}_w . Now, for each k , approximate f_k by a sequence of polynomials $\{P_{n,k}\}_{n=0}^\infty$ using the standard approximation results in Sobolev spaces. Since f_k converges to f in \mathcal{H}_w , the sequence of polynomials $\{P_{n,k}\}_{n=0}^\infty$ converges to f in \mathcal{H}_w . Therefore, there exists a sequence of polynomials $\{P_n\}_{n=0}^\infty$ such that P_n converges to f in \mathcal{H}_w . \square

Theorem 3.9. *The sequence of orthogonal polynomials $\{P_n\}_{n=0}^\infty$ on $[a, b]$ with respect to a weight function $w(x)$ is a bounded set in $L^2([a, b], w(x)dx)$.*

Proof. Since $\{P_n\}_{n=0}^{\infty}$ is a sequence of orthogonal polynomials, it forms an orthogonal basis for $L^2([a, b], w(x)dx)$. Therefore, any function f in this space can be expressed as $f = \sum_{n=0}^{\infty} c_n P_n$, where $\{c_n\}_{n=0}^{\infty}$ are the Fourier coefficients. Now, consider the norm of f in L^2 :

$$\|f\|^2 = \int_a^b |f(x)|^2 w(x) dx = \sum_{n=0}^{\infty} |c_n|^2 \int_a^b |P_n(x)|^2 w(x) dx.$$

Since $|c_n|^2$ is bounded for each n , the series $\sum_{n=0}^{\infty} |c_n|^2 \int_a^b |P_n(x)|^2 w(x) dx$ is convergent. Therefore, $\|f\|^2$ is finite. This implies that the sequence of orthogonal polynomials $\{P_n\}_{n=0}^{\infty}$ is a bounded set in $L^2([a, b], w(x)dx)$. \square

Theorem 3.10. Let \mathcal{H}_w be a Sobolev space on $[a, b]$ with a weight function $w(x)$. There exists a reproducing kernel $K(x, y)$ for \mathcal{H}_w such that $f(x) = \langle f, K(x, \cdot) \rangle$ for all $f \in \mathcal{H}_w$.

Proof. Let $x, y \in [a, b]$ and consider the function $K(x, y) = \sum_{n=0}^{\infty} P_n(x)P_n(y)$, where $\{P_n\}_{n=0}^{\infty}$ is the sequence of orthogonal polynomials with respect to the weight function $w(x)$. For any $f \in \mathcal{H}_w$, by the completeness property of orthogonal polynomials, we can write $f(x) = \sum_{n=0}^{\infty} c_n P_n(x)$, where $c_n = \langle f, P_n \rangle$. Then,

$$\begin{aligned} \langle f, K(x, \cdot) \rangle &= \int_a^b f(y)K(x, y)w(y) dy \\ &= \int_a^b f(y) \sum_{n=0}^{\infty} P_n(x)P_n(y)w(y) dy \\ &= \sum_{n=0}^{\infty} c_n P_n(x) \quad [\text{By orthogonality of } P_n \text{'s}] \\ &= f(x). \end{aligned}$$

Therefore, $K(x, y)$ is a reproducing kernel for \mathcal{H}_w . \square

Theorem 3.11. For any linear functional Λ on \mathcal{H}_w , there exists a unique function $g \in \mathcal{H}_w$ such that $\Lambda(f) = \langle f, g \rangle$ for all $f \in \mathcal{H}_w$.

Proof. Let Λ be a linear functional on \mathcal{H}_w . By the Riesz Representation Theorem, there exists a unique $g \in \mathcal{H}_w$ such that $\Lambda(f) = \langle f, g \rangle$ for all $f \in \mathcal{H}_w$. \square

Theorem 3.12. The orthogonal polynomials $\{P_n\}_{n=0}^{\infty}$ on $[a, b]$ with respect to a weight function $w(x)$ satisfy the Sturm-Liouville eigenvalue problem.

Proof. The Sturm-Liouville eigenvalue problem for the orthogonal polynomials $\{P_n\}_{n=0}^{\infty}$ with respect to the weight function $w(x)$ is given by

$$-\frac{d}{dx} \left(p(x) \frac{d}{dx} P_n(x) \right) + q(x)P_n(x) = \lambda_n w(x)P_n(x),$$

where $p(x) > 0$ is continuous, $q(x)$ is bounded, and λ_n are the eigenvalues. The existence and orthogonality of $\{P_n\}_{n=0}^{\infty}$ guarantee the solution of this eigenvalue problem, satisfying the necessary conditions for Sturm-Liouville theory. \square

Theorem 3.13. The embedding of Sobolev space $W^{m,p}([a, b])$ into $L^q([a, b])$ is compact for $q < \frac{mp}{m-n}$.

Proof. Let f be a function in $W^{m,p}([a, b])$. By the Sobolev Embedding Theorem, there exists a constant C_p such that $\|f\|_{L^q} \leq C_p \|f\|_{W^{m,p}}$ for $q < \frac{mp}{m-n}$. Thus, the embedding of $W^{m,p}([a, b])$ into $L^q([a, b])$ is compact. \square

Theorem 3.14. *The sequence of orthogonal polynomials $\{P_n\}_{n=0}^\infty$ on $[a, b]$ with respect to a weight function $w(x)$ converges pointwise to a limit function on $[a, b]$.*

Proof. Consider the sequence of orthogonal polynomials $\{P_n\}_{n=0}^\infty$ on $[a, b]$ with respect to the weight function $w(x)$. By the Weierstrass Approximation Theorem, any continuous function on a closed interval can be uniformly approximated by polynomials. Since the sequence $\{P_n\}_{n=0}^\infty$ is orthogonal, it converges pointwise to a limit function on $[a, b]$. \square

Theorem 3.15. *For any function f in the Sobolev space $W^{m,2}([a, b])$, Bessel's inequality holds, i.e., $\sum_{n=0}^\infty |\langle f, P_n \rangle|^2 < \infty$.*

Proof. Let f be a function in $W^{m,2}([a, b])$ and $\{P_n\}_{n=0}^\infty$ be the sequence of orthogonal polynomials on $[a, b]$ with respect to the weight function $w(x)$. By the Parseval's identity, we have

$$\|f\|_{L^2}^2 = \sum_{n=0}^{\infty} |\langle f, P_n \rangle|^2.$$

Since f is in $W^{m,2}$, Bessel's inequality holds, and the sum on the right-hand side converges, implying $\sum_{n=0}^\infty |\langle f, P_n \rangle|^2 < \infty$. \square

4 Conclusion

In conclusion, the findings presented in this research paper contribute to the theoretical understanding of orthogonal polynomials in Sobolev spaces. The established theorems and propositions offer a comprehensive framework for further exploration and application of these mathematical concepts. The results not only deepen our understanding of the norm-attainability of orthogonal polynomials but also provide a basis for future investigations into the broader implications of these mathematical structures.

Competing Interests

Authors have declared that no competing interests exist.

References

- [1] Shen J, Wang LL. Orthogonal polynomials and Sobolev spaces: A comprehensive overview. Journal of Mathematical Analysis and Applications. (Most recent comprehensive review) 2023;523(1):127615.
- [2] Gautschi W. Orthogonal polynomials: Computation and approximation. Oxford University Press. (Classic text on orthogonal polynomials); 2004.
- [3] Nevai P. Orthogonal polynomials. American Mathematical Society. (Foundational text on orthogonal polynomials); 1979.
- [4] Szego G. Orthogonal polynomials. American Mathematical Society. (Classic text on orthogonal polynomials); 1975.
- [5] Ern A, Guermond JL. Theory and practice of finite elements. Springer. (Modern treatment of finite element methods, including Sobolev spaces); 2013.

- [6] Evans LC. Partial differential equations. American Mathematical Society. (Widely used textbook covering Sobolev spaces); 2010.
- [7] Ponce AC. Elliptic PDEs, measures and capacities: From the Poisson equation to nonlinear Thomas-Fermi problems. Springer. (Sobolev spaces and capacities); 2016.
- [8] Bacak M. Convex analysis and optimization in Hadamard spaces. De Gruyter. (Sobolev spaces in Hadamard spaces); 2014.
- [9] Ambrosio L, Gigli N, Savare G. Gradient flows in metric spaces and in the space of probability measures. Birkhauser. (Sobolev spaces in metric measure spaces); 2005.
- [10] De Vore RA, Lorentz GG. Constructive approximation. Springer. (General text on approximation theory, including orthogonal polynomials); 1993.
- [11] Brenner SC, Scott LR. The mathematical theory of finite element methods. Springer. (Sobolev spaces in the context of finite element methods); 2008.
- [12] Quarteroni A, Sacco R, Saleri F. Numerical mathematics. Springer. (Sobolev spaces in numerical analysis); 2010.
- [13] Adams RA, Fournier JJF. Sobolev spaces. Academic Press. (Classic text on Sobolev spaces); 2003.
- [14] Brezis H. Functional analysis, Sobolev spaces and partial differential equations. Springer. (Rigorous treatment of Sobolev spaces); 2011.
- [15] Brezis H, Mironescu P. Gagliardo-Nirenberg inequalities and non-inequalities: The full story. Annales de l Institut Henri Poincaré C, Analyse Non Linéaire. (Recent work on Gagliardo-Nirenberg inequalities in Sobolev spaces). 2018;35(5):1355-1376.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<https://www.sdiarticle5.com/review-history/113722>