

Research Article

Quantum Influence of Topological Defects on a Relativistic Scalar Particle with Cornell-Type Potential in Cosmic String Space-Time with a Spacelike Dislocation

Faizuddin Ahmed 

Ajmal College of Arts and Science, Dhubri, 783324 Assam, India

Correspondence should be addressed to Faizuddin Ahmed; faizuddinahmed15@gmail.com

Received 22 April 2020; Revised 23 May 2020; Accepted 1 June 2020; Published 19 June 2020

Academic Editor: Edward Sarkisyan-Grinbaum

Copyright © 2020 Faizuddin Ahmed. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We study the relativistic quantum of scalar particles in the cosmic string space-time with a screw dislocation (torsion) subject to a uniform magnetic field including the magnetic quantum flux in the presence of potential. We solve the Klein-Gordon equation with a Cornell-type scalar potential in the considered framework and obtain the energy eigenvalues and eigenfunctions and analyze a relativistic analogue of the Aharonov-Bohm effect for bound states.

1. Introduction

In relativistic quantum mechanics, the study of spin-0 scalar particles via the Klein-Gordon equation on curved the background with the cosmic string has been of current research interest. Several authors have investigated the physical properties of a series of background with Gödel-type geometries, such as the relativistic quantum dynamics of a scalar particle [1, 2], spin-0 massive charged particles in the presence of a uniform magnetic field with the cosmic string [3], quantum influence of topological defects [4], linear confinement of a scalar particle with the cosmic string [5–7]. Furthermore, the relativistic quantum dynamics of spin-0 scalar particles was investigated in [8, 9] and observed the similarity of the energy eigenvalues with the Landau levels [10]. The relativistic quantum dynamics of a scalar particle in the presence of a homogeneous magnetic field within the Kaluza-Klein theory was investigated in [11]. The relativistic quantum dynamics of spin-0 massive charged particles in a four-dimensional curved space-time with the cosmic string was studied in [12]. Survey on the Klein-Gordon equation in the Gödeltype space-times was studied in [13]. Furthermore, spin-0 scalar particle [14], Klein-Gordon oscillator [15], generalized Klein-Gordon oscillator subject to a

Coulomb-type scalar potential [16], linear confinement of scalar particle [17], quantum effects on spin-0 charged particles with a Coulomb-type scalar, and vector potentials [18] in $(1+2)$ -dimensional rotational symmetry space-time backgrounds have been investigated. In addition, the relativistic quantum dynamics of spin-half particles have also been investigated (e.g., [1, 19–22]). The Dirac equation in $(1+2)$ -dimensional rotational symmetry space-time was investigated in [23].

The cosmic string space-time in the polar coordinates (t, r, φ, z) is described by the following line element [19, 24–29]:

$$ds^2 = -dt^2 + dr^2 + \alpha^2 r^2 d\varphi^2 + dz^2, \quad (1)$$

where $\alpha = 1 - 4\mu$ is the topological parameter with μ being the linear mass density of the cosmic string. In cosmic string space-time, the parameter μ assumes values in the interval $0 < \mu < 1$ within the general relativity [30, 31]. Furthermore, in the cylindrical symmetry, we have that $0 < r < \infty$, $0 \leq \varphi \leq 2\pi$, and $-\infty < z < \infty$. Cosmic string may have been produced by phase transitions in the early universe [32] as it is predicted in the extensions of the standard model

of particle physics [24, 25]. Several authors have studied the relativistic quantum mechanics in the cosmic string space-time (e.g., [19, 33–48]).

Various potentials have been used to investigate the bound state solutions to the relativistic wave-equations. Among them, much attention has given on the Cornell potential. The Cornell potential, which consists of a linear potential plus a Coulomb potential, is a particular case of the quark-antiquark interaction, one more harmonic type term [49]. The Coulomb potential is responsible for the interaction at small distances and linear potential leads to the confinement. Recently, the Cornell potential has studied in the ground state of three quarks [50]. However, this type of potential is worked on spherical symmetry; in cylindrical symmetry, which is our case, this type of potential is known as Cornell-type potential [3]. Investigation of the relativistic wave-equations with this type of potential is the Klein-Gordon scalar field in spinning cosmic string space-time [48], relativistic quantum dynamics of scalar particle subject to a uniform magnetic field in cosmic string spacetime [19], spin-0 scalar particle in $(1+2)$ -dimensional rotational symmetry space-time [51], Aharonov-Bohm effect for bound states [52], quantum effects of confining potential on the Klein-Gordon oscillator [53], effects of potential on a position-dependent mass system [54], etc. Other investigations with this type of interaction are in [55–58]. The Cornell-type potential is given by

$$S = \frac{\eta_c}{r} + \eta_L r, \quad (2)$$

where η_c, η_L are the potential parameters.

In [19], authors studied the relativistic quantum dynamics of bosonic charged particles in the presence of external fields in a cosmic string space-time. They solved the Klein-Gordon equation and obtained the relativistic energy eigenvalues and wave-function. In addition, they introduced a Cornell-type scalar potential by modifying the mass term in the Klein-Gordon equation and obtained the bound state solution of the relativistic quantum system. In [59], authors studied a spin-0 scalar massive charged particle in the presence of external fields including a magnetic quantum flux in the space-time with a spacelike dislocation under the influence of linear potential. They solved the Klein-Gordon equation and evaluated the energy eigenvalues and analyze a relativistic analogue of the Aharonov-Bohm effect for bound states. In addition, they introduced a linear scalar potential by modifying the mass term in the Klein-Gordon equation and obtained the bound state solution of the relativistic quantum system. In [60], authors investigated a spin-0 scalar charged particles in the presence of external fields including a magnetic quantum flux in the space-time with a spacelike dislocation subject to a Coulomb-type potential. They solved the Klein-Gordon equation and evaluated the bound state solution of the relativistic quantum system and analyze a relativistic analogue of the Aharonov-Bohm effect for bound states.

Our main motivation in this work is to investigate a relativistic analogue of the Aharonov-Bohm effect [61, 62] for bound states of a relativistic scalar charged particle subject to a homogeneous magnetic field including a magnetic quantum flux in the presence of a Cornell-type potential in the cosmic string space-time with a spacelike dislocation. We solve the Klein-Gordon equation in the considered framework and obtain the relativistic energy eigenvalues and eigenfunctions and analyze the effects on the eigenvalues. In addition, we check the role play by the torsion parameter in this relativistic system and see that the presence of the torsion parameter modifies the energy levels and breaks their degeneracy in comparison to the result obtained in the cosmic string space-time case.

2. Bosonic Charged Particles: The KG-Equation

In [63, 64], examples of topological defects in the space-time associated with torsion are given. We start this section by considering the cosmic string space-time with a spacelike dislocation, whose line element is given by $(x^0 = t, x^1 = r, x^2 = \varphi, x^3 = z)$

$$ds^2 = -dt^2 + dr^2 + \alpha^2 r^2 d\varphi^2 + (dz + \chi d\varphi)^2, \quad (3)$$

where $\alpha > 0$ is the cosmic string, and χ is the dislocation (torsion) parameter. For zero torsion parameter, $\chi \rightarrow 0$, the metric (3) reduces to the cosmic string space-time. Furthermore, for $\chi \rightarrow 0$ and $\alpha \rightarrow 1$, the study space-time reduces to Minkowski flat space metric in cylindrical coordinates. Topological defects associated with torsion have investigated in solid state [30, 65–68], quantum scattering [69], bound states solutions [52, 59, 70], and in relativistic quantum mechanics [71, 72].

The metric tensor for the space-time (3) to be

$$g_{\mu\nu}(x) = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \alpha^2 r^2 + \chi^2 & \chi \\ 0 & 0 & \chi & 1 \end{pmatrix} \quad (4)$$

with its inverse

$$g^{\mu\nu}(x) = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \frac{1}{\alpha^2 r^2} & -\frac{\chi}{\alpha^2 r^2} \\ 0 & 0 & -\frac{\chi}{\alpha^2 r^2} & 1 + \frac{\chi^2}{\alpha^2 r^2} \end{pmatrix} \quad (5)$$

The metric has a signature $(-, +, +, +)$, and the determinant of the corresponding metric tensor $g_{\mu\nu}$ is

$$\det g = -\alpha^2 r^2. \quad (6)$$

The relativistic quantum dynamics of spin-0 charged scalar particles of mass m is described by the Klein-Gordon (KG) equation [19, 73].

$$\left[\frac{1}{\sqrt{-g}} D_\mu (\sqrt{-g} g^{\mu\nu} D_\nu) - m^2 \right] \Psi = 0, \quad (7)$$

where the minimal substitution is defined by

$$D_\mu \equiv \partial_\mu - i e A_\mu, \quad (8)$$

where e is the electric charge, and A_μ is the electromagnetic four-vector potential by

$$A_\mu = (0, \vec{A}), \quad \vec{A} = (0, A_\phi, 0). \quad (9)$$

We choose the angular component of electromagnetic four-vector potential [52, 55, 57–60, 71, 74].

$$A_\phi = -\frac{1}{2} \alpha B_0 r^2 + \frac{\Phi_B}{2\pi}, \quad \vec{B} = \vec{\nabla} \times \vec{A} = -B_0 \hat{k}. \quad (10)$$

Here, $\Phi_B = \text{const.}$ is the internal quantum magnetic flux [75, 76] through the core of the topological defects [77]. It is noteworthy that the Aharonov-Bohm effect [61, 62] has been investigated in several branches of physics, such as in, graphene [78], Newtonian theory [79], bound states of massive fermions [80], scattering of dislocated wave fronts [81], torsion effects on a relativistic position-dependent mass system [52, 59], Kaluza-Klein theory [11, 54, 82–86], and nonminimal Lorentz-violating coupling [87].

If one introduces a scalar potential by modifying the mass term in the form $m \rightarrow m + S(r)$ [88] into the above equation, then we have

$$\left[\frac{1}{\sqrt{-g}} D_\mu (\sqrt{-g} g^{\mu\nu} D_\nu) - (m + S)^2 \right] \Psi = 0, \quad (11)$$

Several authors have studied the relativistic wave-equations with various kinds of potentials such as linear, Coulomb-type, and Cornell-type (e.g., [3, 5, 7, 19, 51–53, 56, 57, 59, 60]).

Using the equation (3), Eq. (11) becomes

$$\left[-\frac{\partial^2}{\partial t^2} + \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial}{\partial r} \right) + \frac{1}{\alpha^2 r^2} \left(\frac{\partial}{\partial \phi} - i e A_\phi - \chi \frac{\partial}{\partial z} \right)^2 + \frac{\partial^2}{\partial z^2} - (m + S)^2 \right] \Psi = 0. \quad (12)$$

Since the line-element (3) is independent of t, ϕ, z , it is appropriate to choose the following ansatz for the function Ψ

$$\Psi(t, r, \phi, z) = e^{i(-Et + l\phi + kz)} \psi(r), \quad (13)$$

where E is the energy of charged particle, $l = 0, \pm 1, \pm 2, \dots \in Z$ is the eigenvalues of z -component of the angular momentum operator, and k is a constant.

Substituting Eq. (13) into the Eq. (12), we obtain the following radial wave-equation for $\psi(r)$:

$$\left[\frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} + E^2 - \frac{1}{\alpha^2 r^2} (l - e A_\phi - k \chi)^2 - k^2 - (m + S)^2 \right] \psi(r) = 0. \quad (14)$$

Substituting the Eq. (10) and scalar potential (2) into the Eq. (14), we obtain

$$\left[\frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} + \lambda - \omega^2 r^2 - \frac{j^2}{r^2} - \frac{a}{r} - br \right] \psi(r) = 0, \quad (15)$$

where

$$\begin{aligned} \lambda &= E^2 - m^2 - k^2 - 2\eta_c \eta_L - \frac{2m\omega_c}{\alpha} (l - k\chi - \Phi), \\ \omega &= \sqrt{m^2 \omega_c^2 + \eta_L^2}, \\ j &= \sqrt{\frac{(l - k\chi - \Phi)^2}{\alpha^2} + \eta_c^2}, \\ a &= 2m\eta_c, \\ b &= 2m\eta_L, \\ \omega_c &= \frac{eB_0}{2m} \end{aligned} \quad (16)$$

is called the cyclotron frequency of the particle moving in the magnetic field.

Transforming $x = \sqrt{\omega} r$ into the Eq. (15), we obtain the following equation:

$$\psi''(x) + \frac{1}{x} \psi'(x) + \left[\zeta - x^2 - \frac{j^2}{x^2} - \frac{\eta}{x} - \theta x \right] \psi(x) = 0, \quad (17)$$

where we have defined

$$\zeta = \frac{\lambda}{\omega}, \quad \eta = \frac{a}{\sqrt{\omega}}, \quad \theta = \frac{b}{\omega^{3/2}}. \quad (18)$$

We now use appropriate boundary conditions to investigate the bound state solutions in this problem. It is required that the wave-functions must be regular both at $x \rightarrow 0$ and $x \rightarrow \infty$. Suppose the possible solution to the Eq. (17) is

$$\psi(x) = x^j e^{-\frac{1}{2}(\theta+x)x} H(x), \quad (19)$$

where $H(x)$ is an unknown function. Substituting the solution (19) into the Eq. (17), we obtain

$$H''(x) + \left[\frac{\gamma}{x} - \theta - 2x \right] H'(x) + \left[-\frac{\beta}{x} + \Theta \right] H(x) = 0, \quad (20)$$

where

$$\begin{aligned} \gamma &= 1 + 2j, \\ \Theta &= \zeta + \frac{\theta^2}{4} - 2(1+j), \\ \beta &= \eta + \frac{\theta}{2}(1+2j). \end{aligned} \quad (21)$$

Equation (20) is the biconfluent Heun's differential equation [3, 5, 7, 12, 19, 48, 51–54, 57, 59, 71, 74, 85, 86, 89, 90] with $H(x)$ is the Heun polynomial function.

The above equation (20) can be solved by the Frobenius method. Writing the solution as a power series expansion around the origin [91]:

$$H(x) = \sum_{i=0}^{\infty} c_i x^i. \quad (22)$$

Substituting the power series solution (22) into the Eq. (20), we get the following recurrence relation for the coefficients:

$$c_{n+2} = \frac{1}{(n+2)(n+2+2j)} [\{\beta + \theta(n+1)\} c_{n+1} - (\Theta - 2n) c_n]. \quad (23)$$

And the various coefficients are

$$\begin{aligned} c_1 &= \left(\frac{\eta}{1+2j} + \frac{\theta}{2} \right) c_0, \\ c_2 &= \frac{1}{4(1+j)} [(\beta + \theta) c_1 - \Theta c_0]. \end{aligned} \quad (24)$$

As the function $H(x)$ has a power series expansion around the origin in Eq. (22), then, the relativistic bound states solution can be achieved by imposing that the power series expansion becomes a polynomial of degree n . Through the recurrence relation Eq. (23), we can see that the power series expansion $H(x)$ becomes a polynomial of degree n by imposing the following two conditions [3, 5, 7, 12, 19, 48, 51–54, 57, 59, 71, 74, 85, 86, 92, 93].

$$\begin{aligned} \Theta &= 2n, \quad (n = 1, 2, \dots) \\ c_{n+1} &= 0. \end{aligned} \quad (25)$$

By analyzing the first condition $\Theta = 2n$, we get the second-degree equation of the energy eigenvalues $E_{n,l}$:

$$\begin{aligned} E_{n,l}^2 &= \frac{2m\omega_c}{\alpha} (l - k\chi - \Phi) + 2\omega \left(n + 1 + \sqrt{\frac{(l - k\chi - \Phi)^2}{\alpha^2} + \eta_c^2} \right) \\ &\quad - \frac{m^2 \eta_L^2}{\omega^2} + m^2 + k^2 + 2\eta_c \eta_L. \end{aligned} \quad (26)$$

The wave-functions are given by

$$\psi_{n,l}(x) = x \sqrt{\frac{(l - k\chi - \Phi)^2}{\alpha^2} + \eta_c^2} e^{-\frac{1}{2} \left(\frac{2m\eta_L}{\omega^2} + x \right) x} H(x). \quad (27)$$

Note that the Eq. (26) does not represent the general expression for eigenvalues problem. One can obtain the individual energy eigenvalues one by one, that is, E_1, E_2, E_3 by imposing the additional recurrence condition $c_{n+1} = 0$ on the eigenvalue. The solution with Heun's Equation makes it possible to obtain the eigenvalues one by one as done in [3, 5, 7, 12, 19, 48, 51–54, 57, 59, 71, 74, 85, 86, 92, 93] but not explicitly in the general form by all eigenvalues n . With the aim of obtaining the energy levels of the stationary states, let us discuss a particular case of $n = 1$. This means that we want to construct a polynomial of the first degree to $H(x)$. With $n = 1$, we have $\Theta = 2$ and $c_2 = 0$ which implies from Eq. (24)

$$\begin{aligned} c_1 &= \frac{2}{(\beta + \theta)} c_0 \Rightarrow \frac{\eta}{1+2j} + \frac{\theta}{2} = \frac{2}{(\beta + \theta)} \Rightarrow \omega_{1,l}^3 \\ &\quad - \frac{a^2}{2(1+2j)} \omega_{1,l}^2 - a b \left(\frac{1+j}{1+2j} \right) \omega_{1,l} - \frac{b^2}{8} (3+2j) = 0. \end{aligned} \quad (28)$$

a constraint on the physical parameter $\omega_{1,l}$. The relation given in Eq. (28) gives the possible values of the parameter $\omega_{1,l}$ that permit us to construct first degree polynomial to $H(x)$ for $n = 1$. Note that its values change for each quantum number n and l , so we have labeled $\omega \rightarrow \omega_{n,l}$. Besides, since this parameter is determined by the frequency or the magnetic field B_0 , hence, the magnetic field $B_0^{1,l}$ is so adjusted that the Eq. (28) can be satisfied and the first degree polynomial to $H(x)$ can be achieved, where we have simplified our notation by labeling:

$$\omega_{c1,l} = \frac{1}{m} \sqrt{\omega_{1,l}^2 - \eta_L^2} \leftrightarrow B_0^{1,l} = \frac{2}{e} \sqrt{\omega_{1,l}^2 - \eta_L^2}. \quad (29)$$

It is noteworthy that a third-degree algebraic equation (28) has at least one real solution, and it is exactly this solution that gives us the allowed values of the magnetic field for the lowest state of the system, which we do not write because its expression is very long. We can note, from Eq. (29) that the possible values of the magnetic field depend on the quantum numbers and the potential parameter. In addition, for each relativistic energy level, we have different relation of the magnetic field associated to the Cornell-type potential and quantum numbers of the system $\{l, n\}$. For this reason, we have labeled the parameters ω_c , and B_0 in Eqs. (28) and (29).

Therefore, the ground state energy levels for $n = 1$ is given by

$$E_{1,l} = \pm \left\{ \frac{2m\omega_{c1,l}}{\alpha} (l - k\chi - \Phi) + 2m\omega_{1,l} \left(2 + \sqrt{\frac{(l - k\chi - \Phi)^2}{\alpha^2} + \eta_c^2} \right) - \frac{m^2 \eta_L^2}{\omega_{1,l}^2} + m^2 + k^2 + 2\eta_c \eta_L \right\}^{\frac{1}{2}}, \quad (30)$$

Then, by substituting the real solution $\omega_{1,l}$ from Eq. (28) into the Eq. (30), it is possible to obtain the allowed values of the relativistic energy levels for the radial mode $n = 1$ of a position dependent mass system. We can see that the lowest energy state is defined by the real solution of the algebraic equation Eq. (28) plus the expression given in Eq. (30) for the radial mode $n = 1$, instead of $n = 0$. This effect arises due to the presence of Cornell-type potential in the system. Note that it is necessary physically that the lowest energy state is $n = 1$ and not $n = 0$; otherwise, the opposite would imply that $c_1 = 0$, which requires that the rest mass of the scalar particle be zero that is contrary to the proposal of this investigation.

For $\Phi_B \neq 0$ and $\chi \neq 0$, we can observe in Eq. (30) there exists an effective angular momentum $l \rightarrow l' = (l/\alpha) (l - \Phi - k\chi)$. Thus, the relativistic energy levels depend on the geometric phase [75, 76] as well as the torsion parameter. This dependence of the energy levels on the geometric quantum phase gives rise to the well-known effect called as the Aharonov-Bohm effect for bound states [11, 52, 54, 59, 61, 62, 82–86]. Besides, we have that $E_{n,l}(\Phi_B + \Phi_0) = E_{n,l \mp \tau}(\Phi_B)$ where $\Phi_0 = \pm(2\pi/e)\tau$ with $\tau = 0, 1, 2, \dots$, which means the relativistic energy eigenvalues (30) is a periodic function of the Aharonov-Bohm geometric quantum phase.

The ground state wave function for $n = 1$ is given by

$$\psi_{1,l} = x \sqrt{\frac{(l - k\chi)^2}{\alpha^2} + \eta_c^2} e^{-\frac{1}{2} \left(\frac{2m\eta_L}{\omega_{1,l}^{3/2}} + x \right)} \times (c_0 + c_1 x), \quad (31)$$

where

$$c_1 = \frac{1}{\sqrt{\omega_{1,l}}} \left[\frac{2m\eta_c}{1+2j} + \frac{m\eta_L}{\omega_{1,l}} \right] c_0. \quad (32)$$

2.1. Interactions with Linear Scalar Potential. We discuss a case corresponds to $\eta_c \rightarrow 0$, that is, only a linear scalar potential in the considered relativistic quantum systems. Therefore, the radial wave-equation (15) becomes we obtain

$$\psi''(r) + \frac{1}{r} \psi'(r) + \left[\tilde{\lambda} - \omega^2 r^2 - \frac{j^2}{r^2} - br \right] \psi(r) = 0, \quad (33)$$

where $\tilde{\lambda} = E^2 - m^2 - k^2 - (2m\omega_c/\alpha)(l - k\chi - \Phi)$.

By changing the variable $x = \sqrt{\omega}r$, Eq. (33) becomes

$$\psi''(x) + \frac{1}{x} \psi'(x) + \left[\frac{\tilde{\lambda}}{\omega} - x^2 - \frac{j^2}{x^2} - \theta r \right] \psi(x) = 0, \quad (34)$$

Substituting the solution Eq. (19) into the Eq. (34), we obtain

$$H''(x) + \left[\frac{\gamma}{x} - \theta - 2x \right] H'(x) + \left[-\frac{\tilde{\beta}}{x} + \tilde{\Theta} \right] H(x) = 0, \quad (35)$$

where

$$\begin{aligned} \gamma &= 1 + 2j, \\ \tilde{\Theta} &= \frac{\tilde{\lambda}}{\omega} + \frac{\theta^2}{4} - 2(1+j), \\ \tilde{\beta} &= \frac{\theta}{2}(1+2j). \end{aligned} \quad (36)$$

Equation (35) is the biconfluent Heun's differential equation [3, 5, 7, 12, 19, 48, 51–54, 57, 59, 71, 74, 85, 86, 89, 90] with $H(x)$ is the Heun polynomials function.

Substituting the power series solution (22) into the Eq. (35), we get the following recurrence relation for the coefficients:

$$c_{n+2} = \frac{1}{(n+2)(n+2+2j)} \left[\left\{ \tilde{\beta} + \theta(n+1) \right\} c_{n+1} - \left(\tilde{\Theta} - 2n \right) c_n \right]. \quad (37)$$

And the various coefficients are

$$\begin{aligned} c_1 &= \frac{\theta}{2} c_0, \\ c_2 &= \frac{1}{4(1+j)} \left[\left(\tilde{\beta} + \theta \right) c_1 - \tilde{\Theta} c_0 \right]. \end{aligned} \quad (38)$$

The power series expansion $H(x)$ becomes a polynomial of degree n by imposing the following two conditions [3, 5, 7, 12, 19, 48, 51–54, 57, 59, 71, 74, 85, 86, 92, 93].

$$\begin{aligned} \tilde{\Theta} &= 2n, \quad (n = 1, 2, \dots) \\ c_{n+1} &= 0. \end{aligned} \quad (39)$$

By analyzing the first condition, we obtain the following energy eigenvalues expression $E_{n,l}$:

$$\begin{aligned} E_{n,l}^2 &= m^2 + k^2 + \frac{2m\omega_c}{\alpha} (l - k\chi - \Phi) \\ &+ 2\omega \left(n + 1 + \frac{|l - k\chi - \Phi|}{\alpha} \right) - \frac{m^2 \eta_L^2}{\omega^2}, \end{aligned} \quad (40)$$

where $n = 1, 2, \dots$.

The ground state energy levels associated with the radial mode $n = 1$ is given by

$$E_{1,l}^2 = m^2 + k^2 + \frac{2m\omega_{c1,l}}{\alpha} (l - k\chi - \Phi) + 2\omega_{1,l} \left(n + 1 + \frac{|l - k\chi - \Phi|}{\alpha} \right) - \frac{m^2 \eta_L^2}{\omega_{1,l}^2}, \quad (41)$$

where by using Eq. (39) for $n = 1$, we obtain the following constraint

$$\begin{aligned} \omega_{1,l} &= [m^2 \eta_L^2 (3 + 2j)]^{\frac{1}{3}}, \\ \omega_{c1,l} &= \frac{1}{m} \sqrt{[m^2 \eta_L^2 (3 + 2j)]^{\frac{2}{3}} - \eta_L^2}, \\ j &= \sqrt{\frac{(l - k\chi - \Phi)^2}{\alpha^2} + \eta_c^2}. \end{aligned} \quad (42)$$

Equation (41) is the ground state energy levels associated with the radial mode $n = 1$ of a relativistic scalar charged particle in the presence of an external uniform magnetic field including a magnetic quantum flux in the cosmic string space-time with a spacelike dislocation. For $\alpha \rightarrow 1$, the energy eigenvalues Eq. (40) reduce to the result obtained in [59]. We can see that the presence of cosmic string parameter (α) shifts the energy levels in comparison to those in [59]. Thus, by comparing the energy eigenvalues Eq. (30) with Eq. (41), we have the presence of an extra linear potential that modifies the relativistic spectrum of energy.

2.2. Interactions with Coulomb-Type Potential. We discuss another case corresponds to $\eta_L \rightarrow 0$, that is, only Coulomb-type scalar potential in the considered relativistic quantum systems. Therefore, the radial wave-equation (15) becomes we obtain

$$\psi''(r) + \frac{1}{r} \psi'(r) + \left[\tilde{\lambda} - m^2 \omega_c^2 r^2 - \frac{j^2}{r^2} - \frac{a}{r} \right] \psi(r) = 0, \quad (43)$$

where $\tilde{\lambda}$ is given earlier.

Let us define $x = \sqrt{m\omega_c} r$, then Eq. (43) becomes

$$\psi''(x) + \frac{1}{x} \psi'(x) + \left[\tilde{\lambda} - x^2 - \frac{j^2}{x^2} - \frac{\delta}{r} \right] \psi(x) = 0, \quad (44)$$

where $\delta = a/\sqrt{m\omega_c}$. By imposing that $\psi(x) \rightarrow 0$ when $x \rightarrow 0$ and $x \rightarrow \infty$, we have

$$\psi(x) = x^j e^{-\frac{x^2}{2}} H(x). \quad (45)$$

By substituting Eq. (45) into the Eq. (44), we obtain the following equation for $H(x)$:

$$H'(x) + \left[\frac{1+2j}{x} - 2x \right] H'(x) + \left[\frac{\tilde{\lambda}}{m\omega_c} - 2 - 2j - \frac{\delta}{x} \right] H(x) = 0. \quad (46)$$

Equation (46) is the biconfluent Heun's differential equation [3, 5, 7, 12, 19, 48, 51–54, 57, 59, 71, 74, 85, 86, 89, 90] with $H(x)$ is the Heun polynomials function.

Substituting the power series solution (22) into the Eq. (46), we get the following recurrence relation for the coefficients:

$$c_{n+2} = \frac{1}{(n+2)(n+2+2j)} \left[\delta c_{n+1} - \left(\frac{\tilde{\lambda}}{m\omega_c} - 2 - 2j - 2n \right) c_n \right]. \quad (47)$$

And the various coefficients are

$$\begin{aligned} c_1 &= \frac{\delta}{1+2j} c_0, \\ c_2 &= \frac{1}{4(1+j)} \left[\delta c_1 - \left(\frac{\tilde{\lambda}}{m\omega_c} - 2 - 2j \right) c_0 \right]. \end{aligned} \quad (48)$$

The power series expansion $H(x)$ becomes a polynomial of degree n by imposing the following two conditions [3, 5, 7, 12, 19, 48, 51–54, 57, 59, 71, 74, 85, 86, 92, 93].

$$\begin{aligned} \frac{\tilde{\lambda}}{m\omega_c} - 2 - 2j &= 2n, \quad (n = 1, 2, \dots) \\ c_{n+1} &= 0. \end{aligned} \quad (49)$$

By analyzing the first condition, we obtain the following equation of eigenvalues $E_{n,l}$:

$$E_{n,l} = \pm \sqrt{m^2 + k^2 + 2m\omega_c \left[n + 1 + j + \frac{1}{\alpha} (l - k\chi - \Phi) \right]} \quad (n = 1, 2, \dots). \quad (50)$$

For $n = 1$, the ground state energy levels are given by

$$E_{1,l} = \pm \sqrt{m^2 + k^2 + 2m\omega_{c1,l} \left[2 + j + \frac{1}{\alpha} (l - k\chi - \Phi) \right]}, \quad (51)$$

where by using Eq. (49) for $n = 1$, we obtain the following constraint

$$\omega_{c1,l} = \frac{2m\eta_c^2}{1+2j} \leftrightarrow B_0^{1,l} = \frac{4m^2\eta_c^2}{e(1+2j)}. \quad (52)$$

Here, the magnetic field $B_0^{1,l}$ is so adjusted that the Eq. (52) can be satisfied, and a polynomial of first degree to $H(x)$ can be achieved.

Equation (51) is the energy levels associated with the radial mode $n = 1$ of a relativistic scalar charged particle in the presence of an external uniform magnetic field including a magnetic quantum flux in the cosmic string space-time with a spacelike dislocation. For $\alpha \rightarrow 1$, the energy levels Eq. (51) reduces to the result obtained in [60]. Thus, we can see that the presence of the cosmic string parameter (α)

shifts the energy levels in comparison to those in [60]. Thus, by comparing the energy eigenvalue expression Eq. (30) with Eq. (51), we can see that the presence of an extra Coulomb-type potential modifies the relativistic energy spectrum of the system.

2.3. Without Torsion Parameter. We discuss here zero torsion parameter, $\chi \rightarrow 0$, in the considered relativistic quantum systems.

Therefore, the radial wave-equation (15) becomes

$$\left[\frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} + \lambda_0 - \omega^2 r^2 - \frac{j_0^2}{r^2} - \frac{a}{r} - br \right] \psi(r) = 0, \quad (53)$$

where

$$\begin{aligned} \lambda_0 &= E^2 - k^2 - m^2 - \frac{2m\omega_c}{\alpha} (l - \Phi) - 2\eta_c \eta_L, \\ j_0 &= \sqrt{\frac{(l - \Phi)^2}{\alpha^2} + \eta_c^2}. \end{aligned} \quad (54)$$

Transforming a new variable $x = \sqrt{\omega} r$ into the Eq. (53), we obtain

$$\left[\frac{d^2}{dx^2} + \frac{1}{x} \frac{d}{dx} + \frac{\lambda_0}{\omega} - x^2 - \frac{j_0^2}{x^2} - \frac{\eta}{x} - \theta x \right] \psi(r) = 0, \quad (55)$$

Suppose the possible solution to the Eq. (55) is

$$\psi(x) = x^{j_0} e^{-\frac{1}{2}(\theta+x)x} H(x). \quad (56)$$

Substituting the solution (19) into the Eq. (17), we obtain

$$\begin{aligned} H''(x) + \left[\frac{1+2j_0}{x} - \theta - 2x \right] H'(x) \\ + \left[-\frac{\beta_0}{x} + \frac{\lambda_0}{\omega} + \frac{\theta^2}{4} - 2 - 2j_0 \right] H(x) = 0, \end{aligned} \quad (57)$$

where $\beta_0 = \eta + (\theta/2)(1 + 2j_0)$.

Equation (57) is the biconfluent Heun's differential equation [3, 5, 7, 12, 19, 48, 51–54, 57, 59, 71, 74, 85, 86, 89, 90] with $H(x)$ is the Heun polynomial function.

Substituting the power series solution (22) into the Eq. (57), we get the following recurrence relation for the coefficients:

$$\begin{aligned} c_{n+2} &= \frac{1}{(n+2)(n+2+2j_0)} \left[\left\{ \eta + \theta \left(n + j_0 + \frac{3}{2} \right) \right\} c_{n+1} \right. \\ &\quad \left. - \left(\frac{\lambda_0}{\omega} + \frac{\theta^2}{4} - 2 - 2j_0 - 2n \right) c_n \right]. \end{aligned} \quad (58)$$

And the various coefficients are

$$\begin{aligned} c_1 &= \left(\frac{\eta}{1+2j_0} + \frac{\theta}{2} \right) c_0, \\ c_2 &= \frac{1}{4(1+j)} \left[\left\{ \eta + \theta \left(j_0 + \frac{3}{2} \right) \right\} c_1 - \left(\frac{\lambda_0}{\omega} + \frac{\theta^2}{4} - 2 - 2j_0 \right) c_0 \right]. \end{aligned} \quad (59)$$

The power series expansion $H(x)$ becomes a polynomial of degree n by imposing the following two conditions [3, 5, 7, 12, 19, 48, 51–54, 57, 59, 71, 74, 85, 86, 92, 93].

$$\begin{aligned} \frac{\lambda_0}{\omega} + \frac{\theta^2}{4} - 2 - 2j_0 &= 2n \quad (n = 1, 2, \dots) \\ c_{n+1} &= 0. \end{aligned} \quad (60)$$

By analyzing the first condition, we obtain the following second-degree energy eigenvalues expression $E_{n,l}$:

$$\begin{aligned} E_{n,l}^2 &= m^2 + k^2 + \frac{2m\omega_c(l-\Phi)}{\alpha} \\ &\quad + 2m\omega \left(n+1 + \sqrt{\frac{(l-\Phi)^2}{\alpha^2} + \eta_c^2} \right) \\ &\quad + 2\eta_c \eta_L - \frac{m^2 \eta_L^2}{\omega^2}. \end{aligned} \quad (61)$$

Equation (61) is the energy eigenvalues of a relativistic scalar particle with an external uniform magnetic field including a magnetic quantum flux in the cosmic string space-time subject to Cornell-type scalar potential. For zero magnetic quantum flux, $\Phi_B \rightarrow 0$, the energy eigenvalues (61) is consistent with the result in [19]. Thus, we can see that the energy eigenvalue expression Eq. (26) get modify in comparison to those in [19] due to the presence of the magnetic quantum flux Φ_B as well the torsion parameter χ which break the degeneracy of the energy spectrum.

To obtain the individual energy levels, we impose the additional recurrence condition $c_{n+1} = 0$. For example, $n = 1$, we have from (59)

$$\omega_{1,l}^3 - \frac{a^2}{2(1+2j_0)} \omega_{1,l}^2 - ab \left(\frac{1+j_0}{1+2j_0} \right) \omega_{1,l} - \frac{b^2}{8} (3+2j_0) = 0, \quad (62)$$

a constraint on the physical parameter $\omega_{1,l}$. The relation given in Eq. (62) gives the possible values of the parameter $\omega_{1,l}$ that permit us to construct first degree polynomial to $H(x)$ for $n = 1$. Note that its values change for each quantum number n and l , so we have labeled $\omega \rightarrow \omega_{n,l}$. Besides, since this parameter is determined by the frequency or the magnetic field B_0 , hence, the magnetic field $B_0^{1,l}$ is so adjusted that the Eq. (62) can be satisfied and the first degree polynomial to $H(x)$ can be achieved, where we have simplified our notation by labeling:

$$\omega_{c,1,l} = \frac{1}{m} \sqrt{\omega_{1,l}^2 - \eta_L^2} \leftrightarrow B_0^{1,l} = \frac{2}{e} \sqrt{\omega_{1,l}^2 - \eta_L^2}. \quad (63)$$

Note that the equation (62) has at least one real solution, and it is exactly this solution that gives us the allowed values of the magnetic field for the lowest state of the system, which we do not write because its expression is very long. We can note, from Eq. (63), that the possible values of the magnetic field depend on the quantum numbers and the potential parameter.

The ground state energy levels for $n = 1$ is

$$E_{1,l} = \pm \left\{ m^2 + k^2 + 2\eta_c \eta_L + \frac{2m\omega_{c,1,l}}{\alpha} (l - \Phi) + 2m\omega_{1,l} \left(2 + \sqrt{\frac{(l - \Phi)^2}{\alpha^2} + \eta_c^2} \right) - \frac{m^2 \eta_L^2}{\omega_{1,l}^2} \right\}^{\frac{1}{2}}, \quad (64)$$

Then, by substituting the real solution $\omega_{1,l}$ from Eq. (62) into the Eq. (64), it is possible to obtain the allowed values of the relativistic energy levels for the radial mode $n = 1$ of a position dependent mass system. We can see that the lowest energy state is defined by the real solution of the algebraic equation Eq. (62) plus the expression given in Eq. (64) for the radial mode $n = 1$, instead of $n = 0$. This effect arises due to the presence of Cornell-type potential in the system.

The ground state wave function is

$$\psi_{1,l} = x^{\sqrt{\frac{(l-\Phi)^2}{\alpha^2} + \eta_c^2}} e^{-\frac{1}{2} \left(\frac{2m\eta_L}{\omega_{1,l}^{3/2} + x} \right) x} (c_0 + c_1 x), \quad (65)$$

where

$$c_1 = \frac{1}{\omega_{1,l}^{1/2}} \left[\frac{2m\eta_c}{1 + 2j_0} + \frac{m\eta_L}{\omega_{1,l}} \right] c_0. \quad (66)$$

In subsection 2.1–2.3, we can see that the relativistic energy eigenvalues depend on the geometric quantum phase [75, 76]. This dependence of the energy eigenvalues on the geometric quantum phase gives rise to a relativistic analogue of the Aharonov-Bohm effect for bound states [11, 52, 54, 59, 61, 62, 82–86]. Besides, we have that $E_{n,l}(\Phi_B + \Phi_0) = E_{n,l \mp \tau}(\Phi_B)$ where $\Phi_0 = \pm(2\pi/e)\tau$ with $\tau = 0, 1, 2 \dots$.

3. Conclusions

In [19], authors studied the relativistic quantum dynamics of bosonic charged particle in the presence of external fields in the cosmic string space-time subject to a Cornell-type potential. In [59], authors studied a scalar field in the presence of external fields including a magnetic quantum flux in the space-time with a spacelike dislocation subject to a linear potential. In [60], authors investigated a spin-0 massive charged particle in the presence of external fields including a magnetic quantum flux in the space-time with a spacelike dislocation subject to a Coulomb-type potential.

In this paper, we have investigated quantum effects of torsion and topological defects that stem from a space-time with a spacelike dislocation under the influence of a Cornell-type potential in the relativistic quantum system. By solving the Klein-Gordon equation subject to a uniform magnetic field including a magnetic flux in the presence of a Cornell-type potential, we have obtained the energy eigenvalues Eq. (26) and corresponding eigenfunctions Eq. (27). By imposing the additional recurrence condition $c_{n+1} = 0$, we have obtained the individual energy levels and corresponding wave-function, as for example, $n = 1$ and others are in the same way. The presence of the torsion parameter modifies the energy levels and breaks their degeneracy. We have discussed three cases (subsection 2.1) for zero Coulomb potential, $\eta_c \rightarrow 0$, (subsection 2.2) zero linear potential, $\eta_L \rightarrow 0$, and (subsection 2.3) zero torsion parameter $\chi \rightarrow 0$ in the considered relativistic quantum system. In subsection 2.1–2.2, we have seen that for $\alpha \rightarrow 1$, the energy eigenvalues Eq. (40) and Eq. (51) are consistent with those results obtained in [59, 60], respectively. Thus, the presence of the cosmic string parameter shifts the energy levels in comparison to those obtained in [59, 60]. Furthermore, by comparing the energy eigenvalues Eq. (26) with those results obtained in [59, 60], we have seen that the presence of an extra potential term as well the cosmic string parameter modifies the energy eigenvalues. In sub-section 2.3, for zero-zero magnetic quantum flux, $\Phi_B \rightarrow 0$, we have seen that the energy eigenvalues Eq. (61) is consistent with those results obtained in [19]. Hence, the relativistic energy eigenvalues Eq. (61) is the extended result in comparison to those in [19] due to the presence of a magnetic quantum flux Φ_B . In addition, by comparing the energy eigenvalues Eq. (26) with the result obtained in [19], we have seen that the presence of the torsion parameter χ as well as the magnetic quantum flux Φ_B modify the energy eigenvalues where the degeneracy of the energy levels is broken by the torsion parameter.

We have seen in each case that the angular quantum number l is shifted, $l \rightarrow l_{eff} = (1/\alpha)(l - \Phi - k\chi)$, an effective angular quantum number. Thus, the relativistic energy eigenvalues obtained in subsection 2.1–2.2 depends on the geometric quantum phase [75, 76] as well as the torsion parameter and only the geometric quantum phase in subsection 2.3. This dependence of the relativistic energy eigenvalues on the geometric quantum phase gives rise to a relativistic analogue of the Aharonov-Bohm effect for bound states [11, 52, 54, 59, 61, 62, 82–86]. Besides, we have that $E_{n,l}(\Phi_B + \Phi_0) = E_{n,l \mp \tau}(\Phi_B)$ where $\Phi_0 = \pm(2\pi/e)\tau$ with $\tau = 0, 1, 2 \dots$.

It is well known in nonrelativistic quantum mechanics that the Landau quantization is the simplest system that would work with the studies of quantum Hall effect. Therefore, the relativistic quantum systems analyzed in this work would be used for investigating the influence of torsion, the cosmic string as well as the Cornell-type potential on the thermodynamic properties of quantum systems [94–97], searching a relativistic analogue of the quantum Hall effect [98, 99], and the displaced Fock states [100] in a topological defects space-time with a spacelike dislocation.

Data Availability

No data has been used to prepare this paper.

Conflicts of Interest

The author declares that there is no conflict of interest regarding publication this paper.

Acknowledgments

The author sincerely acknowledges the anonymous kind referee(s) for their valuable comments and suggestion which have greatly improved the present paper.

References

- [1] B. D. B. Figueiredo, I. D. Soares, and J. Tiomno, "Gravitational coupling of Klein-Gordon and Dirac particles to matter vorticity and spacetime torsion," *Classical and Quantum Gravity*, vol. 9, no. 6, pp. 1593–1617, 1992.
- [2] F. Ahmed, "The energy-momentum distributions and relativistic quantum effects on scalar and spin-half particles in a Gödel-type space-time," *European Physical Journal C: Particles and Fields*, vol. 78, no. 7, p. 598, 2018.
- [3] Z. Wang, Z. W. Long, C. Y. Long, and M. L. Wu, "Relativistic quantum dynamics of a spinless particle in the Som-Raychaudhuri space-time," *The European Physical Journal Plus*, vol. 130, no. 3, p. 36, 2015.
- [4] J. Carvalho, A. M. d. M. Carvalho, and C. Furtado, "Quantum influence of topological defects in Gödel-type space-times," *The European Physical Journal C*, vol. 74, no. 6, article 2935, 2014.
- [5] R. L. L. Vitória, C. Furtado, and K. Bakke, "Linear confinement of a scalar particle in a Gödel-type spacetime," *European Physical Journal C: Particles and Fields*, vol. 78, no. 1, p. 44, 2018.
- [6] F. Ahmed, "Comment on "Linear confinement of a scalar particle in a Gödel-type space-time"," *European Physical Journal C: Particles and Fields*, vol. 79, no. 8, article 682, 2019.
- [7] F. Ahmed, "Linear confinement of a scalar and spin-0 particle in a topologically trivial flat Gödel-type space-time," *European Physical Journal C: Particles and Fields*, vol. 79, no. 2, p. 104, 2019.
- [8] N. Drukker, B. Fiol, and J. Simón, "Gödel-type universes and the Landau problem," *Journal of Cosmology and Astroparticle Physics*, vol. 2004, no. 10, p. 012, 2004.
- [9] S. Das and J. Gegenberg, "Gravitational non-commutativity and Gödel-like spacetimes," *General Relativity and Gravitation*, vol. 40, no. 10, pp. 2115–2129, 2008.
- [10] C. Furtado, B. G. C. da Cunha, F. Moraes, E. R. B. de Mello, and V. B. Bezerra, "Landau levels in the presence of disclinations," *Physics Letters A*, vol. 195, no. 1, pp. 90–94, 1994.
- [11] J. Carvalho, A. M. . M. Carvalho, E. Cavalcante, and C. Furtado, "Klein-Gordon oscillator in Kaluza-Klein theory," *European Physical Journal C: Particles and Fields*, vol. 76, no. 7, article 365, 2016.
- [12] F. Ahmed, "Relativistic quantum dynamics of spin-0 massive charged particle in the presence of external fields in 4D curved space-time with a cosmic string," *The European Physical Journal Plus*, vol. 135, no. 1, article 108, 2020.
- [13] H. Sobhani, H. Hassanabadi, and W. S. Chung, "A survey on the Klein-Gordon equation in the Gödel-type space-times," *International Journal of Geometric Methods in Modern Physics*, vol. 15, no. 3, article 1850037, 2018.
- [14] F. Ahmed, "Scalar quantum particle in $(1+2)$ -dimensions Gürses space-time and the energy-momentum distributions," *Annals of Physics*, vol. 401, no. 193, pp. 193–201, 2019.
- [15] F. Ahmed, "The Klein-Gordon oscillator in $(1+2)$ -dimensions Gürses space-time backgrounds," *Annals of Physics*, vol. 404, no. 1, pp. 1–9, 2019.
- [16] F. Ahmed, "The generalized Klein-Gordon oscillator with Coulomb-type potential in $(1+2)$ -dimensions Gürses space-time," *General Relativity and Gravitation*, vol. 51, no. 5, article 69, 2019.
- [17] F. Ahmed, "Linear confinement of spin-0 relativistic quantum particles in $(1+2)$ -dimensions Gürses space-time," *Annals of Physics*, vol. 411, no. 167941, p. 167941, 2019.
- [18] F. Ahmed, "Quantum effects on spin-0 massive charged particles with Coulomb-type potentials in $(1+2)$ -dimensions Gürses space-time," *General Relativity and Gravitation*, vol. 51, no. 10, article 129, 2019.
- [19] E. R. F. Medeiros and E. R. B. de Mello, "Relativistic quantum dynamics of a charged particle in cosmic string spacetime in the presence of magnetic field and scalar potential," *European Physical Journal C: Particles and Fields*, vol. 72, no. 6, article 2051, 2012.
- [20] F. Ahmed, "The Dirac equation in a class of topologically trivial flat Gödel-type space-time backgrounds," *European Physical Journal C: Particles and Fields*, vol. 79, no. 6, article 534, 2019.
- [21] M. de Montigny, S. Zare, and H. Hassanabadi, "Fermi field and Dirac oscillator in a Som-Raychaudhuri space-time," *General Relativity and Gravitation*, vol. 50, no. 5, article 47, 2018.
- [22] P. Sedaghatnia, H. Hassanabadi, and F. Ahmed, "Dirac fermions in Som-Raychaudhuri space-time with scalar and vector potential and the energy momentum distributions," *European Physical Journal C: Particles and Fields*, vol. 79, no. 6, article 541, 2019.
- [23] F. Ahmed, "The Dirac equation in $(1+2)$ -dimensional Gürses space-time backgrounds," *The European Physical Journal Plus*, vol. 134, no. 10, article 518, 2019.
- [24] A. Vilenkin and E. P. S. Shellard, *Cosmic Strings and Other Topological Defects*, Cambridge Monographs on Mathematical Physics, Cambridge University Press, Cambridge, 1994.
- [25] M. B. Hindmarsh and T. W. B. Kibble, "Cosmic strings," *Reports on Progress in Physics*, vol. 58, no. 5, pp. 477–562, 1995.
- [26] A. Vilenkin, "Gravitational field of vacuum domain walls," *Physics Letters B*, vol. 133, no. 3-4, pp. 177–179, 1983.
- [27] A. Vilenkin, "Cosmic strings and domain walls," *Physics Reports*, vol. 121, no. 5, pp. 263–315, 1985.
- [28] W. A. Hiscock, "Exact gravitational field of a string," *Physical Review A*, vol. 31, no. 12, pp. 3288–3290, 1985.
- [29] B. Linet, "The static metrics with cylindrical symmetry describing a model of cosmic strings," *General Relativity and Gravitation*, vol. 17, no. 11, pp. 1109–1115, 1985.
- [30] M. O. Katanaev and I. V. Volovich, "Theory of defects in solids and three-dimensional gravity," *Annals of Physics*, vol. 216, no. 1, pp. 1–28, 1992.

- [31] C. Furtado and F. Moraes, "On the binding of electrons and holes to disclinations," *Physics Letters A*, vol. 188, no. 4-6, pp. 394–396, 1994.
- [32] T. W. B. Kibble, "Topology of cosmic domains and strings," *Journal of Physics A*, vol. 9, no. 8, pp. 1387–1398, 1976.
- [33] L. C. N. Santos and C. C. Barros Jr., "Relativistic quantum motion of spin-0 particles under the influence of noninertial effects in the cosmic string space-time," *European Physical Journal C: Particles and Fields*, vol. 78, no. 1, article 13, 2018.
- [34] M. Hosseini, H. Hassanabadi, S. Hassanabadi, and P. Sedaghatnia, "Klein-Gordon oscillator in the presence of a Cornell potential in the cosmic string space-time," *International Journal of Geometric Methods in Modern Physics*, vol. 16, no. 4, article 1950054, 2019.
- [35] M. Hosseinpour, H. Hassanabadi, and F. M. Andrade, "The DKP oscillator with a linear interaction in the cosmic string space-time," *European Physical Journal C: Particles and Fields*, vol. 78, no. 2, article 93, 2018.
- [36] M. Hosseinpour, F. M. Andrade, E. O. Silva, and H. Hassanabadi, "Scattering and bound states for the Hulthén potential in a cosmic string background," *European Physical Journal C: Particles and Fields*, vol. 77, no. 5, article 270, 2017.
- [37] H. Sobhani, H. Hassanabadi, and W. S. Chung, "Effects of cosmic string framework on the thermodynamical properties of anharmonic oscillator using the ordinary statistics and the q -deformed superstatistics approaches," *European Physical Journal C: Particles and Fields*, vol. 78, no. 2, article 106, 2018.
- [38] M. Hosseinpour and H. Hassanabadi, "Scattering states of Dirac equation in the presence of cosmic string for Coulomb interaction," *International Journal of Modern Physics A: Particles and Fields; Gravitation; Cosmology; Nuclear Physics*, vol. 30, no. 21, article 1550124, 2015.
- [39] M. de Montigny, M. Hosseinpour, and H. Hassanabadi, "The spin-zero Duffin-Kemmer-Petiau equation in a cosmic-string space-time with the Cornell interaction," *International Journal of Modern Physics A: Particles and Fields; Gravitation; Cosmology; Nuclear Physics*, vol. 31, no. 36, article 1650191, 2016.
- [40] S. Zare, H. Hassanabadi, and M. de Montigny, "Non-inertial effects on a generalized DKP oscillator in a cosmic string space-time," *General Relativity and Gravitation*, vol. 52, no. 3, article 25, 2020.
- [41] M. Hosseinpour, H. Hassanabadi, and M. de Montigny, "The Dirac oscillator in a spinning cosmic string spacetime," *European Physical Journal C: Particles and Fields*, vol. 79, no. 4, article 311, 2019.
- [42] H. Hassanabadi, S. Zare, and M. de Montigny, "Relativistic spin-zero bosons in a Som-raychaudhuri space-time," *General Relativity and Gravitation*, vol. 50, no. 8, article 104, 2018.
- [43] M. Hosseinpour and H. Hassanabadi, "The DKP oscillator in spinning cosmic string background," *Advances in High Energy Physics*, vol. 2018, Article ID 2959354, 8 pages, 2018.
- [44] H. Hassanabadi, W. S. Chung, S. Zare, and H. Sobhani, "Investigation of DKP equation for spin-zero system in the presence of Gödel-type background space-time," *European Physical Journal C: Particles and Fields*, vol. 78, no. 1, article 83, 2018.
- [45] M. Hosseinpour and H. Hassanabadi, "DKP equation in a rotating frame with magnetic cosmic string background," *The European Physical Journal Plus*, vol. 130, no. 11, article 236, 2015.
- [46] H. Hassanabadi, A. Afshardoost, and S. Zarrinkamar, "On the motion of a quantum particle in the spinning cosmic string space-time," *Annals of Physics*, vol. 356, pp. 346–351, 2015.
- [47] A. Afshardoost and H. Hassanabadi, "Non-central potentials in the cosmic string space-time," *Canadian Journal of Physics*, vol. 94, no. 1, pp. 71–74, 2016.
- [48] M. Hosseinpour, H. Hassanabadi, and M. de Montigny, "Klein-Gordon field in spinning cosmic-string space-time with the Cornell potential," *International Journal of Geometric Methods in Modern Physics*, vol. 15, no. 10, article 1850165, 2018.
- [49] M. K. Bahar and F. Yasuk, "Exact Solutions of the Mass-Dependent Klein-Gordon Equation with the Vector Quark-Antiquark Interaction and Harmonic Oscillator Potential," *Advances in High Energy Physics*, vol. 2013, Article ID 814985, 6 pages, 2013.
- [50] C. Alexandrou, P. de Forcrand, and O. Jahn, "The ground state of three quarks," *Nuclear Physics B - Proceedings Supplements*, vol. 119, pp. 667–669, 2003.
- [51] F. Ahmed, "Relativistic quantum motion of spin-0 particles with a Cornell-type potential in $(1+2)$ -dimensional Gürses spacetime backgrounds," *Modern Physics Letters A*, vol. 34, no. 38, article 1950314, 2019.
- [52] F. Ahmed, "Klein-Gordon Oscillator in the Presence of External Fields in a Cosmic Space- Time with a Space-Like Dislocation and Aharonov-Bohm Effect," *Advances in High Energy Physics*, vol. 2020, Article ID 5691025, 10 pages, 2020.
- [53] R. L. L. Vitória and K. Bakke, "Relativistic quantum effects of confining potentials on the Klein-Gordon oscillator," *The European Physical Journal Plus*, vol. 131, no. 2, article 36, 2016.
- [54] E. V. B. Leite, H. Belich, and R. L. L. Vitória, "Effects of the Cornell-type potential on a position-dependent mass system in Kaluza-Klein theory," *Advances in High Energy Physics*, vol. 2019, Article ID 6740360, 7 pages, 2019.
- [55] M. S. Cunha, C. R. Muniz, H. R. Christiansen, and V. B. Bezerra, "Relativistic Landau levels in the rotating cosmic string spacetime," *European Physical Journal C: Particles and Fields*, vol. 76, no. 9, article 512, 2016.
- [56] R. L. L. Vitória and H. Belich, "Effects of a linear central potential induced by the Lorentz symmetry violation on the Klein-Gordon oscillator," *European Physical Journal C: Particles and Fields*, vol. 78, no. 12, article 999, 2018.
- [57] R. L. L. Vitória and H. Belich, "A central potential with a massive scalar field in a Lorentz symmetry violation environment," *Advances in High Energy Physics*, vol. 2019, Article ID 1248393, 13 pages, 2019.
- [58] A. L. C. de Oliveira and E. R. B. de Mello, "Exact solutions of the Klein-Gordon equation in the presence of a dyon, magnetic flux and scalar potential in the spacetime of gravitational defects," *Classical and Quantum Gravity*, vol. 23, no. 17, pp. 5249–5263, 2006.
- [59] R. L. L. Vitória and K. Bakke, "Aharonov-Bohm effect for bound states in relativistic scalar particle systems in a spacetime with a spacelike dislocation," *International Journal of Modern Physics D: Gravitation; Astrophysics and Cosmology*, vol. 27, no. 2, article 1850005, 2018.

- [60] R. L. L. Vitória and K. Bakke, "On the interaction of the scalar field with a Coulomb-type potential in a spacetime with a screw dislocation and the Aharonov-Bohm effect for bound states," *The European Physical Journal Plus*, vol. 133, no. 11, article 490, 2018.
- [61] M. Peshkin and A. Tonomura, *The Aharonov-Bohm effect*, vol. 340 of Lecture Notes in Physics, Springer, Berlin, Germany, 1989.
- [62] V. B. Bezerra, "Gravitational analogs of the Aharonov-Bohm effect," *Journal of Mathematical Physics*, vol. 30, no. 12, pp. 2895–2899, 1989.
- [63] R. A. Puntigam and H. H. Soleng, "Volterra distortions, spinning strings, and cosmic defects," *Classical and Quantum Gravity*, vol. 14, no. 5, pp. 1129–1149, 1997.
- [64] V. B. Bezerra, "Global effects due to a chiral cone," *Journal of Mathematical Physics*, vol. 38, no. 5, pp. 2553–2564, 1997.
- [65] H. Kleinert, *Gauge Fields in Condensed Matter*, vol. 2, World Scientific Publishing Co, Singapore, 1989.
- [66] C. Furtado and F. Moraes, "Landau levels in the presence of a screw dislocation," *Europhysics Letters (EPL)*, vol. 45, no. 3, pp. 279–282, 1999.
- [67] J. Wang, K. Ma, K. Li, and H. Fan, "Deformations of the spin currents by topological screw dislocation and cosmic dispiration," *Annals of Physics*, vol. 362, no. 327, pp. 327–335, 2015.
- [68] C. Filgueiras, M. Rojas, G. Aciole, and E. O. Silva, "Landau quantization, Aharonov-Bohm effect and two-dimensional pseudoharmonic quantum dot around a screw dislocation," *Physics Letters A*, vol. 380, no. 45, pp. 3847–3853, 2016.
- [69] G. D. A. Marques, V. B. Bezerra, C. Furtado, and F. Moraes, "Quantum effects due to a magnetic flux associated to a topological defect," *International Journal of Modern Physics A: Particles and Fields; Gravitation; Cosmology; Nuclear Physics*, vol. 20, no. 26, pp. 6051–6064, 2012.
- [70] C. Furtado, V. B. Bezerra, and F. Moraes, "Quantum scattering by a magnetic flux screw dislocation," *Physics Letters A*, vol. 289, no. 3, pp. 160–166, 2001.
- [71] R. L. L. Vitória, "Noninertial effects on a scalar field in a spacetime with a magnetic screw dislocation," *European Physical Journal C: Particles and Fields*, vol. 79, no. 10, article 844, 2019.
- [72] R. L. L. Vitória and K. Bakke, "Rotating effects on the scalar field in the cosmic string spacetime, in the spacetime with space-like dislocation and in the spacetime with a spiral dislocation," *European Physical Journal C: Particles and Fields*, vol. 78, no. 3, article 175, 2018.
- [73] L. C. N. Santos and C. C. Barros Jr., "Scalar bosons under the influence of noninertial effects in the cosmic string spacetime," *European Physical Journal C: Particles and Fields*, vol. 77, no. 3, article 186, 2017.
- [74] R. L. L. Vitória and K. Bakke, "Torsion effects on a relativistic position-dependent mass system," *General Relativity and Gravitation*, vol. 48, no. 12, article 161, 2016.
- [75] Y. Aharonov and D. Bohm, "Significance of electromagnetic potentials in the quantum theory," *Physical Review*, vol. 115, no. 3, pp. 485–491, 1959.
- [76] G. d. A. Marques, C. Furtado, V. B. Bezerra, and F. Moraes, "Landau levels in the presence of topological defects," *Journal of Physics A: Mathematical and General*, vol. 34, no. 30, pp. 5945–5954, 2001.
- [77] C. Furtado and F. Moraes, "Harmonic oscillator interacting with conical singularities," *Journal of Physics A: Mathematical and General*, vol. 33, no. 31, pp. 5513–5519, 2000.
- [78] R. Jackiw, A. I. Milstein, S.-Y. Pi, and I. S. Terekhov, "Induced current and Aharonov-Bohm effect in graphene," *Physical Review B*, vol. 80, no. 3, article 033413, 2009.
- [79] M. A. Anacleto, I. G. Salako, F. A. Brito, and E. Passos, "Analogue Aharonov-Bohm effect in neo-Newtonian theory," *Physical Review D*, vol. 92, no. 12, article 125010, 2015.
- [80] V. R. Khalilov, "Bound states of massive fermions in Aharonov-Bohm-like fields," *European Physical Journal C: Particles and Fields*, vol. 74, no. 1, p. 2708, 2014.
- [81] C. Coste, F. Lund, and M. Umeki, "Scattering of dislocated wave fronts by vertical vorticity and the Aharonov-Bohm effect. I. Shallow water," *Physical Review E*, vol. 60, no. 4, pp. 4908–4916, 1999.
- [82] C. Furtado, V. B. Bezerra, and F. Moraes, "Aharonov-Bohm effect for bound states in Kaluza-Klein theory," *Modern Physics Letters A*, vol. 15, no. 4, pp. 253–258, 2011.
- [83] C. Furtado, F. Moraes, and V. B. Bezerra, "Global effects due to cosmic defects in Kaluza-Klein theory," *Physical Review D*, vol. 59, no. 10, article 107504, 1999.
- [84] E. V. B. Leite, H. Belich, and K. Bakke, "Aharonov-Bohm effect for bound states on the confinement of a relativistic scalar particle to a Coulomb-type potential in Kaluza-Klein theory," *Advances in High Energy Physics*, vol. 2015, Article ID 925846, 6 pages, 2015.
- [85] E. V. B. Leite, R. L. L. Vitória, and H. Belich, "Effects of the Kaluza-Klein theory on a Klein Gordon particle with position-dependent mass," *Modern Physics Letters A*, vol. 34, no. 38, article 1950319, 2019.
- [86] F. Ahmed, "The generalized Klein-Gordon oscillator in the background of cosmic string space-time with a linear potential in the Kaluza-Klein theory," *European Physical Journal C: Particles and Fields*, vol. 80, no. 3, article 211, 2020.
- [87] H. Belich, E. O. Silva, M. M. Ferreira, and M. T. D. Orlando, "Aharonov-Bohm-Casher problem with a nonminimal Lorentz-violating coupling," *Physical Review D*, vol. 83, no. 12, article 125025, 2011.
- [88] W. Greiner, *Relativistic Quantum Mechanics: Wave Equations*, Springer-Verlag, Berlin, Germany, 3rd edition, 2000.
- [89] A. Ronveaux, *Heuns Differential Equations*, Oxford University Press, Oxford, 1995.
- [90] S. Y. Slavyanov and W. Lay, *Special Functions: A Unified Theory Based in Singularities*, Oxford University Press, New York, 2000.
- [91] G. B. Arfken and H. J. Weber, *Mathematical Methods for Physicists*, Elsevier Academic Press, London, 2005.
- [92] A. Vercin, "Two anyons in a static uniform magnetic field. Exact solution," *Physics Letters B*, vol. 260, no. 1-2, pp. 120–124, 1991.
- [93] J. Myrheim, E. Halvorsen, and A. Vercin, "Two anyons with Coulomb interaction in a magnetic field," *Physics Letters B*, vol. 278, no. 1-2, pp. 171–174, 1992.
- [94] H. Hassanabadi and M. Hosseinpour, "Thermodynamic properties of neutral particle in the presence of topological defects in magnetic cosmic string background," *European Physical Journal C: Particles and Fields*, vol. 76, no. 10, article 553, 2016.
- [95] M. Eshghi and H. Mehraban, "Study of a 2D charged particle confined by a magnetic and AB flux fields under the radial

- scalar power potential,” *The European Physical Journal Plus*, vol. 132, no. 3, article 121, 2017.
- [96] A. N. Ikot, B. C. Lutfuoglu, M. I. Ngwueke, M. E. Udoh, S. Zare, and H. Hassanabadi, “Klein-Gordon equation particles in exponential-type molecule potentials and their thermodynamic properties in D dimensions,” *The European Physical Journal Plus*, vol. 131, no. 12, article 419, 2016.
- [97] H. Hassanabadi, S. Sargolzaeipor, and B. H. Yazarloo, “Thermodynamic properties of the three-dimensional Dirac oscillator with Aharonov-Bohm field and magnetic monopole potential,” *Few-Body Systems*, vol. 56, no. 2-3, pp. 115–124, 2015.
- [98] R. E. Prange and S. M. Girvin, *The Quantum Hall Effect*, Springer-Verlag, New York, 1990.
- [99] D. Chowdhury and B. Basu, “Effect of a cosmic string on spin dynamics,” *Physical Review D*, vol. 90, no. 12, article 125014, 2014.
- [100] J. L. de Melo, K. Bakke, and C. Furtado, “Geometric quantum phase for displaced states for a particle with an induced electric dipole moment,” *EPL (Europhysics Letters)*, vol. 115, no. 2, article 20001, 2016.