



Assessment and Mapping of Micronutrient Cation Distribution in Agricultural Research Station Soil: A Geospatial Approach

Kamlesh Bhil ^{a*}, M. K. Sharma ^a, Neha Meena ^a,
Narendra Danga ^a, Rajesh Meena ^a, Tushar ^b,
Sarita Choudhary ^b and Kamal Gurjar ^b

^a Department of Soil Science and Agricultural Chemistry, College of Agriculture - Umedganj, AU - Kota (Rajasthan - 324 001), India.

^b Department of Soil Science and Agricultural Chemistry, Rajasthan College of Agriculture, MPUAT-Udaipur (Rajasthan - 313001), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i183393

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/103486>

Original Research Article

Received: 21/05/2023

Accepted: 24/07/2023

Published: 31/07/2023

ABSTRACT

Aim: The objective was to construct fertility maps of the area, specifically focusing on micronutrient cations. The research aims to provide valuable insights into the spatial distribution of micronutrients, aiding in the development of targeted agricultural practices and informed decision-making.

Study Design: Soil sampling, soil analysis and soil mapping.

Place and Duration of Study: Agricultural Research Station, Umedganj-Kota in 2019 and 2020

*Corresponding author: E-mail: kamleshbhilag7073@gmail.com;

Methodology: The study involved the collection of 300 soil samples, with precise coordinate's locked using GPS technology. Soil variability maps for soil micronutrient Zn, Fe, Mn and Cu were generated by using Arc GIS 10.5 on the basis of latitude and longitude of sample site by using hand GPS device and analyzed sample data.

Results: The available-Zn of the soils ranged from 0.03 to 4.39 mg kg⁻¹ with a mean value 0.61 mg kg⁻¹, the soil falls under low (64.67%) to medium (30.33%) in available Zn. The available-Fe of the soils ranged from 1.09 to 12.54 mg kg⁻¹ with a mean value 4.29 mg kg⁻¹, the soils fall under low (58.00%) and medium (31.67%) categories for available Fe. The available-Mn of the soils ranged from 2.27-22.31 mg kg⁻¹ with a mean value 5.74 mg kg⁻¹, the soils of the farm falls under medium (28.33%) to high (69.34%) in available Mn. The available-Cu of the soils ranged from 0.32-2.25 mg kg⁻¹ with a mean value 1.09 mg kg⁻¹, the soil comes under high (57.00%) and very high (30.33%) categories for available Cu.

Conclusion: Spatial variability map it was found that a high amount of Zn, Mn, Fe present in the soil of field number 14 and 12A compared to other fields. The possible cause behind this may be due to adoption of organic farming practices in field number 14 and execution of integrated farming system modules in field number 12A from some year.

Keywords: GPS; arc GIS 10.5; organic farming practices and integrated farming system.

1. INTRODUCTION

Soil is a finite, non-renewable natural resource that defines the feasibility of implementing an agricultural development program [1]. Micronutrients are vital for optimal plant growth and development, despite being required in small quantities. Zinc (Zn), iron (Fe), manganese (Mn), copper (Cu), boron (B), molybdenum (Mo), and nickel (Ni) are essential micronutrients. While discussions on soil fertility often focus on macronutrients like nitrogen, phosphorus, and potassium, the significance of micronutrients should not be underestimated. These micronutrients play critical roles in various physiological and biochemical processes within plants. Ziaean and Malakouti, [2] found that Fe, Mn, Zn and Cu fertilization significantly increased grain yield, straw yield, 1000-grain weight, and the number of grains per spikelet. Zinc is essential for enzyme activity and growth hormone synthesis [3], while iron is crucial for the respiration, photosynthesis and nitrogen fixation processes [4]. Manganese is essential for oxidation and reduction activities, as well as electron transport in photosynthesis [5]. Copper is involved in protein and lignin synthesis Yruela, [6]. The availability and distribution of micronutrient cations significantly influence plant health and productivity. Inadequate levels can lead to nutrient deficiencies, causing stunted growth, reduced yields, and increased susceptibility to pests and diseases. Conversely, excessive concentrations can lead to toxicity and hinder plant growth.

Geographical Information System (GIS) is a powerful tool that integrates diverse spatial

information, such as agro-climate zones, land use, and soil management. GPS satellites collect data transmitted to ground-based receivers, which is then utilized by GIS software to analyze and utilize the information. By utilizing GPS technology for soil sample collection, soil variability maps can be created, enabling the measurement of nutrient status and guiding efficient fertilizer use. GIS techniques facilitate the generation of site-specific nutrient recommendations, enhancing precision in agricultural practices.

Soil fertility mapping involves the systematic collection and analysis of soil samples to assess geographical representation showing diversity of essential nutrients, pH levels, organic matter content, and other soil properties. This mapping provides valuable information about the fertility status of agricultural fields, aiding in the development of targeted soil management strategies

2. MATERIALS AND METHOD


Agricultural Research Station, Ummedganj-Kota is situated in Agro-climatic zone V "Humid South Eastern Plain" which falls in the south eastern part of Rajasthan. According to Agro-ecological region map brought out by National Bureau of Soil Survey and Land Use Planning (NBSS & LUP, Nagpur), Kota falls in Agro-ecological region No 06 "Hot Semi-Arid Eco-Region with Shallow and Medium (Dominant) Black Soils" which covered 9.5% of the area of the country. Agricultural Research farm Ummedganj, Kota is located at

the 75°25' N latitude 25°13' E longitude and an altitude of 258 m above mean sea level.

Soil sampling was carried out after harvesting of Rabi crops (in the months of April to May, 2019). Total area of the farm was 107 ha, which was officially marked as farm and mostly under cultivation. Total 300 Geo-referenced surface (0-15 cm) soil samples representing the field were collected following standard methods from 18 blocks/ fields of Agricultural Research Station, Ummедganj-Kota. Soil variability maps for various soil properties were produced using the latitude and longitude, the point shape files showing the location of the

observation were generated by using Arc GIS 10.5.

The collected soil samples were dried in shade, cursed and sieved for laboratory analysis. The micronutrients (Fe, Mn, Zn and Cu) were extracted using 0.005M DTPA, 0.01M calcium chloride and 0.1M triethanolamine buffered at pH 7.3 [7]. DTPA extractant has the ability to chelate Zn, Cu, Fe and Mn in competition with Ca⁺⁺ and Mg⁺⁺ and unlike most chelating agents. Take 10 g soil and add 20 ml of DTPA extractant than shake for 2 hrs in a mechanical shaker and filter it. Concentration of micronutrients measured by Atomic Absorption Spectrophotometer (AAS) in the filtrate.



Map 1. Sampling points at agricultural research station farm, Ummедganj, Kota

Table 1. Six tier ratings of DTPA extractable micronutrients in soil

S. NO.	Ratings	Fe	Mn	Zn	Cu
		Mg kg ⁻¹	Mg kg ⁻¹	Mg kg ⁻¹	Mg kg ⁻¹
1.	Very low	<2.5	<1	<0.3	<0.1
2.	Low	2.5-4.5	1-2	0.3-0.6	0.1-0.2
3.	Medium	4.5-9	2-4	0.6-1.2	0.2-0.4
4.	Moderately high	9-18	4-8	1.2-1.8	0.4-0.8
5.	High	18-27	8-16	1.8-2.4	0.8-1.2
6.	Very high	>27	>16	>2.4	>1.2
	Critical limits	10	2.0	0.6	0.2

[8]

3. DATA ANALYSIS AND INTERPRETATION

Table 2. Distribution of available Zn and Fe in the soils of agricultural research station, Ummedganj-Kota

Available-Zn (mg kg ⁻¹)	No. of samples	% Samples	Available-Fe (mg kg ⁻¹)	No. of samples	% Samples
Very low (<0.3)	1	0.33	Very low (<2.5)	23	7.67
Low (0.3-0.6)	194	64.67	Low (2.5-4.5)	174	58.00
Medium (0.6-1.2)	91	30.33	Moderate (4.5-9)	95	31.67
Moderately high (1.2-1.8)	9	3.00	Moderately high (9-18)	8	2.66
High (1.8-2.4)	4	1.33	High (18-27)	Nil	Nil
Very high (>2.4)	1	0.33	Very high (>27)	Nil	Nil
General statistics					
Range	0.03-4.39		Range	1.09-12.54	
Mean	0.61		Mean	4.29	
Standard deviation	0.38		Standard deviation	1.57	
CV%	61.73		CV%	36.60	

Table 3. Distribution of available Mn and Cu in the soils of agricultural research station, Ummedganj-Kota


Available-Mn (mg kg ⁻¹)	No. of samples	% Samples	Available-Cu (mg kg ⁻¹)	No. of samples	% Samples
Very low (<1)	Nil	Nil	Very low (<0.1)	Nil	Nil
Low (1-2)	Nil	Nil	Low (0.1-0.2)	Nil	Nil
Medium (2-4)	85	28.33	Medium (0.2-0.4)	5	1.67
Moderately high (4-8)	164	54.67	Moderately high (0.4-0.8)	33	11.00
High (8-16)	44	14.67	High (0.8-1.2)	171	57.00
Very high (>16)	7	2.33	Very high(>1.2)	91	30.33
General statistics					
Range	2.27-22.31		Range	0.30-2.25	
Mean	5.74		Mean	1.09	
Standard deviation	3.11		Standard deviation	0.29	
CV%	54.15		CV%	26.88	

4. SPATIAL INTERPOLATION TECHNIQUES USED FOR FERTILITY MAPPING

Spatial interpolation is the process of using points with known values to estimate values at other unknown points. These techniques helped to generate full area micronutrient fertility maps through use of some point of soil sampling.

Table 4. Extent of area under different Fe- availability category of the soils of agricultural research station, Ummedganj-Kota

Fe-Availability Category	Available-Fe, mg kg ⁻¹	Ranking of the class on the basis of area
I	1.09-3.38 (77)*	II
II	3.39-5.67 (188)	I
III	5.68-7.96 (26)	III
IV	7.97-10.25 (6)	IV
V	10.26-12.54 (3)	V



Map 2. Spatial distribution of available iron in the soils of agricultural research station, Ummedganj-Kota

Table 5. Extent of area under different zinc availability category of the soils of agricultural research station, Ummedganj-Kota


Zn-Availability Category	Available-Zn (mg kg ⁻¹)	Ranking of the class on the basis of area
I	0.03-0.90 (263)*	I
II	0.91-1.77 (32)	II
III	1.78-2.65 (4)	III
IV	2.66-3.52 (0)	V
V	3.53-4.39 (1)	IV

* In bracket, number of soil samples under the category


Table 6. Extent of area under different Mn availability category of the soils of agricultural research station, Ummedganj-Kota

Mn-Availability Category	Available-Mn, mg kg ⁻¹	Ranking of the class on the basis of area
I	2.27-6.28 (206)*	I
II	6.29-10.29 (79)	II
III	10.30-14.29 (7)	IV
IV	14.30-18.30 (3)	V
V	18.31-22.31(5)	III

* In bracket, number of soil samples under the category




Map 3. Spatial Distribution of available zink in the soils of agricultural research station, Ummédganj-Kota



Map 4. Spatial distribution of available manganese in the soils of agricultural research station, Ummédganj-Kota

Table 7. Extent of area under different Cu- availability category of the soils of agricultural research station, Umedganj-Kota

Cu-Availability Category	Available-Cu, mgkg ⁻¹	Ranking of the class on the basis of area
I	0.30-0.69 (21) *	IV
II	0.70-1.08 (135)	I
III	1.09-1.47 (113)	II
IV	1.48-1.86 (29)	III
V	1.87-2.25 (2)	V



Map 5. Spatial distribution of available copper in the soils of agricultural research station, Umedganj-Kota

5. RESULTS AND DISCUSSION

5.1 Available Zn

The available-Zn in the soils of ARS ranged from 0.03 to 4.39 mg kg⁻¹ with a mean value 0.61 mg kg⁻¹, on the basis of analyzed data, the soils of ARS farm falls under low (64.67%) to medium (30.33%) in available Zn. Zinc deficiency is widespread not only in soils and crops of India [9] but also in many countries of the world [10]. Since, the soils were alkaline and rich in CaCO₃, zinc may be precipitated as hydroxides and carbonates under alkaline pH range. Therefore, their solubility and mobility may be decreased

resulting in reduced availability. In contrast, Zn extractability from soil is negatively related to [11] and calcium carbonate content in soil [12]. Therefore, low plant availability of Zn can be expected in calcareous soils. On the basis of soil analysis data, the field number 3 and 14 (1.20 and 1.07 mg kg⁻¹) have higher amount of Zn compared to other fields. The results are in close agreement with the findings of Rajendiran et al., [13] in the Alirajpur district of Madhya Pradesh, Kumar et al., [14] in the Theni district of Tamil Nadu, Amara et al., [15] of the soils of Boguru micro watershed in Karnataka state of India. Similar finding in calcareous soils were reported by Tagore et al., [16].

5.2 Available Fe

The available-Fe in the soils of ARS ranged from 1.09 to 12.54 mg kg⁻¹ with a mean value 4.29 mg kg⁻¹. The soils of ARS farm falls under low (58.00%) and medium (31.67%) categories in available Fe. Iron deficiency can be explained by high soil pH (7.7) and little soil organic carbon (SOC) [17], which is also indicated by the negative correlation between Fe and pH and a positive correlation between Fe and SOC. Low Fe content may be due to precipitation of Fe²⁺ by CaCO₃ and decrease the availability. On the basis of soil analysis data that field number 14 (10.08 mg kg⁻¹) contains a higher amount of available Fe compare to other fields, because continuous organic farming practices are performed in field number 14 and integrated farming system modules executed in field number 12A for some years. The results are in close agreement with the findings of Amara et al., [15] of the soils of Bogur micro watershed in Karnataka state of India, Madhu et al., [18] in the soils of Jamuna Paar regions of Allahabad District, Uttar Pradesh. This trend of finding was also noticed by Sharma et al., [19] in central India.

5.3 Available Mn

The available-Mn in the soils of ARS ranged from 2.27-22.31 mg kg⁻¹ with a mean value 5.74 mg kg⁻¹, the soils of ARS farm falls under medium (28.33%) to high (69.34%) in available Mn. Manganese deficiency usually does not occur in black soils because a sizeable portion of Mn is bound with manganese oxide which may be readily available [20]. On the basis of soil analysis data, the field number 14 (10.08 mg kg⁻¹) have contained higher amount of available Mn compare to other fields because organic farming practices applied in this field from some years. The results are in close agreement with the findings of Amara et al., [15] in the soils of Bogur micro watershed in Karnataka state of India, Dash et al., [21] regional research and technology transfer station (RRTTS) and Krishi vigyan kendra (KVK) farm, Dhenkanal located in the mid-central table land Agro climatic zone of Odisha, India. Sood et al., [22] reported similar trends in Muktsar district of Punjab.

5.4 Available Cu

The available-Cu in the soils of Agriculture Research Station (ARS) ranged from 0.30-2.25 mg kg⁻¹ with a mean value 1.09 mg kg⁻¹, the soils

of ARS farm falls under high (57.00%) and very high (30.33%) categories in available Cu. Reason behind this, sufficient amount of Cu content in soils due to low availability of organic carbon, because copper formed chelates and has negative correlation with the OC. According to soil analytical data and spatial variability map the soils of field number 19 was found highest in amount of Cu content (1.48 mg kg⁻¹) at ARS, Kota. The results are in close agreement with the findings of Amara et al., [15] of the soil of Bogur micro watershed in Karnataka state of India, Madhu et al., [18] in the soils of Jamuna Paar region of Allahabad District, Uttar Pradesh. Kumar et al., [23] while studying spatial variability in Sekhawati region of Thar desert, India came up with similar results.

6. CONCLUSION

It can be concluded from above result that available- Zn, Fe, Mn and Cu ranged from 0.03 to 4.39 mg kg⁻¹, 1.09 to 12.54 mg kg⁻¹, 2.27-22.31 mg kg⁻¹ and 0.30-2.25 mg kg⁻¹ with mean value 0.61 mg kg⁻¹, 4.29 mg kg⁻¹, 5.74 mg kg⁻¹ and 1.09 mg kg⁻¹, respectively. Spatial variability map it was found that a high amount of Zn, Mn, Fe present in the soil of field number 14 and 12A compared to other fields. The possible cause behind this may be due to adoption of organic farming practices in field number 14 and execution of integrated farming system modules in field number 12A from some year. Cu was the highest amount evaluated in the soils of field number 19. Soil fertility index mean value was observed for Zn-1.21, Fe-1.15, Mn-1.96 and Cu-2.58, as a result, soil fertility class for Zn and Fe were low, medium for Mn and high fertility class for Cu. This data is also very useful for providing a base for making sound fertilizers recommendations for various crops and crop planning in near future. Spatial variability analysis of nutrients is one of key factor precision agriculture management.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Vedadri U, Naidu MVS. Characterization, classification and evaluation of soils in semi-arid ecosystem of Chillakur Mandal in SPSR Nelloredistrict of Andhra Pradesh.

- Journal of the Indian Society of Soil Science. 2018;66:9-19.
2. Ziaean AH, Malakouti MJ. Effect of micronutrient application on wheat production in calcareous soils. Prepared for the Second National Conference on Optimum Utilization of Chemical Fertilizers and Pesticide in Agriculture, Karaj, Iran; 2001.
 3. Mengel K, Kirkby EA. Principles of plant nutrition. Bern. International Potash Institute. 1987;687-695.
 4. Rout GR, Sahoo S. Role of iron in plant growth and metabolism. Reviews in Agricultural Science. 2015;3:1-24.
 5. Mukhopadhyay MJ, Sharma A. Manganese in cell metabolism of higher plants. The Botanical Review. 1991;57: 117-149.
 6. Yruea I. Copper in plants: acquisition, transport and interactions. Functional Plant Biology. 2009;36(5):409-430.
 7. Lindsay WL, Norvel WA. Development of a DTPA test for zinc, iron, manganese and copper. Soil Science Society American Journal. 1978;4:421-428.
 8. Katkar RN, Patil DB. Available micronutrient content in soils of Vidarbha. Souvenir of state level seminar in Department of Science and Agriculture Chemistry, Dr. PDKV, Akola Organized During. Jan., 2-3, 2010;161-166.
 9. Takkar PN, Singh MV, Ganeshmurthy AN. In. Plant Nutrient Needs, Supply, Efficiency and Policy Issues, 2000-2025, National Academy of Agricultural Sciences, New Delhi. 1997;238-264.
 10. Cakmak. Presented during FAI Annual Seminar at New Delhi, India; 2008.
 11. Yang Xw, Tian Xh, Lu Xc, Cao Yx, Chen Zh. Impacts of phosphorus and zinc levels on phosphorus and zinc nutrition and phytic acid concentration in wheat (*Triticum aestivum* L.). Journal of the Science of Food & Agriculture. 2011; 91:2322-2328.
 12. Iratkar AG, Giri JD, Kadam MM, Giri JN, Dabhade MB. Distribution of DTPA extractable micronutrients and their relationship with soil properties in soil of Parsori watershed of Nagpur district of Maharashtra. Asian Journal of Soil Science. 2014;9(2):297-299
 13. Rajendiran S, Dotaniya ML, Coumar MV, Sinha NK, Kundu S, Srivastava S, et al. Evaluation of soil fertility status of Alirajpur: A most backward tribal district of Madhya Pradesh, India. Annals of Plant and Soil Research. 2018;20(1):16–21.
 14. Kumar M, Kar A, Raina P, Singh SK, Moharana PC, Chauhan JS. Spatial variability of available soil nutrients in the Sekhawati region of Thar Desert, India; 2019.
 15. Amara DMK, Patil PL, Kamara MK, Saidu DH. Assessment of soil fertility status using nutrient index approach. Academia Journal of Agricultural Research. 2017;5(2):028-038.
 16. Tagore GS, Sethy SK, Kulhare PS, Sharma GD. Characterization of spatial Variability of micro nutrients in soils: Classical Vs. Geo-Statistical Approach. Communications in Soil Science and Plant Analysis. 2023;54(4):472-487.
 17. Foughifar H, Jafarzadeh AA, Torabi H, Pakpour A, Miransari M. Using geostatistics and geographic information system techniques to characterize spatial variability of soil properties, including micronutrients. Communications in Soil Science and Plant Analysis. 2013;44(8): 1273-1281.
 18. Madhu BM, David AA. Soil fertility status of Jamuna Paar region of Allahabad District, Uttar Pradesh, India. International Journal of Current Microbiology and Applied Science. 2017;6(12):3069-3076.
 19. Sharma RP, Chattaraj S, Vasu D, Karthikeyan K, Tiwary P, Naitam RK, Nimkar AM. Spatial variability assessment of soil fertility in black soils of central India using geostatistical modeling. Archives of Agronomy and Soil Science. 2021;67(7): 876-888.
 20. Singh SS. Crop management under irrigated and rainfed conditions. Kalyani Publication. New Delhi, Ludhiana. 1988; 450.
 21. Dash PK, Mishra A, Saren S, Revathi B, Sethy SK. Preparation of GPS and GIS based soil fertility maps and identification of soil related crop production constraints of soil related crop production constraints of RRTTS and KVK farm, Dhenkanal located in the mid-central table land agro climatic zone of Odisha, India. International Journal of Chemical Studies. 2018;6(5): 934-943.
 22. Sood A, Sharma PK, Tur NS, Nayyar VK. Micronutrient status and their spatial variability in soils of Muktsar district of Punjab—a GIS approach. Journal of the Indian Society of Soil Science. 2009; 57(3):300-306.

23. Kumar MV, Saliha B, Kannan P, Mahendran PP. Delineation and geographic information system (GIS) mapping of soil nutrient status of sugarcane growing tracts of Theni district, Tamil Nadu. *African Journal of Agricultural Research*. 2015;10(33):3281-3291.

© 2023 Bhil et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/103486>