



Comparative Study of the SBA and MOL Methods. Application to Some Systems of Nonlinear Partial Differential Equations

Abdarahman Abakar Himeda ^{a*}, Djibet Mbainguesse ^a,
Bakari Abbo ^a and Youssouf Pare ^b

^aUniversity of N'Djamena, Tchad.

^bLaboratoire d'Analyse Numérique, Informatique et de Biomathématiques (L.A.N.I.BIO), École Normale Supérieure, Burkina Faso.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMCS/2023/v38i121854

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/110122>

Received: 15/10/2023

Accepted: 18/12/2023

Published: 23/12/2023

Original Research Article

Abstract

The paper aims at solving two systems of nonlinear partial differential equations, namely the Fisher-Murray and the Fitz Hugh-Nagumo which are two different mathematical models often used to study ecological and biological phenomena. These systems of equations are solved using the numerical Method of Lines (MOL) and the computed solution are compared with the ones obtained from SBA method (combination of Adomian method, Picard and successive approximation) by means of Matlab routines. The results showed the accuracy and the efficiency of two methods.

*Corresponding author: E-mail: abakarhimedaabdarahman@gmail.com;

J. Adv. Math. Com. Sci., vol. 38, no. 12, pp. 20-29, 2023

Keywords: Mots clés: MOL; systems of nonlinear PDE Fitz Hugh-Nagumo; systems of nonlinear PDE Fisher-murray; Matlab.

2010 Mathematics Subject Classification: 34A05, 34A08, 42A10, 40A30, 65B10.

1 Introduction

System of nonlinear equations arise in many fields of applied mathematics and engineering. The Fitz Hugh-Nagumo and Fisher-Murray systems are two examples of nonlinear systems of equations. The Fisher-Murray system models the propagation of biological limits, the diffusion of a population in an environment. This is derived from a diffusion reaction equation whereas Fitz Hugh-Nagumo model describes neuronal oscillations and excitability. The motivation behind the determination of solution to these systems will lead us to an understanding of the patterns of propagation, developpement and behaviour of population in the case of Fisher-Murray system, as well as for the mode of operation of neuronal oscillations for the Fitz Hugh-Nagumo model. These two models help researchers to predict, understand and control a wide range of natural and biological phenomena. The methods to find their solutions are of fondamental importance. As analytical solution are rarely available, the research of efficient numerical methods are essential. Consider the following system of nonlinear partial differential equations ([1]; [2]; [3]) of type

$$\begin{cases} u_t = k_1 u_{xx} + R(u, u_x, v, v_x), & (x, t) \in \Omega \times [0, T] \\ v_t = k_2 v_{xx} + Q(u, u_x, v, v_x), & (x, t) \in \Omega \times [0, T] \end{cases} \quad (1.1)$$

where $u = u(x, t)$ and $v = v(x, t)$ are dependant variable of independant spatial variable $x \in \Omega = [0, L]$ and temporal $t \in [0, T]$, R and Q are the reaction nonlinear functions of u, u_x, v, v_x . The contants coefficients $k_1 > 0$ and $k_2 > 0$ are the thermal diffusibility of the media. For brevity $u_t = \frac{\partial u}{\partial t}$ and $u_{xx} = \frac{\partial^2 u}{\partial x^2}$. The system of equations (1.1) will be solved on the spatial interval $[0, L]$ subject to boundary conditions for u

$$\begin{cases} u(0, t) = a(t) \\ u_x(L, t) = b(t) \end{cases} \quad (1.2)$$

and for v

$$\begin{cases} v(0, t) = \alpha(t) \\ v_x(L, t) = \beta(t), t \in [0, T] \end{cases} \quad (1.3)$$

and the initial conditions

$$\begin{cases} u(x, 0) = s(x) \\ v(x, 0) = r(x), x \in [0, L] \end{cases} \quad (1.4)$$

The boundary condiions (1.2) and (1.3) give the values of the two solution and their flux at the two ends of space domain as function of t . The equations in (1.4) specify the initial conditions.

We aim at solving numerically this problem, using the Method of Lines(MOL) ([4], [5], [6]; [7], [8]) and compared with SBA method. To solve the nonlinear systems of partial differential equation, we want to transform them into an ordinary differential equation . To achieve this, we need to eliminate the space variable by discretization and retain the time variable, thus creating an ordinary differential equation. The Somé Blaise Abbo(SBA) method ([9], [10], [11], [12]) is an efficient algorithm used by researchers to solve partial differential equations and ordinary differential equations. This method meets the challenges of Adomian polynomial calculations. The basic idea is to see the consistency between the analytical Somé Blaise Abbo method and the numerical lines method, and to analyse which of the two methods provides a less costly solution.

The paper is organized as follows. In section 2 we describe the Method Of Lines. In section 3, we apply these methods for solving two numerical example: the system of non linear equations of FitzHugh-Nagumo and the system of nonlinear equations of Fisher-Murray.

2 Method

We have the choice to discretize both in space and time to obtain a set of nonlinear algebraic equations(AEs) or to discretize only the space derivative and obtaining a set of non linear differential algebraic equations(DAEs). The last approach is retained in this work.The method of lines(MOL) ([13],[14],[15],[16],[9]) is a general way to convert a partial differential equation(PDE) [17];[18] in the form of system of ordinary differential equations(ODE) see [19], [14] [20],[21] The derivatives with respect to the space variables in PDE are discretized to obtain a system of ODEs in time variable ([3]). A suitable ODE solver ([22] [20]) is used for the solution of ODE system. This method is give a very accurate numerical solution for linear and non linear PDE. We define a uniform mesh $0 = x_0 < x_1 < \dots < x_N = L$ with

$$x_i = (i - 1) h, \quad i = 1, 2, \dots, N, \quad h = \frac{L}{N - 1} \quad (2.1)$$

to approximate (1.1) along $x = x_i$ with

$$\begin{cases} u_t(x_i, t) &= k_1 u_{xx}(x_i, t) + R(u(x_i, t), u_x(x_i, t), v(x_i, t), v_x(x_i, t)), \\ v_t(x_i, t) &= k_2 v_{xx}(x_i, t) + Q(u(x_i, t), u_x(x_i, t), v(x_i, t), v_x(x_i, t)), \end{cases} \quad i = 1, 2, \dots, N - 1 \quad (2.2)$$

Let $u_i(t) = u(x_i, t)$ and $v_i(t) = v(x_i, t)$. The equation (1.1) can be dscretized on the uniform mesh (2.1), using the finite difference method [23] [14] [24] – [27] with the central difference approximation to obtain

$$\begin{cases} \frac{du_i}{dt}(t) &= k_1 \frac{u_{i+1}(t) - 2u_i(t) + u_{i-1}}{h^2} + R_i, \quad i = 1, 2, \dots, N - 1 \\ \frac{dv_i}{dt}(t) &= k_2 \frac{v_{i+1}(t) - 2v_i(t) + v_{i-1}(t)}{h^2} + Q_i, \quad i = 1, 2, \dots, N - 1 \end{cases} \quad (2.3)$$

where

$$R_i = R(u_i, \delta p_i, v_i, \delta q_i), \quad Q_i = Q(u_i, \delta p_i, v_i, \delta q_i) \quad (2.4)$$

and

$$\delta p_i = \frac{u_{i+1} - u_{i-1}}{2h}, \quad \delta q_i = \frac{v_{i+1} - v_{i-1}}{2h} \quad (2.5)$$

The boundary conditions (1.2) can be discretized to give

$$\begin{cases} u_0(t) &= a(t) \\ u_{N+1} &= u_{N-1} + 2hb(t) \end{cases} \quad (2.6)$$

and for v

$$\begin{cases} v_0(t) &= \alpha(t) \\ v_{N+1} &= v_{N-1} + 2h\beta(t) \end{cases} \quad (2.7)$$

For initial condition,the discretization of (1.4) give

$$\begin{cases} u_i(0) &= s_i, \quad 1 \leq i \leq N \\ v_i(0) &= r_i, \quad 1 \leq i \leq N \end{cases} \quad (2.8)$$

By introducing (2.6) and (2.7) in (2.3) by taking $i = 1$ and $i = N$, we get

$$\begin{cases} \frac{du_1}{dt}(t) &= \frac{k_1}{h^2} (u_2(t) - 2u_1(t) + a(t)) + R_1 \\ \frac{dv_1}{dt}(t) &= \frac{k_2}{h^2} (v_2(t) - 2v_1(t) + \alpha(t)) + Q_1 \end{cases} \quad (2.9)$$

and

$$\begin{cases} \frac{du_N}{dt}(t) &= \frac{k_1}{h^2} (2u_N(t) + 2u_{N-1}(t) + 2hb(t)) + R_N \\ \frac{dv_N}{dt}(t) &= \frac{k_2}{h^2} (2v_N(t) + 2v_{N-1}(t) + 2h\beta(t)) + Q_N \end{cases} \quad (2.10)$$

The equation

$$\begin{cases} \frac{du_i}{dt}(t) = k_1 \frac{u_{i+1}(t) - 2u_i(t) + u_{i-1}}{h^2} + R_i, i = 2, \dots, N-1 \\ \frac{dv_i}{dt}(t) = k_2 \frac{v_{i+1}(t) - 2v_i(t) + v_{i-1}(t)}{h^2} + Q_i, i = 2, \dots, N-1 \end{cases} \quad (2.11)$$

can be added to others in (2.9) and (2.10).

2.1 Vectoriel and matricial form

We let

$$\mathbf{w} = [u_1(t), u_2(t), \dots, u_{N-1}(t), u_N(t), v_1(t), v_2(t), \dots, v_N(t)]^T \quad (2.12)$$

$$\mathbf{F}(\mathbf{w}) = \begin{bmatrix} \frac{k_1}{h^2} (u_2(t) - 2u_1(t) + a(t)) + R_1 \\ \dots \\ k_1 \frac{u_{i+1}(t) - 2u_i(t) + u_{i-1}}{h^2} + R_i \\ \dots \\ \frac{k_1}{h^2} (2u_N(t) + 2u_{N-1}(t) + 2hb(t)) + R_N \\ \frac{k_2}{h^2} (v_2(t) - 2v_1(t) + \alpha(t)) + Q_1 \\ \dots \\ k_2 \frac{v_{i+1}(t) - 2v_i(t) + v_{i-1}(t)}{h^2} + Q_i \\ \dots \\ \frac{k_2}{h^2} (2v_N(t) + 2v_{N-1}(t) + 2h\beta(t)) + Q_N \end{bmatrix} \quad (2.13)$$

where $2 \leq i \leq N-1$. Using (2.8) as initial condition, and noting by

$$\mathbf{g} = [r_1(t), r_2(t), \dots, r_{N-1}(t), r_N(t), s_1(t), s_2(t), \dots, s_N(t)]^T \quad (2.14)$$

the assciate vectoriel form we then obtain

$$\mathbf{w}(0) = \mathbf{g}, \quad (2.15)$$

The equations (2.9) – (2.11) and (2.15) give the following autonomous system of ordinary differential equation.

$$\begin{cases} \frac{d\mathbf{w}}{dt}(t) = \mathbf{F}(\mathbf{w}(t)), t > 0 \\ \mathbf{w}(0) = \mathbf{g} \end{cases} \quad (2.16)$$

The MOL approximation replaces a PDE system in (1.1) with an initial-value ODE system in (2.16). This ODE system is integrated using a standard routine. In this way, the solution take avantage of the progress in ODE numerical integrators available in Matlab like RK4 or ode15s, ode 23tb, ... for stiff system of ODE.

3 Numerical Experiments

In this section, we solve some examples to show the efficiency of the method of lines and compare the resulting numerical solution with the one obtained by the SBA method.

Example 3.1. The first example consider the FITZ Hugh-Nagumo sytem of equations (1.1) – (1.2)

$$\begin{cases} u_t = \zeta u_{xx} + 2u + v, \\ v_t = \zeta v_{xx} + uv + v^2 \end{cases}$$

where $L = \pi, T = 10, k_1 = k_2 = \zeta, R(u, v) = 2u + v, Q(u, v) = uv + v^2$ with initial conditions

$$\begin{cases} u(x, 0) &= \theta_1 \cos x + \theta_2 \sin x \\ v(x, 0) &= -(\theta_1 \cos x + \theta_2 \sin x) \end{cases}$$

and boundary conditions

$$\begin{cases} u(0, t) &= |\theta_1| \exp(|1 - \zeta| t) \\ v(0, t) &= -|\theta_1| \exp(|1 - \zeta| t) \end{cases}$$

and

$$\begin{cases} u(\pi, t) &= -|\theta_1| \exp(|1 - \zeta| t) \\ v(\pi, t) &= |\theta_1| \exp(|1 - \zeta| t) \end{cases}$$

$$\begin{cases} u(\pi, t) &= -|\theta_1| \exp(|1 - \zeta| t) \\ v(\pi, t) &= |\theta_1| \exp(|1 - \zeta| t) \end{cases}$$

The parameters $\theta_1 = .05; \theta_2 = 1.5; \zeta = 0.5$ are considered for the demonstration The solutions give by the application of SBA method ([9], [10], [11], [12]) give $u(x, t) = |\theta_1 \cos x + \theta_2 \sin x| \exp(|1 - \zeta| t)$ and $v(x, t) = -|\theta_1 \cos x + \theta_2 \sin x| \exp(|1 - \zeta| t)$ wich can be used for the comparison of our method.




Fig. 1. MOL Solution for u and v at a fixed time




Fig. 2. MOL Solution for u




Fig. 3. MOL Solution for v




Fig. 4. SBA Solution for u




Fig. 5. SBA Solution for v

Example 3.2. The second example consider the sytem of equations (1.1) – (1.2) de type Fisher-Murray

$$\begin{cases} u_t = u_{xx} + [u.u_x]_x + u^2 - v^2, \\ v_t = v_{xx} - v_x^2 + u^2 \end{cases}$$

With $L = \pi, T = 10, k_1 = k_2 = 1, R(u, u_x, v, v_x) = [u.u_x]_x + u^2 - v^2, Q(u, u_x, v, v_x) = -v_x^2 + u^2$ with initial conditions

$$\begin{cases} u(x, 0) = \sin x \\ v(x, 0) = \cos x \end{cases}$$

and boundary conditions

$$\begin{cases} u(0, t) = 0 \\ v(0, t) = \exp(-t) \\ \frac{\partial u}{\partial x}(\pi, t) = -\exp(-t) \\ \frac{\partial v}{\partial x}(\pi, t) = 0 \end{cases}$$

The semi-analytical solution computed using SBA method is given by $u(x, t) = \sin x \exp(-t), v(x, t) = \cos x \exp(-t)$ to be compared with MOL numerical method. The different numerical analysis for MOL method has been undertaken by dividing the spatial domain $\Omega = [0, \pi]$, using $N=101$ with $h = \frac{\pi}{N-1}$ and replacing derivatives using the finite difference method for order two. For resulting ODE, we ressort to ODE solver ode15s wich is convenient for stiff problem in the interval $[0, .1]$. The comparison was made by confronting the graph provide by MOL and SBA methods.




Fig. 6. MOL Solution for u and v at a fixed time




Fig. 7. MOL Solution for u




Fig. 8. MOL Solution for v




Fig. 9. SBA Solution for u




Fig. 10. SBA Solution for v

The comparison is made using the graphs from the method of lines and those from the Somé Blaise Abbo method, in fact the graphs represent the semi-analytical solutions and the numerical solutions.

4 Conclusion

This paper investigated MOL method for solving the one-dimensional systems of nonlinear partial differential equations and compared the resulting solution of another semi-analytical SBA method. The method of MOL proceeds in two separate steps. Firstly, spatial derivatives are replaced with finite difference, using finite difference method, finite element method, finite volume method, spectral method and the resulting systems of ordinary differential equations is integrated over time. The availability of high-quality numerical algorithm for solution of stiff system of odes facilitated the computation of the desired results. For our paper we have chosen the finite difference method for discretization in space because of the simplicity to implement in Matlab code and the non-complexity of domain. The use of other spatial discretization methods is not ruled out for future work.

Competing Interests

Authors have declared that no competing interests exist.

References

- [1] Ablowitz MJ, Clarkson PA. Solitons nonlinear evolution equations and inverses scattering. Cambridge Univ. Press, Cambridge; 1991.
- [2] Thambynayagam RKM. The diffusion handbook: Applied solutions for engineers, McGraw-Hill Professional; 2011.
- [3] John Fritz. Partial differential equations. Springer Science. 1991;22.
- [4] Ahmad I, Berzins M. MOL solvers for hyperbolic PDEs with source terms. Math. Comput. Simulat. 2001;56(2):115-125.
- [5] Erdem A, Pamuk S. The method of lines for the numerical solution of mathematical model for capillary formation: The role of tumor angiogenic factor in the extra-cellular matrix. Appl. Math. Comput. 2007;186:891-897.
- [6] Shakeri F, Dehghan M. The method of lines for solution of the one-dimensional wave equation subject to an integral conservation conditions. Comput. and Math. with Appl. 2008;56:2175-2188.
- [7] Sharaf AA, Bakoda HO. A good spatial discretisation in the method of lines. Appl. Math. Comput. 2005;171(2):1253-1263.
- [8] Younes A, Konz M, Fahs M, Zidane A, Huggenberger P. Modelling variable density flow problems in heterogeneous porous media using the method of lines and advanced spatial discretisation methods. Math. Comput. Simulat. 2011;81:2346-2355.
- [9] Some B. Méthode SBA de Résolution des Modèles Mathématiques en environnement. Editions Universitaires Européennes; 2018.
- [10] Bakari A. Nouvel algorithm de Résolution des equations Différentielles Ordinaires(EDO) et des equations aux Dérivées Partielles(EDP) non linéaires, Thèse de Doctorat Unique, Université de Ouagadougou, Burkina Faso; 2007.
- [11] Kamate A, Bationo JY, Djibet M, Paré Y. Analytical solutions of classical and fractional Navier stokes equations by the SBA methods. Journal of Mathematics Research. 2022;14(4).
- [12] Kamaté A, Djibet M, Bationo JY, Bakari A, Youssouf P. Analytical solution of some nonlinear fractional integro-differential equations of Fredholm second kind by a new approximation technique of the numerical SBA method. International Journal of Numerical Methods and Applications; 2022.
- [13] Teschl G. Ordinary differential equations and dynamical systems. American Mathematical Society; 1991.

- [14] Smith GD. Nonlinear solution of partial differential equation. 3rd ed. Oxford University Press, Oxford; 1985.
- [15] Schiesser WE, Griffiths GW. A compendium of partial differential equation model: Methods of lines analysis with matlab. Cambridge University Press; 2009.
- [16] Brezis H, Sibony M. Methode d'approximation et d'Itération pour les opérateurs monotones. Archives for Rational Mechanics and Analysis. 1979;41(4).
- [17] Book DL. Finite difference techniques for vectorised fluid dynamics calculations. Springer, New York; 1981.
- [18] Richtmyer RD, Morton KW. Difference methods for initial values problems. 2nd. ed. Wiley, New York; 1967.
- [19] Ablowitz MJ, Ramany A, Segur H. A connection between nonlinear evolution equations and ordinary differential equations of p type. J. Math. Physique. 1980;21;715-721.
- [20] Schiesser WE. The numerical methods of lines. Academic Press; 1991.
ISSN - 12-624130-9
- [21] Hamdi S, Schiesser WE, Griffiths GW. Methods of lines. Scholarpedia. 2007;2(7)2859.
- [22] Gear CW. Numerical initial value problems in ordinary differential equations. Prentice-Hall, Englewood Cliffs; 1971.
- [23] Mitchell AA, Griffiths DF. Finite difference methods in partial differential equations. Wiley, Chichester; 1980.
- [24] Lambert JD. Computational methods in ordinary differential equations. Wiley, Chichester; 1973.
- [25] Korn GA. Interactive solution of partial differential equations by the methods of lines. Math. Comput. Simulat. 1999;48(1-2):129-138.
- [26] Samarski AA. Some problems in modern theory of differential equations. Differents uravn. 1980;16;121-1228.
- [27] The numerical method of lines. Academic Press. New York. 1991.

© 2023 Himeda et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<https://www.sdiarticle5.com/review-history/110122>