



# Assessment of Soil Particle Distribution, Primary Nutrient Status and Their Response to Environmental Factors Across the Bengal Gram Productive Regions of Sangareddy, Telangana

Venkatesh. B <sup>a\*</sup>, Pavan Chandra Reddy. K <sup>b++</sup>,  
Kiran Reddy. G <sup>c#</sup> and Satish. P <sup>d†</sup>

<sup>a</sup> Department of Soil Science and Agricultural Chemistry, College of Agriculture, Rajendranagar, PJTSAU, Hyderabad, 500030, Telangana, India.

<sup>b</sup> Department of Soil Science and Agricultural Chemistry, AICRP on STCR, Institute of Soil Health Management, Agricultural Research Institute (ARI), PJTSAU, Rajendranagar, Hyderabad, 500030, India.

<sup>c</sup> Department of Soil Science and Agricultural Chemistry, AICRP on Integrated Farming System, Rajendranagar, Hyderabad, 500030, India.

<sup>d</sup> Department of Agronomy-College Farm, College of Agriculture, Rajendranagar, Hyderabad, 500030, India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. 'All authors read and approved the final manuscript.*

## **Article Information**

DOI: <https://doi.org/10.9734/ijecc/2024/v14i54182>

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/116985>

<sup>++</sup> Principal Scientist;

<sup>#</sup> Scientist;

<sup>†</sup> Professor;

\*Corresponding author: E-mail: [bolluvenkatesh97@gmail.com](mailto:bolluvenkatesh97@gmail.com);

**Cite as:** Venkatesh. B, Pavan Chandra Reddy. K, Kiran Reddy. G, & Satish. P. (2024). Assessment of Soil Particle Distribution, Primary Nutrient Status and Their Response to Environmental Factors Across the Bengal Gram Productive Regions of Sangareddy, Telangana. *International Journal of Environment and Climate Change*, 14(5), 205-213. <https://doi.org/10.9734/ijecc/2024/v14i54182>

## ABSTRACT

The present study was carried out to know the fertility status of rabi grown bengal gram soils of Sanga Reddy district, Telangana state by thorough field survey during the year 2022-2023. Total of 150 soil samples (50 samples from each productivity regions) from soil surface (0 - 15 cm depth) across three productivity regions high (592-613 kg acre<sup>-1</sup>), medium (571-592 kg/acre) and low (<571 kg acre<sup>-1</sup>) were collected from the study area before sowing of the crop and analyzed for the soil particle size distribution, available Nitrogen(N), Phosphorous(P) and Potassium(K). Their status was quantified and analytical data was interpreted and statistical parameters like range, mean, standard deviation and coefficient of variation were calculated. The sand, silt and clay content in high, medium and low productivity regions averaged 32.1 %, 26.7%, 41.1 %; 33.7 %, 26.1 %, 40.1 % and 37.4 %, 25.1 %, 37.2 % respectively. The soil available N, P and K showed mean values of 224.3 kg ha<sup>-1</sup>, 209.4 kg ha<sup>-1</sup> and 204.0 kg ha<sup>-1</sup>; 32.0 kg ha<sup>-1</sup>, 28.4 kg ha<sup>-1</sup>, 24.8 kg ha<sup>-1</sup> and 316.4 kg ha<sup>-1</sup>, 311.0 kg ha<sup>-1</sup>, 306.1 kg ha<sup>-1</sup> in high, medium and low productivity regions. Low productivity areas had the largest mean sand concentration, which influenced soil particle aggregation. Silt content varied greatly, particularly in medium and low productivity zones, whereas clay content varied the most in high productivity regions. Available N decreased from high to medium and medium to low production areas. Fluctuations in available P and K showed a wide distribution within each category, as evidenced by large standard deviation and coefficient of variation values across productivity regions, particularly for potassium, emphasising the importance of considering diverse soil conditions and factors influencing potassium availability when developing fertilisation strategies.

*Keywords: Productivity; particle size analysis; primary nutrients; available nitrogen; available phosphorus; available potassium.*

## 1. INTRODUCTION

Soil is undoubtedly one of the most important component of the environment, yet it is perhaps one of the most undervalued, misused and abused of the Earth's resources. To a large extent soils determine the agricultural potential of an area, they influence many geomorphological and hydrological processes. Soil's physical, chemical and biological properties significantly impact its functions, by supplying nutrients for plant growth, regulating water supply, recycling raw materials and serving as a habitat for soil organisms [1].

Soil texture has huge significance for soil development. It can help us to understand the age of the soil and soil development process. Practically, all soil properties and as a whole fertility, depend on it. Soil properties are influenced by texture as, water holding capacity, aeration, susceptibility to erosion, organic matter content, cation exchange capacity (CEC), pH buffering capacity etc. The cultivation, character

and impact of the soil cultivation tools and the fertilization depend to a great extent on the soil texture. It is natural that the soil morphology is determined by its texture. Therefore, the studying of the texture in the field and laboratory conditions is the first necessary stage of research on soil as natural body. Dilkova. [2].

Soil fertility, in addition to physical qualities, influences plant growth it's the concentrations of water, micronutrients, organic and inorganic elements, available nitrogen (N), available phosphorous (P), and available potassium (K). The NPK ratio is a key indicator of crop productivity, quality, and yield that distinguishes balanced from imbalanced fertilisation. Thus, balanced fertiliser application is critical for high crop yields [3]. Soil chemical fertility, particularly nutrient shortage, contributes significantly to soil degradation and limits soil production, stability, and sustainability [4]. The balance of organic and inorganic elements in soil is maintained by nutrient cycling, which is impacted by numerous inputs and losses. Long-term fertiliser trials [5]

revealed decreasing yields when chemical fertilisers are applied continuously, owing to reduced nutrient supplying power and soil fertility, which affects the growing food needs of people worldwide [6], which added for sustainable agricultural production.

Bengal gram is also called as chickpea or gram (*Cicer arietinum*. L) in South Asia and Garbanzo bean in western countries. It is traditionally cultivated in many parts of the world in a wide range of agro-climatic environments. It helps in fixing nitrogen from atmosphere which improves organic matter content in the soil. Bengal gram is a major food legume and important pulse crop. This accounts for about 46.0 percent of the total production of pulses and 65.39 percent of the total world production of bengal gram. Hence, it is known as king of pulse crops.

India is the largest producer of bengal gram followed by Pakistan, Turkey and Iran. India produces around 10 to 11 million tonnes and contributes around 70 percent of the total world production. Bengal gram is the most largely produced pulse crop in India accounting for a share of 40 percent of the total pulse production.

But the bengal gram soil's fertility and quality were being depleted as a result of continuous cropping in areas like Telangana due to prolonged improper management practices that led to the destruction of soil properties because of monoculture, uneven application of inorganic

fertilizer, and excessive reclamation making it challenging to maintain bengal gram productivity. Achieving increased crop productivity, implementing effective soil conservation measures, and fostering sustainable agriculture management require a comprehensive understanding and examination of soil conditions. Therefore, considering the importance of above soil properties and their influence to identify the differences between bengal gram soils with varying productivity, this study has been undertaken to uncover variations in soil properties that help in development of sustainable land use management and the improvement of grain yield in bengal gram cultivated areas.

## 2. MATERIALS AND METHODS

### 2.1 Location of Study Site

Sangareddy district is located in the northern region of the Indian state of Telangana about 50 miles west from Hyderabad. It falls between 17.611629 and 17° 36' 41.8644" N latitude. and 78.081810 and 78° 4' 54.5160" E longitude and lies on 524m above sea level. The district is spread over an area of 4,4645 square kms. Agriculture is the most predominant sector of the district economy. The gross cropped area of the district is 2,65,290 hectares. The study site of bengalgram soil regions of Sangareddy district, Telangana, India, is illustrated in Figs. 1 and 2.




Fig. 1. Location of the study site




Fig. 2. Map of study area

## 2.2 Soil Sample Collection

The present study involved the survey and collection of soil samples from the study area where *rabi* bengal gram was cultivated by dividing the mandals of the district into low (<571 kg acre<sup>-1</sup>), medium (571-592 kg acre<sup>-1</sup>) and high (592-613 kg acre<sup>-1</sup>) productive regions based on the area and yield data gathered from the district agriculture office, Sangareddy. A total of 150 samples were collected, specifically 50 from each productive region from the topsoil layer (0 – 15 cm deep) before sowing of the *rabi* crop, 2022-2023 by stratified random sampling method. These samples were then air-dried, sieved through 2 mm sieve, stored in plastic bags and subjected to further laboratory analysis.

## 2.3 Laboratory Analysis

The collected soil samples were analysed for the parameters *i.e.*, soil texture by Bouyoucos hydrometer method that elucidates the particles size distribution as described by Piper, [7]. The analysis for available nitrogen was carried out by alkaline permanganate method [8], available phosphorus by olsen's method using colorimeter at 660 nm [9] and available potassium by

extraction with neutral normal ammonium acetate using flame photometer [10].

## 3. RESULTS AND DISCUSSION

### 3.1 Computation of Particle Size Analysis in the Three Productivity Regions

In terms of soil physical features, Table 1 and Fig 3 show the particle size distribution of soils classified as clay, clay loamy, gravelly clay, sandy clay loam, and sandy clay, revealing the diversity of the research area's soils. Sand concentration in high-productivity locations varied between 12.4% and 61.1%, with an average of 32.1%, a standard deviation of 9.4, and a coefficient of variation of 29.4%. The range for medium productivity regions was 14.6% to 61.1%, with an average of 33.7%, a standard deviation of 10.2, and a coefficient of variation of 30.2%. The low productivity locations ranged from 20.4% to 57.1%, with an average of 37.5%, a standard deviation of 8.4, and a coefficient of variation of 22.4%.

In terms of silt content, high productivity locations ranged from 9.9% to 34.1%, with an average of 26.7%, a standard deviation of 4.7, and a coefficient of variation of 17.7 percent. The range


for medium productivity regions was 9.9% to 30.8%, with an average of 26.1%, a standard deviation of 5.0, and a coefficient of variation of 19.2%. Low productivity locations showed a range of 5.0% to 31.2%, with an average of 25.1%, a standard deviation of 6.35, and a high coefficient of variation of 28.7%.

deviation of 9.4 and a 23.0% coefficient of variation. In medium productivity areas, the range was 28.9% to 55.8%, averaging at 40.1% with a standard deviation of 8.2 and a 20.6% coefficient of variation. Low productivity regions displayed a range of 19.9% to 62.0%, with an average of 37.2%, standard deviation of 7.94 and a 20.3% coefficient of variation. Fig 3 provides a graphical representation of distribution of soil particles in three productive regions.

On the examination of clay content, high productivity regions showcased a range of 24.5% to 69.7%, averaging at 41.1% with a standard

**Table 1. Descriptive statistics of particle size analysis among the three Bengal gram productivity regions**

<b>High Productivity Regions (n = 50)</b>					
<b>Soil Property</b>	<b>Minimum</b>	<b>Maximum</b>	<b>Mean</b>	<b>SD</b>	<b>CV</b>
Sand (%)	12.40	61.12	32.14	9.46	29.43
Silt (%)	9.97	34.10	26.73	4.75	17.75
Clay (%)	24.56	69.70	41.13	9.49	23.07
<b>Medium Productivity Regions (n = 50)</b>					
<b>Soil Property</b>	<b>Minimum</b>	<b>Maximum</b>	<b>Mean</b>	<b>SD</b>	<b>CV</b>
Sand (%)	14.60	61.12	33.73	10.22	30.29
Silt (%)	9.97	30.80	26.15	5.04	19.26
Clay (%)	28.90	55.80	40.11	8.27	20.62
<b>Low Productivity Regions (n = 50)</b>					
<b>Soil Property</b>	<b>Minimum</b>	<b>Maximum</b>	<b>Mean</b>	<b>SD</b>	<b>CV</b>
Sand (%)	20.40	57.15	37.48	8.41	22.44
Silt (%)	5.00	31.20	25.12	6.35	28.71
Clay (%)	19.90	62.00	37.20	7.94	20.39




**Fig. 3. Particle size distribution among the three productivity Bengal gram regions**



**Table 2. Descriptive statistics of primary nutrients among the three bengalgram productivity regions**

<b>High Productivity Regions (n = 50)</b>					
<b>Soil Property</b>	<b>Minimum</b>	<b>Maximum</b>	<b>Mean</b>	<b>SD</b>	<b>CV</b>
Available Nitrogen (kg ha <sup>-1</sup> )	170.60	307.33	224.34	31.42	14.01
Available Phosphorus (kg ha <sup>-1</sup> )	12.38	55.13	32.09	11.28	35.14
Available Potassium (kg ha <sup>-1</sup> )	196.88	434.25	316.49	55.27	17.46
<b>MEDIUM PRODUCTIVITY REGIONS (n = 50)</b>					
<b>Soil Property</b>	<b>Minimum</b>	<b>Maximum</b>	<b>Mean</b>	<b>SD</b>	<b>CV</b>
Available Nitrogen (kg ha <sup>-1</sup> )	155.55	292.28	209.40	28.12	13.43
Available Phosphorus (kg ha <sup>-1</sup> )	12.60	52.67	28.44	8.55	30.07
Available Potassium (kg ha <sup>-1</sup> )	198.0	423.00	311.01	59.79	19.22
<b>Low Productivity Regions (n = 50)</b>					
<b>Soil Property</b>	<b>Minimum</b>	<b>Maximum</b>	<b>Mean</b>	<b>SD</b>	<b>CV</b>
Available Nitrogen (kg ha <sup>-1</sup> )	111.64	265.93	204.09	37.34	18.30
Available Phosphorus (kg ha <sup>-1</sup> )	14.25	48.58	24.85	8.42	33.89
Available Potassium (kg ha <sup>-1</sup> )	121.50	417.38	306.15	59.57	19.46

\*SD – Standard Deviation, CV – Coefficient of Variation



**Fig. 4. Depiction of primary nutrients across three bengal gram productivity regions**

Thus, it can be inferred that soils of three regions contributed to their unique composition. Sand content varied across productivity regions, with the highest mean observed in low productivity areas. Silt content varied significantly, with the highest coefficient of variation observed in medium and low productivity regions. Clay content displays variations, with the highest mean observed in high productivity regions and a notable coefficient of variation in low productivity areas. High productivity regions tend to have lower mean sand and high silt, clay contents compared to the other two, favouring the aggregation of soil particles. The similar results

were observed in high, medium and low productive paddy soils in the research conducted by Liu et al., [11]. Particle size distribution that helps in the determination of soil texture, which in turn shows its impact on root production and, consequently, water and nutrient uptake. Notably, clay soil proves advantageous for higher grain yield due to its ability to retain water and nutrients, resulting in heavier seeds, and improved grain filling compared to sandy soil [12]. The variability in particle size distribution across different productivity levels emphasizes the importance of soil aggregation in the effective agricultural management.

## 3.2 Computation of Primary Nutrients in the Three productivity Regions

### 3.2.1 Available nitrogen

Nitrogen (N) is a major fertiliser for agriculture and food production. Nitrogen (N) is often the most limiting factor in crop production. Hence, application of fertilizer nitrogen results in higher biomass yields, protein yield and concentration in plant tissue is commonly increased. Information about nitrogen (N) availability in soil is important for optimizing N fertilizer recommendations.

In high-productivity zones, available nitrogen (N) in the current study (Table 2) ranged from 170.60 to 307.33 kg ha<sup>-1</sup>, with a mean of 224.34 kg ha<sup>-1</sup>, standard deviation of 31.42, and coefficient of variation of 14.01%. In medium productivity regions, the range was 155.55 to 292.28 kg ha<sup>-1</sup> with a mean of 209.40 kg ha<sup>-1</sup>, with standard deviation and coefficient of variation of 28.12 and 13.43%, respectively, and in low productivity regions, the range was 111.64 to 265.93 kg ha<sup>-1</sup> with a mean of 204.09 kg ha<sup>-1</sup>, with standard deviation of 37.34 and coefficient of variation 18.30%. The mean available N value followed a decreasing trend from high to medium and medium to low productivity regions whereas the standard deviation and coefficient of variation values indicated a moderate level of variability in N levels across the three productivity regions, with the highest variability in high and medium productivity areas. The categorization suggests a low to medium level of nitrogen availability in the soils across the studied regions.

### 3.2.2 Available phosphorus

Phosphorus, as a macronutrient, is required for the conversion of sugars and starches, photosynthesis, energy transmission, and nutrient movement inside plants. In the current study, available P in high productivity locations ranged from 12.38 to 55.13 kg ha<sup>-1</sup>, with a mean of 32.09 kg ha<sup>-1</sup>, a standard deviation of 11.28, and a coefficient of variation of 35.14%. The available phosphorus in medium productivity regions ranged from 12.60 to 52.67 kg ha<sup>-1</sup>, with a mean of 28.44 kg ha<sup>-1</sup>, a standard deviation of 8.55, and a coefficient of variation of 30.07%. In low production zones, the range fluctuated between 14.25 to 48.58 kg ha<sup>-1</sup>, with a mean of 24.85 kg ha<sup>-1</sup>, standard deviation of 8.42 and coefficient of variation 33.89%, respectively.

(Table 2). These outcomes are in line with the findings of Sannidi et al., [13]. This observed variability in available P levels emphasizes the importance of agricultural management practices based on the specific phosphorus dynamics within each productivity category.

### 3.2.3 Available potassium

Potassium (K) is the second most abundant nutrient in plant photosynthetic tissues after nitrogen. It plays a fundamental role in plant function, especially in water-use efficiency and economy. It plays a particularly crucial role in a number of physiological processes vital to growth, yield, quality, and stress resistance of crops [14,15].

The available K concentration in soils of high productivity locations ranged from 196.88 to 434.25 kg ha<sup>-1</sup>, with a mean of 316.49 kg ha<sup>-1</sup>, and standard deviation and coefficient of variation of 55.27 and 17.46%, respectively. Medium productivity zones have available K levels ranging from 198.0 to 423.00 kg ha<sup>-1</sup>, with a mean of 311.01 kg ha<sup>-1</sup>, 59.79 standard deviation, and 19.22% coefficient of variation. Similarly, low productivity locations displayed K levels ranging from 121.50 to 417.38 kg ha<sup>-1</sup>, with a mean of 306.15 kg ha<sup>-1</sup>, having standard deviation and coefficient of variation of 59.57 and 19.46%, respectively (Table 2) and are categorised from medium to high similar to the findings reported by Vasu et al., [16]. The Fig 4. provides a graphical representation of values of primary nutrients in the three productive regions. The large standard deviation and coefficient of variation values across productivity regions indicate a wide range of K levels within each group. Soil composition, weathering, and other environmental factors may all contribute to this heterogeneity. When adopting fertilisation techniques, farmers and agronomists should take into account the variety of soil conditions and factors impacting potassium availability.

## 4. CONCLUSION

The results showed the variation in sand, silt, and clay concentration, highlighting the varied soil compositions that contribute to the specific character of each productivity region. Notably, low productivity areas had the largest mean sand concentration, which

influenced soil particle aggregation and, thus, crop yield. Silt content varied significantly, especially in medium and low production zones, whereas clay content varied, with the greatest mean in high productivity areas, promoting soil particle aggregation. The study area's soils had a low to medium level of nitrogen availability, which declined from high to medium to low productivity regions. The observed changes in available phosphorus levels underline the importance of agricultural management systems that account for phosphorus dynamics within each productivity category. Potassium levels appear to be widely dispersed within each category, as evidenced by high standard deviation and coefficient of variation values across the productivity regions. When developing fertilisation methods, farmers and soil scientists should consider a wide range of soil conditions and factors that influence potassium availability. Particle size distribution, which aids in the measurement of soil texture, influences water and nutrient uptake, resulting in an impact on root formation and, consequently yield.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Vasu D, Singh SK, Ray SK, Duraisami VP, Tiwary P, Chandran P, Nimkar AM, Anantwar SG. Soil quality index (SQI) as a tool to evaluate crop productivity in semi-arid Deccan plateau, India. *Geoderma*. 2016b; 282:70-79.
2. Dilkova R. Physical properties of the soil in: Lectures of the soil science, Project of FAO - Bulgaria TCP/BUL/4502(T), Sofia. 1998; 231-248.
3. Tale, K.S and Ingole, S. A review on role of physico-chemical properties in soil quality. *Chemical Science Review and Letters*. 2015;4(13):57-66.
4. Hartemink AE. Land use change in the tropics and its effect on soil fertility. In *Proceedings 19th World Congress of Soil Science*, Brisbane, Australia. 2010; (55-58).
5. Ladha JK, Dawe D, Pathak H, Padre AT, Yadav RL, Singh B, Singh Y, Singh Y, Singh, P, Kundu AL, Sakal R. How extensive are yield declines in long-term rice–wheat experiments in Asia? *Field Crops Research*. 2003;81(2-3): 159 -180.
6. Shahid M, Nayak AK, Shukla AK, Tripathi R A, Kumar S, Mohanty P, Bhattacharyya R. Raja, Panda BB.. Long-term effects of fertilizer and manure applications on soil quality and yields in a sub-humid tropical rice-rice system. *Soil Use and Management*. 2013;29(3):322–32.
7. Piper CS. *Soil and plant analysis*. Hans Publishers, Bombay 1966;137-153.
8. Subbaiah BV, Asija GL. A rapid procedure for the estimation of available nitrogen in soils. *Current Science*. 1956;25: 259-260.
9. Olsen SR, Cole CV, Watanabe FS, Dean LA. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. *USDA Circular*. Washington. 1954;939.
10. Jackson ML. *Soil chemical analysis*. Prentice-Hall of India Private Limited, New Delhi; 1973.
11. Liu Z, Zhou W, Shen J, Li S, Ai C, Soil quality assessment of yellow clayey paddy soils with different productivity. *Biology and Fertility of Soils*. 2014;50: 537-548.
12. Dou, F., Soriano, J., Tabien, R.E. and Chen, K. Soil texture and cultivar effects on rice (*Oryza sativa*, L.) grain yield, yield components and water productivity in three water regimes. *PloS one*. 2016;11(3): e0150549.
13. Sannidi S, Bindu GM, Neelima TL, Umadevi M. Soil quality mapping in the groundnut belt of erstwhile Mahabubnagar District, Telangana, India using GIS. *Current Science*. 2022;122(5): 600.
14. Sekhon GS. Potassium in Indian soils and crops. *Proceedings of the Indian National Science Academy*. 1999;B65 (3&4):83-108.
15. Srinivasarao Ch, Vittal KPR, Tiwari KN, Gajbhiye PN, Kundu S, Pharande AL., Reddy YY, Shankar MA, Potassium supplying characteristics of twenty-one soil profiles under diverse rainfed production systems. *Journal of the Indian Society of Soil Science*. 2007b;55: 14-22.



16. Vasu, D., Singh, S.K., Duraisami, V.P., Jangir, A and Butte, P.S. Fertility status of cotton (*Gossypium hirsutum* L.) growing soils in Thimmajipet mandal, Mahabubnagar district, Telangana. *Andhra Pradesh Journal of Agricultural Sciences*. 2016a;2(1): 10-15.

---

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/116985>