



# Evaluation of Antimicrobial Efficacy of Hydrazide-Hydrazone Derivatives against Food Borne Pathogens

**Zébré Arthur Constant <sup>a</sup>, Tiekoura Konan Bertin <sup>b</sup>,  
Coulibaly Bakary <sup>a\*</sup>, Coulibali Siomenan <sup>c</sup>,  
N'dri N'Guessan Alexandre <sup>a</sup>, Konaté Ibrahim <sup>a</sup>,  
Koffi-Nevry Rose <sup>d</sup> and Connil Nathalie <sup>e</sup>**

<sup>a</sup> Department of Biochemistry-Microbiology, Agrovalorisation Laboratory, University Jean Lorougnon Guédé, Daloa, Côte d'Ivoire.

<sup>b</sup> Department of Bacteriology-virology, National Antibiotic Reference Centre, Institut Pasteur, Côte d'Ivoire.

<sup>c</sup> Department of Sciences of Structures of Matter and Technology, Laboratory of Constitution and Reaction of Matter, Félix Houphouët-Boigny University, Abidjan, Côte d'Ivoire.

<sup>d</sup> Laboratory of Biotechnology and Food Microbiology, Department of Food Science and Technology, Nangui Abrogoua University, Abidjan, Côte d'Ivoire.

<sup>e</sup> Research Unit-Bacterial Communication and Anti-infectious Strategies (CBSA, UR4312), Rouen Normandie University, Evreux, France.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: <https://doi.org/10.9734/mrji/2024/v34i71455>

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/117522>

**Original Research Article**

**Received: 28/03/2024**

**Accepted: 02/06/2024**

**Published: 08/06/2024**

\*Corresponding author: E-mail: [benipoho7@gmail.com](mailto:benipoho7@gmail.com);

## ABSTRACT

Bacteria are a real public health problem because of their involvement in many diseases. Their resistance to antibiotics has become one of the most important problems in the fight against infectious diseases worldwide. The objective of this work was to evaluate the antibacterial activity of hydrazide-hydrazone derivatives against pathogenic strains isolated from food using the agar diffusion method in cups and the macro-dilution method in liquid medium.

In this work, a survey was conducted among food vendors on the four security corridors of the city of Daloa. A structured questionnaire was administered to 127 vendors and a total of 60 food samples were collected from the four corridors and analysed.

The results indicate that the majority of food sales at these security corridors are carried out by women (88.12%). The vendors were mostly adults (59.84%) aged between 15 and 35, with no formal education (50.39%). Microbiological analysis of the 2 types of food showed the presence of enterobacteria above the recommended thresholds. Concerning the antimicrobial effectiveness explored, four substances (AEV5, AB2, AB3 and AB5) revealed antibacterial activity. The MBC/MIC ratio of all these substances being less than 4 indicates that these synthesised substances all have bactericidal activity. As a result, substances derived from S-alkylation and Mannich bases can be used to combat bacterial infections.

**Keywords:** *Biological properties; hydrazide-hydrazone derivatives; food poisoning; bacterial strains; enterobacteria; food; Daloa.*

## 1. INTRODUCTION

Food is the first consumer good used to satisfy the primary needs of every human being. The quality of this food is vital to the health of consumers [1]. Food is the main source of microorganisms when it is not eaten in a healthy way. Every year, several million people suffer from food-borne illnesses [2,1]. These food-borne illnesses are generally known as food poisoning. They are diseases that are often infectious and accidental, contracted as a result of ingesting food contaminated with pathogenic bacteria. Such contamination generally results from inadequate food handling, preservation or cooking methods [3]. These infectious diseases are a global concern. The number of cases of food poisoning is constantly rising around the world. Every year, there are more than 600 million cases of food poisoning in the world [4,5]. There have been numerous cases of food poisoning in Côte d'Ivoire. In May 2022, in the town of Bondoukou, in the east of the country, around a hundred people aged between 2 and 32 suffered from food poisoning after eating a dairy product sold in an ice cream parlour [6]. Food-borne diseases are therefore a major concern for governments [7]. Most are caused by bacteria such as *Listeria monocytogenes*, *Escherichia coli*, *Staphylococcus aureus*, *Salmonella* spp., *Clostridium* spp. and *Campylobacter* spp [8,9]. Antibiotics are most commonly used to treat this condition. Unfortunately, the misuse of antibiotics and the horizontal transfer of resistance genes have led to the emergence of bacteria resistant to certain

antibiotics. Among the concerns raised by this issue, bacterial resistance to antibiotics has been recognised by the World Health Organisation as one of the most serious threats to global health, food security and development. Its spread could change medicine in the coming years. Antibiotic resistance is all the more worrying given that the use of antibiotics is often uncontrolled [10]. Faced with the constant threat of infectious agents, it is essential to find new antimicrobial molecules. Hydrazide-hydrazone derivatives could offer an alternative for better management of cases of food poisoning [11]. Indeed, these substances are a potential source for strengthening the fight against bacteria that are increasingly resistant to antibiotics. The main objective of this work is to evaluate the antibacterial activity of synthesised hydrazide-hydrazone derivatives on strains isolated from street foods sold in the commune of Daloa.

The specific objectives of the present study were: to identify the presence of germs responsible for food poisoning in foods sold on the four security corridors of the city of Daloa, highlight the antibacterial effect of hydrazide-hydrazone derivatives on pathogenic strains and determine the antibacterial parameters (MIC, CMB) of these substances.

## 2. MATERIALS AND METHODS

### 2.1 Materials

#### 2.1.1 Presentation of the study area

Daloa is the capital of the Haut Sassandra region. Its position at the crossroads of the roads

linking respectively the West and the North-West to the rest of the country, on the one hand, to neighboring Guinea and Liberia to the West, to Mali and Burkina-Faso in the North, on the other hand, gives it the status of a crossroads city [12].

## 2.2 Biological Material

### 2.2.1 Bacterial strains

The microbial support consists of:

- ❖ Bacterial strains isolated and identified from foodstuffs sold on the various corridors. A total of six (6) strains were selected. These were four strains of *E.coli* coded *E.coli* A1, *E.coli* A2, *E.coli* A3, *E.coli* A4 and two strains of Enterobacteriaceae coded *Ent* P1, *Ent* P2.
- ❖ A strain of *E. coli* resistant to third generation cephalosporins (C3G ®) from the bio-collection of the National Reference Center for Antibiotics of the Institute Pasteur (Abidjan, Côte d'Ivoire) used for quality control of the tests.

**Hydrazide-hydrazone derivatives and mannich bases:** 4 codified substances (AEV3, AEV5, AEV6, AEV8) derived from benzimidazole S-alkylation and 5 codified mannich base substances (AB2, AB3, AB4, AB5 and AB7) synthesised and described as new bioactive molecules were provided by the Laboratory of Constitution and Reaction of Matter, Department of Sciences of Structures of Matter and Technology, Félix Houphouët-Boigny University, Abidjan, Côte d'Ivoire for this study.

### 2.3 Methods

A socio-demographic survey of 127 sellers on the different security corridors (Man, Issia, Bouaflé and Vavoua) was carried out over a period of one month (from March 31 to April 30, 2022). The survey sought information on the age, gender and level of education of the vendors. It also looked at the hygienic state of food preparation, conditions of sale, conditions of food preparation and storage, and the level of knowledge about hygiene practices among male and female vendors.

### 2.4 Sample Collection and Transport

Food samples were collected from four security corridors in the city of Daloa, namely the Issia corridor (South), the Vavoua corridor (North-West), the Bouaflé corridor (East) and the Man

corridor (West). A total of 30 solid food samples and 30 liquid food samples were collected from vendors. The samples were packaged in labelled sterile jars, then placed in a cooler containing an ice accumulator and transported to the laboratory.

**Isolation and identification of bacteria from the food samples:** The preparation of the mother suspension of samples and decimal dilutions was carried out in accordance with standard NF EN ISO 6887-1, 2017, which defines the general rules for the preparation of the mother suspension and dilutions. Culture media were prepared and used in accordance with the manufacturers' instructions. Two types of inoculation were carried out. Surface inoculation for the detection of *E.coli* on Rapid' *E. coli* medium (Bio-Rad, France) and inoculation in the mass with VRBL (Biokar, France) and VRBG (Biokar, France) media for the detection of Enterobacteriaceae. Cultures were incubated at 37°C for 24 h. Colonies were observed and counted, identified and preserved in 10% glycerol.

### 2.5 Preparation of Solutions of Hydrazide-Hydrazone Derivatives and Mannich Bases

Stock solutions of the chemical substances were prepared at a concentration of 16 µg/mL. The substances in powder form were first diluted in ethanol at 96°C.

### 2.6 Sensitivity Testing and Determination of Inhibition Diameters

The Mueller-Hinton agar diffusion technique described by Bssaibis et al. [13] was used in this study. This technique involved obtaining log phase colonies transplanted onto nutrient agar using the streak method. A bacterial inoculum estimated at 10<sup>6</sup> CFU/mL adjusted by opacity to 0.5 McFarland was prepared in physiological water (NaCl 0.9%). of this inoculum was then swabbed onto Muller-Hinton agar (GMH, Bio-Rad, France). Cups were made by sinking the large end of a sterile Pasteur pipette into the inoculated Muller-Hinton agar and left to dry. These plates were then filled with 50 µL of the chemical preparation and incubated at 37°C for 24 h. The inhibition diameters were then determined by caliper measurement. The effectiveness of the substances was assessed according to the criteria of Ponce et al. [14].

Thus, a substance is said to be ineffective if the inhibition diameter is less than 8 mm, effective if the diameter is between 9 and 14 mm, very effective for a diameter between 15 and 19 mm and extremely effective if the diameter is greater than 20 mm.

## **2.7 Determination of Antibacterial Parameters Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC)**

The determination of the antibacterial parameters was made for substances with an inhibition diameter greater than 9 mm according to the macro-dilution method in a liquid medium described by [15].

## **2.8 Preparation of the bacterial inoculum**

A bacterial inoculum was prepared from a fresh colony in 10 mL of sterile MH broth and incubated at 37°C for 3 h. Then 0.3 mL of this pre-culture was diluted in 10 mL of sterile MH broth and homogenised.

## **2.9 Preparation of Test Concentration Ranges**

A concentration range for the test substances was obtained using the double dilution method. A stock solution with a concentration of 1000 µg/mL (2mg in 2ml of 70% ethanol) was prepared. A series of 2-fold dilutions were made from this solution to obtain concentration ranges from 500 µg/mL to 7.81 µg/mL.) Then, in sterile haemolysis tubes, 1 mL of each concentration range of the substance was brought into contact with 1 mL of bacterial inoculum. In a growth control tube, 1 mL of sterile distilled water was added to the inoculum, while 2 mL of sterile Muller-Hinton broth (MHB) was added to the sterility control. The preparations were incubated for 24 h at 37°C. Afterwards, observation with the naked eye was carried out and the lowest concentration for which no bacterial growth was observed with the naked eye corresponded to the MIC.

The MBC is determined by transferring a sample of the tubes showing no growth to a fresh medium devoid of the molecules to be tested. This involved diluting the starting inoculum from  $10^{-1}$  to  $10^{-4}$  and inoculating these different dilutions using a calibrated 2 µL loop in 5 cm long streaks on Muller-Hinton agar, then incubating at 37°C for 24 h (preparation A). After

reading the MICs, the contents of tubes in which there was no visible growth were used to inoculate the GMH on 5 cm long streaks (preparation B). The MIC was determined by comparing the bacterial growth of preparations A and B. The smallest concentration in the tube with less than 0.01% viable bacteria compared to the initial inoculum is the MBC.

The CMB/CMI report made it possible to clarify the mode of action of the chemical substance.

According to Nayaka et al. [16], the substance is bactericidal when its CMB is equal to its MIC or if the CMB/MIC ratio is less than or equal to 4. It is said to be bacteriostatic when its CMB is greater than its MIC or if the CMB/MIC ratio is higher than 4. When this ratio is equal to 32, the strain is said to be tolerant.

## **2.10 Statistical Analysis**

All tests were carried out in triplicate and the numerical values obtained are expressed as the arithmetic mean plus the corresponding statistical standard deviation. STATISTICA 7.1 software was used for these analyses. Analyses of variance (ANOVA) based on multiple DUNCAN tests with a significance level of 5% were performed in order to compare the variables measured on different strains with the substances tested. Means, standard deviations and graphs were produced using EXCEL 2016.

## **3. RESULTS**

### **3.1 Socio-demographic Profile of the People Surveyed and their Level of Knowledge About Hygiene**

The results of the survey of food vendors from the four corridors are shown in Table 1. In terms of gender, the results show that 11.82% of vendors are men and 88.12% are women. In terms of level of education, vendors are poorly educated. Only 7.08% had completed higher education, 27.55% had attended secondary school, 14.96% had completed primary school, and 50.39% had no education at all. In terms of age, 15.74% of sellers are under 14, 59.84% are between 15 and 35, and 24.42% are over 36. In terms of knowledge of hygiene, the results show that 54.5% of respondents have some knowledge of hygiene and 45.5% have no knowledge of hygiene.

### 3.2 Antibacterial Effect of Chemicals Derived from Benzimidazole S-Alkylation on Bacterial Strains

Table 2 shows the inhibition diameters of the four benzimidazole S-alkylation substances (AEV3, AEV5, AEV6 and AEV8) on strains isolated from food. It should be noted that the four substances were ineffective on the 4 *E. coli* strains isolated because they had an inhibition diameter of less than 9 mm. The substances AEV5 and AEV8 were highly effective and effective respectively only on the *Ent P2* strain with diameters of 17.3 mm and 10 mm. On the *Ent P2* strain, all the substances in this chemical group were found to have no antibacterial effect.

### 3.3 Effect of Various Mannich-Based Chemicals on the *In Vitro* Growth of Bacterial Strains

The results of the sensitivity of bacterial strains to Mannich-based substances are presented in Table 3. All the substances were ineffective on

the *Ent P2* strain. No zone of inhibition was observed. The results also show that the substances AB2, AB3, AB4 and AB7 are extremely effective on the *E. coli* A2 strain with diameters between 21 and 27 mm and similarly for AB2 and AB3 on the *E. coli* A1 strain with diameters of  $21.0 \pm 0.0$  and  $23.0 \pm 0.0$  respectively. The largest inhibition zone diameters were observed on *E. coli* A2 ( $27.0 \pm 0$  mm) for AB2 (Table 3).

### 3.4 Antibacterial Parameters of Synthesised Chemicals

Table 4 presents the results of the determination of antibacterial parameters (MIC, MBC and MBC/MIC) of the substances that showed an inhibitory action on the bacterial strains by their inhibition diameter. These were AEV5 and AEV8 on the *Ent P1* strain, AB2 on *E. coli* A2, AB3 on *E. coli* A1 and AB5 on *Ent P1*. The MBC/MIC ratio for all the substances was less than 4. The substances all have bactericidal power on the bacterial strains used.

**Table 1. Socio-demographic profiles of vendors on the various corridors**

Socio-demographic profiles		Staff	Frequency (%)
Sex	Male	15	11.82
	Female	112	88.12
Age	0-14 years	20	15.74
	15-35 years	76	59.84
	35 years and over	31	24.42
Level of study	No level of education	64	50.39
	Primary	19	14.96
	Secondary	35	27.55
	Superior	9	7.08
Level of hygiene knowledge	Yes	69	54.5
	No	58	4.5

**Table 2. Diameters\* of the zones of inhibition of chemical substances derived from benzimidazole S-alkylation on bacterial strains**

Bacterial strains	Inhibition diameters (mm)			
	S-alkylation benzimidazole			
	AEV3	AEV5	AEV6	AEV8
<i>E. coli</i> A1	0,8±0,1 <sup>b</sup>	5,3±0,6 <sup>a</sup>	0,6±0,2 <sup>b</sup>	0,5±0,0 <sup>b</sup>
<i>E. coli</i> A2	0,7±0,2 <sup>a</sup>	0,8±0,1 <sup>a</sup>	0,7±0,2 <sup>a</sup>	0,7±0,0 <sup>a</sup>
<i>E. coli</i> A3	0,7±0,1 <sup>b</sup>	7,2±0,6 <sup>a</sup>	0,7±0,2 <sup>b9</sup>	0,5±0,0 <sup>b</sup>
<i>E. coli</i> A4	0,7±0,2 <sup>a</sup>	0,7±0,1 <sup>a</sup>	0,6±0,2 <sup>a</sup>	0,9±0,0 <sup>a</sup>
<i>Ent P1</i>	0,6±0,2 <sup>c</sup>	17,3±0,6 <sup>a</sup>	0,8±0,1 <sup>c</sup>	10±0,0 <sup>b</sup>
<i>Ent P2</i>	0,8±0,1 <sup>a</sup>	0,6±0,2 <sup>a</sup>	0,7±0,2 <sup>a</sup>	0,5±0,1 <sup>a</sup>

\*mean ± SD

NB: Values with the same letter in superscript are statistically identical, at the 5% threshold

**Table 3. Diameters\* of the zones of inhibition of Mannich-based substances on bacterial strains**

Bacterial strains	Inhibition diameters (mm)				
	Mannich-based				
	AB <sub>2</sub>	AB <sub>3</sub>	AB <sub>4</sub>	AB <sub>5</sub>	AB <sub>7</sub>
<i>E. coli</i> A1	21,2±0,8 <sup>b</sup>	23,0±0,0 <sup>a</sup>	—	18,3±0,5 <sup>c</sup>	13,0±1,6 <sup>d</sup>
<i>E. coli</i> A2	27,9±1,4 <sup>a</sup>	24,5±2,5 <sup>b</sup>	21,2±0,8 <sup>c</sup>	16,3±0,2 <sup>d</sup>	22,6±1,4 <sup>bc</sup>
<i>E. coli</i> A3	17,2±0,7 <sup>b</sup>	17,0±0,2 <sup>b</sup>	12,3±0,7 <sup>c</sup>	19,0±0,0 <sup>a</sup>	—
<i>Ent</i> P1	17,0±1,3 <sup>a</sup>	15,3±0,7 <sup>a</sup>	14,0±0,8 <sup>a</sup>	19,0±1,2 <sup>a</sup>	10,0±0,8 <sup>a</sup>
<i>Ent</i> P2	—	—	—	—	—

\*mean ± SD

NB: Values with the same letter in superscript are statistically identical, at the 5% threshold

**Table 4. Antibacterial parameters of benzimidazole S-alkylation derivatives and Mannich bases**

S-alkylation benzimidazole	Mannich-based				
	AEV5	AEV8	AB2	AB3	AB5
<b>Bacterial strains</b>	<i>Ent</i> P1	<i>Ent</i> P1	<i>E. coli</i> A2	<i>E. coli</i> A1	<i>Ent</i> P1
MIC (µg/ mL)	31,25	31,25	125	62,5	62,5
MBC (µg/ mL)	62,5	62,5	250	62,5	125
MBC/MIC	2	2	2	1	2
Effect	Bactericidal	Bactericidal	Bactericidal	Bactericidal	Bactericidal

#### 4. DISCUSSION

The aim of this work was to evaluate the antibacterial activity of the chemical substances synthesised on bacterial strains responsible for food poisoning. The strains were isolated from food sold on the various security corridors in the town of Daloa. The socio-demographic survey on the sale of food at these various corridors revealed that this activity is mainly carried out by women (88.12%), compared with 11.82% for men. Vendors are generally very young, with 59% aged between 15 and 35, and the vast majority have no formal education. These same observations have been made by Khomotso and Chelule [17] and also by Attien et al. [18], in fact the commercial activity of street food in Africa is mostly carried out by women and especially that are linked to catering. Actually, according to Rahman et al [19], the predominance of women in the street food sector could be due to the fact that women are the main breadwinners in the majority of households. The results of the microbiological analysis of the various foodstuffs revealed the presence of microorganisms such as: enterobacteria, faecal coliforms and *E. coli*. These germs have also been found in street foods in the work of certain authors such as Mosupye and Von Holy [20], Attien et al. [21] and

Zébré et al [22]. The presence of these germs in these foods is thought to be due to handling by the vendors. These germs are indicators of poor hygiene. They are evidence of faecal contamination [23]. Hygiene conditions and the practices of food handlers in sales outlets are generally inadequate [21,22]. A lack of understanding of the concept and importance of hygiene is thought to be behind the high levels of contamination in the various samples analysed [24,18]. Consumer health is therefore becoming a global concern. In order to combat food-borne illnesses effectively in the face of the growing threat of bacteria resistant to conventional antibiotics, new chemical molecules derived from benzimidazole S-alkylation and Mannich bases have been synthesised and tested on bacteria isolated from food. The results obtained with the benzimidazole S-alkylation substances show that only the *Ent* P1 strain is sensitive to the AEV5 and AEV8 substances. This observation is less surprising given that enterobacteria are highly resistant to antimicrobials [25,26]. According to Zniber et al. [27], the lack of effect could be explained by taking into account the structure of the molecule tested (the size of the molecules) as well as the physiology and genetic potential of the target cell. But according to Iredell et al. [25],

Enterobacteriaceae resistance to antibiotics, particularly  $\beta$ -lactam antibiotics, is increasingly dominated by the mobilisation of single continuously expressed genes that code for effective drug-modifying enzymes. As for Mannich-based substances, they have almost all shown antibacterial activity on all strains with the exception of the *Ent* P2 strain. The antibacterial activity of these substances has been demonstrated in several previous studies. Indeed, this family of substances is effective on both Gram+ and Gram- bacteria [28]. Determination of the antibacterial parameters shows that the substances have an antibacterial effect. The MBC/MIC ratio is less than 4. These substances could therefore be used as antibacterial agents [29-31].

## 5. CONCLUSION

This work made it possible to demonstrate the effectiveness and *In vitro* antibacterial properties of derivatives resulting from S-alkylation and Mannich-based substances on six bacterial strains (four (4) strains of *E. coli* and two (2) strains of enterobacteria) isolated from foods. The MICs obtained ranged between 31.25 and 125  $\mu\text{g/mL}$ . As for the CMBs, they were between 31.25 and 250  $\mu\text{g/MI}$ . CMB/MIC ratios indicated chemicals with bactericidal activity. These results made it possible to conclude that some of these synthesized chemical compounds have antibacterial properties. Thus, these molecules could contribute to the fight against bacterial infections.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Fung F, Huei-Shyong W, Suresh M. Food safety in the 21st century. *Biomedical Journal*. 2018;41(2):88-95.

2. Kirk DM, Pires M. Sara, Robert E. Black, Caipo M, Crump AJ, Devleeschauwer B, Döpfer D, Fazil A, Fischer-Walker CL, Hald T, Hall JA, Keddy HK, Lake JR, Lanata CF, Torgerson PR, Havelaar AH, Angulo FJ. World health organization estimates of the global and regional disease burden of 22 foodborne bacterial, protozoal, and viral diseases, 2010: A Data Synthesis. *PLoS Med*. 2015;12(12): e1001921.
3. Ferreira A, Petretti C, Vanina B. Les toxi-infections alimentaires collectives, In: *Biologie de l'alimentation Humaine*, France. 2013;477-493.
4. World health organization (WHO). *Collective catering regional publication*, European series Geneva. 1996;71.
5. Fleckenstein JF, Kuhlmann M, Sheikh AP. Acute bacterial gastroenteritis. *Gastroenterol Clin North Am*. 2021;50 (2):283-304.
6. Ministry of health and public hygiene, Ivory Coast. *Food poisoning in Bondoukou*. In: *bondoukou, dispatches, health/AIP*; 2022.
7. Bernard H, Faber M, Wilking H, Haller S, Höhle M, Schielke A, Ducombe T, Sifczyk C, Merbecks SS, Fricke G, Hamouda O, Stark K, Werber D. Large multistate outbreak of norovirus gastroenteritis associated with frozen strawberries, Germany, 2012. *Eurosurveillance*. 2014;19(8):20719.
8. Antunes P, Novais C, Peixe L. Food-to-humans bacterial transmission. *Microbiol Spectrum*. 2020;8(1):1-26.
9. Vieira K. C. de O, Silva HRA. da, Rocha IPM, Barboza E, Eller LKW. Food borne pathogens in the omics era. *Critical Reviews in Food Science and Nutrition*. 2021;62(24):1-16.
10. Pons MJ, Mosquito S, Gomes C, Del VLJ, Ochoa J, Ruiz J. Analysis of quinolone-résistance in commensal and diarrheagenic *Escherichia coli* isolates from infants in Lima, Peru. *Trans R Soc Trop Med Hyg*. 2014;108(1):22-8.
11. Popiolek L. Hydrazide-hydrazones as potential antimicrobial agents: overview of the literature since 2010. *Med Chem Res*. 2017;26:287-301.
12. Bolou GA. Urban sprawl and access to socio-collective services in the city of Daloa (Central-West Ivory Coast).

- Scientific Journal Specializing in Geography. 2021;5:26-46.
13. Bssaibis F, Gmira N, Meziane M. Activité antibactérienne de *Dittrichia viscosa* (L.) W. Greuter. Revue de Microbiologie Industrielle, Santé et environnement. 2009;3:44-55.
  14. Ponce AG, Fritz R, Del Vallec, Rouras I. Antimicrobial activity of essential oils on the native microflora of organic Swisschard. Society of Food Science and Technology. 2003;36(7):679-684.
  15. Koné WM, Atindehou KK, Terreaux C, Hostettmann KURT, Traore D, Dosso M. Traditional medicine in North Côte-d'Ivoire: screening of 50 medicinal plants for antibacterial activity. Journal of Ethnopharmacology. 2004;93(1): 43-49.
  16. Nayaka NMDMW, Sasadara MMV, Sanjaya DA, Yuda, PESK, Dewi NLKAA, Cahyaningsih E, Hartati R. Piper betle (L): Recent review of antibacterial and antifungal properties, safety profiles, and commercial applications. Molécules. 2021; 26(2321): 1-21.
  17. Khomotso JM, Chelule PK. Safe Food handling knowledge and practices of street food vendors in polokwane central business district. Foods. 2020;9(11): 1-10.
  18. Attien YP, Zebre AC, Sina H, Angaman DM, Baba-Moussa L, Adjehi TD. Assessment of the sanitary quality of the dishes sold in street's restaurants with high frequeation in Daloa (Ivory Coast). Int. J. Curr. Microbiol. App. Sci. 2021;10(12):206-215.
  19. Rahman MM, Arif MT, Bakar K, Talib Z. Food safety knowledge, attitude and hygiene practices among the street food vendors in Northern Kuching City, Sarawak. Borneo Sci. 2016;31: 95-103.
  20. Mosupye M, Von Holy A. Microbiological hazard identification and exposure assessment of street food vending in Johannesburg, South Africa Francina. International Journal of Food Microbiology. 2000;61(2000):137-145
  21. Attien P, Toe E, Kouassi KA, Zébré AC, Gomé MN, Sina H. Dadie A. Evaluation of health risks related to the consumption of fish from the Guéssabo river. Food and Nutrition Sciences. 2022; 13(1):55-64.
  22. Zébré AC, Yao PA, Sina H, Gome M, Konate I, Baba-Moussa L. Sanitary investigation on microbial and heavy metals of grilled fish sold in the night street market of Daloa (Ivory Coast). Journal of Advances in Microbiology. 2022;22(12):1-9.
  23. Koffi-Nevry R, Assi CBJ, Assemmand EF, Wognin AS, Kousssemon M. Origin of evidence of fecal contamination of water used for watering lettuce (*Lactuca sativa*) grown in the peri-urban area of Abidjan. Journal of Applied Biosciences. 2012;52: 3669-3675.
  24. Cardinale E, Perrier Gros-Claude JD, Tall F, Gueye EF, Salvat G. Risk factors for contamination of ready-to-eat street-vended poultry dishes in Dakar, Senegal. It. J. Food Microbiol. 2005; 25:157-165.
  25. Iredell J, Brown J, Tagg K. Antibiotic resistance in enterobacteriaceae: Mechanisms and clinical implications. The Bmj. 2016;352:1-19.
  26. Li Y, Wang Z, Ajani JA, Song S. Drug resistance and Cancer stem cells. Cell Communication and Signaling. 2021;19 (1):1-11.
  27. Zniber R, Ahmed M, Redouane A, Mostafa EG, Abdelkrim FM, Abdel IM, Abdelkamel H. Biological activities of benzimidazole derivatives. Romanian Journal of Chemistry. 2009;54(8) :643-650.
  28. Gheorghe R. Mannich bases in medicinal chemistry and drug design. European Journal of Medicinal Chemistry. 2015; 89:743-816.
  29. Abdi RD, Dego KO. Antimicrobial activity of *Persicaria pensylvanica* extract against *Staphylococcus aureus*. European Journal of Integrative Medicine. 2019;29: 1-7.
  30. INS. General population and housing census: Overall results. Ministry of Planning and Development of Côte d'Ivoire, Abidjan (Côte d'Ivoire). 2021; 37.
  31. Konate K, Mavoungou JF, Lepengue AN, Aworet-Samseny RR, Hilou A, Souza A, Dicko MH, M'Batchi B. Antibacterial activity against beta- lactamase producing methicillin and ampicillin-resistants



staphylococcus aureus: fractional determination, Ann. Clin. Microbiol.  
Inhibitory Concentration Index (FICI) Antimicrob. 2012;11(18):1-12.

---

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/117522>