



Effect of Leaf Clipping and Nitrogen Split Application on Bridegroom Rice Productivity

J. Saranya ^a, S. Jawahar ^{b*}, K. Suseendran ^a,
C. Kalaiyarasan ^a and D. Elayaraja ^c

^a Department of Agronomy, Faculty of Agriculture, Annamalai University, Annamalainagar, Tamil Nadu, India.

^b Dr. M. S. S. AC & RI, Tamil Nadu Agricultural University, Thanjavur, Tamil Nadu, India.

^c Department of Soil Science Agricultural Chemistry, Faculty of Agriculture, Annamalai University, Annamalainagar - 608 002, Tamil Nadu, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i92891>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/123337>

Received: 13/07/2024

Accepted: 15/09/2024

Published: 18/09/2024

Original Research Article

ABSTRACT

Bridegroom rice is one the most important indigenous rice variety predominantly grown in Tamil Nadu, India. Lodging in nature and poor responsive to nitrogenous fertilizers affects the yield of this crop. To minimize crop lodging and to enhance the productivity, a field experiment was carried out during Samba 2023 at Experimental Farm, Department of Agronomy, Annamalai University, Tamil Nadu, India to study the effect of leaf clipping and nitrogen split application on bridegroom rice productivity. The experiment was laid out in Factorial randomized block design

*Corresponding author: E-mail: jawa.au@gmail.com;

Cite as: Saranya, J., S. Jawahar, K. Suseendran, C. Kalaiyarasan, and D. Elayaraja. 2024. "Effect of Leaf Clipping and Nitrogen Split Application on Bridegroom Rice Productivity". *Journal of Experimental Agriculture International* 46 (9):937-45. <https://doi.org/10.9734/jeai/2024/v46i92891>.

(FRBD) with three replications. The treatments consisted of two factors viz., Factor I -leaf clipping (L): L₁- No leaf clipping, L₂- leaf clipping on 30 DAT at one third of leaf, L₃- leaf clipping on 45 DAT at one third of leaf and Factor II - split application of N (N) : N₁- no splits (no NPK), N₂- 4 equal splits (basal, 25, 50, 75 DAT), N₃- 5 equal splits (basal 20, 40, 60,80 DAT) and N₄- 6 equal splits (basal, 15, 30, 45, 60 ,75 DAT). Among the treatments, leaf clipping on 45 DAT at one third of leaf along with application of nitrogen in 6 equal split registered its superiority compared to others which recorded the higher values for growth attributes viz., plant height (243cm), number of tillers hill⁻¹ (23.42), root length (33.56cm), root volume (45.19 cc) and dry matter production (12592 kg ha⁻¹), yield attributes viz., number of panicles m⁻² (407.62) and number of grains panicle⁻¹ (158.34) and test weight (42.34g) and yields of rice (grain - 3950 kg ha⁻¹ and straw - 6530 kg ha⁻¹). Thus, it can be concluded that leaf clipping on 45 DAT at one third of leaf along application of nitrogen in 6 equal splits is a viable practice to enhance the growth and yield of bridegroom rice.

Keywords: Leaf clipping; nitrogen split application; bridegroom rice; growth and yield.

1. INTRODUCTION

Rice (*Oryza sativa* L.) is one of the most important staple foods of nearly more than half of the global population and the majority of Asians consumes rice thrice a day. More than 90% of the world's rice is grown and consumed in Asia. The refrain, "rice is life" is the most fitting for India because this crop is essential to the country's food security and provides a living for millions of people. 14.29 Mt of India's total rice output is derived from the dry season (rabi) and the rest from the rainy season (kharif) [1]. According to United Nations estimates, by 2050, there will be 9.7 billion people on the planet. As a result, providing food and nutrition security for this expanding population is getting more difficult. Despite being a cereal with a high caloric and calorie content, rice's grains are deficient in protein, minerals, and other nutrients [2]. Now a day, all over the world, scientists focused their attention on nutritional security than food security to alleviate malnutrition. After green revolution, improved rice varieties replaced our traditional rice due to their high yield potential. But most of the improved rice varieties are lack of vitamins and minerals. At the same times, traditional varieties are a viable source of various agricultural properties as well as sources of many bioactive non-vital nutrients. Traditional rice varieties are rich in nutrients such as vitamin D, calcium, thiamine, riboflavin, glutamic acids, and high in fiber [3]. They are low in fat, sugar, gluten, and contain oryzanol, a molecule that reduces fat production in the body, making them a great diet for diabetics and hypertensive people. Additionally, a few nutritional components are also present in various fragments of traditional rice (viz. endosperm, germ function, and bran) in small quantities with different biological activities called bioactive compounds [4].

Farmers in Tamil Nadu have documented 174 traditional rice varieties as a result of this. Some of the rice varieties are *Karuppukavuni*, *Kullakkar*, *Illupai poo*, *Mapillai samba*, *Kichadi Samba*, *Thuyamalli*, *Kaivara samba*, *Karunguruvai*, *Kothamalli samba*, *Tiruchengode samba*, *Salem senna*, *Sivan samba*, *Thanga samba*, etc., Among them, Indigenous bridegroom rice is one of the most important rice variety. It has a duration of 165 – 170 days and taller than other traditional rice varieties. This variety can withstand flood and drought situations. The water used to wash this milled rice is beneficial for treating stomach and mouth ulcers. Mappillai samba also has a unique name "Bridegroom Rice", which has an interesting historical significance and interesting stories behind it. The story goes back to the early days when grooms were often tasked with lifting heavy rocks to demonstrate their strength. To aid them in this endeavour, the grooms were provided with Mappillai Samba rice, renowned for its ability to enhance physical prowess. It helps diabetes patients because of the low glycemic index value of 66. It is also very high in iron, calcium and fibre. The calcium content of Mappillai samba rice (boiled) is 50.4mg/100g (the value of calcium for parboiled rice is 8.11 mg/100g as per IFCT, 2017). Phytosterols and tocopherols present in this rice are beneficial in reducing several lifestyle disorders [5]. Rightfully So, Mappillai Samba rice is filled with rich iron, fibre, vitamins, and antioxidants, providing enormous benefits to our body. It aids in blood, muscle, neuron, and vein nourishment, improves male fertility, lowers cholesterol, and cures colon cancer [6]. However the productivity is very low due to lodging in nature, poor responsive inorganic fertilizers and inadequate application essential plant nutrients and lack of adaptation improved agronomic management practices. Among them, lodging and inadequate and improper application of

nitrogen greatly affects the yield. Hence, suitable agronomic management practices are needed to increase the yield of bridegroom rice. Some of them are leaf clipping and application of N in split doses.

Leaf clipping is one of the most affordable methods of raising yield by cutting one third of leaf. Generally, it alleviates drought stress in clipped plants, it has also been reported that leaf clipping presumably removes transpiring biomass and conserves soil moisture [7]. In this study the main advantages is to preventing lodging in the event of excessive vegetative growth and giving farmers fresh feed for their work animals without compromising grain output. Therefore, cutting leaves doesn't actually harm grain production. Leaf clipping is provides more green feed to the animals; To overcome the scarcity of green forage, it is necessary to explore the feasibility of using green leaves of rice. It might turn out to be one of the most economical ways to increase the land's total yield [8].

Fertilizers are mainly recommended for crop plants to supply essential nutrient element throughout the growth and development period. Among the all essential nutrient elements, nitrogen is needed most for crop plants. Applying nitrogen on time and in parts allows rice to use it more efficiently throughout the growth season. This is because providing nitrogen to the crop reduces nitrogen losses and provides targeted nutrients at growth peaks. Rice is a poor nitrogen user, nitrogen usage efficiency ranged from 30 to 50 %. One method for effectively using nitrogen throughout the growth season is the split application of nitrogenous fertilizers, which improves nitrogen absorption, decreases denitrification losses, and synchronizes with plant needs [9]. Researches on combined effect of leaf clipping and split application nitrogen on rice is very limited and particularly nil in bridegroom rice. Therefore, an experiment was programmed to study the effect of leaf clipping and application nitrogen in split doses on growth and yield of rice.

2. MATERIALS AND METHODS

A field experiment was carried out during Samba 2023 at Experimental Farm, Department of Agronomy, Annamalai University, Tamil Nadu, India (11.38°N latitude and 79.72°E longitudes with an altitude of +5.79 m MSL) to study the effect of leaf clipping and split application of

nitrogen on productivity of bridegroom rice. The experiment was laid out in Factorial randomized block design (FRBD) with three replications. The treatments consisted of two factors viz., Factor I - leaf clipping (L): L₁- No leaf clipping, L₂- leaf clipping on 30 DAT at one third of leaf, L₃- leaf clipping on 45 DAT at one third of leaf and Factor II - split application of N (N) : N₁- No split (no NPK), N₂ – recommended 4 equal splits (basal,25,50,75 DAT, N₃- 5 equal splits (basal 20, 40, 60, 80 DAT) and N₄ - 6 equal splits (basal, 15, 30, 45, 60 ,75 DAT). The bridegroom rice seeds were collected from indigenous rice growing farmer's at Kollidam block, Mayiladurai District, Tamilnadu field. Recommended 60 kg seeds were treated with Carbendazim@ 2g per kg of seed. Pre-germinated seeds were sown in the nursery. Thirty days old seedlings were transplanted in lines with two seedlings hill⁻¹ by adopting a spacing of 20 × 15 cm to accommodate a plant population of 33.33 plants m⁻². The bridegroom rice was fertilized with 150:50:50 NPK kg ha⁻¹. Entire P₂O₅ was administered as a baseline and K was administered in four equal splits at basal, tillering, panicle initiation and heading stages. Required quantity of N was applied as per the treatments. Need based plant protection measures were carried out for controlling insect pest and disease in rice crop. Growth attributes (plant height, number of productive tillers hill⁻¹, root length, root volume and dry matter production), yield attributes (number of panicles m⁻² and number of grains panicle⁻¹ and 1000 grain weight) and yields of bridegroom rice was recorded at critical stages. Grains were separated by hand threshing, cleaned and sun dried to bring the moisture content to 14 % and the weight of grain and straw were recorded and expressed as kg ha⁻¹. The statistical analysis of the data was followed as per the recommendation of Gomez and Gomez [10]. The critical difference (CD) were arrived at 5% probability level (P = 0.05). Treatment differences that not significant were denoted by 'NS'.

3. RESULTS AND DISCUSSION

3.1 Growth Attributes

Leaf clipping and split application N significantly influenced on the growth attributes viz., plant height, number of productive tillers hill⁻¹, root length, root volume and dry matter production of bridegroom rice (Table 1).

Table 1. Effect of leaf clipping and nitrogen split application on growth attributes of bridegroom rice at harvesting stage

F – I	Plant height (cm)at harvest				Number of tillers hill⁻¹at maximum tillering stage				Root length (cm) at flowering stage				Root volume (cc) at flowering stage				Dry matter production (kg ha⁻¹) at harvest			
F -II	L₁	L₂	L₃	MEAN	L₁	L₂	L₃	MEAN	L₁	L₂	L₃	MEAN	L₁	L₂	L₃	MEAN	L₁	L₂	L₃	MEAN
N₁	186.9	215.6	223.5	208.7	10.6	15.0	16.4	14.69	11.7	20.45	29.78	20.67	20.23	28.34	30.98	26.51	5413	9912	10651	8658
N₂	226.3	230.3	232.6	229.7	16.6	18.7	19.5	18.32	19.2	28.29	31.21	26.25	29.21	34.51	41.41	35.04	8236	10641	11846	10241
N₃	237.5	238.6	240.8	239.0	19.0	20.7	22.8	20.90	25.3	30.42	32.89	29.56	33.45	39.72	43.21	38.79	10464	11872	12384	11573
N₄	239.4	241.2	243	241.2	20.5	22.9	23.4	22.32	28.6	31.91	33.56	31.38	34.78	43.81	45.19	41.26	11126	12414	12592	12044
MEAN	225.0	231.4	235.0		16.2	19.3	20.5		21.2	27.76	31.86		29.41	36.59	40.19		8809	11210	11868	
	L	N	L-N		L	N	L-N		L	N	L-N		L	N	L-N		L	N	L-N	
SEd	2.13	0.45	2.61		1.34	0.41	1.32		3.1	1.41	3.12		2.32	0.41	3.12		346	123	410	
CD (P=0.05)	4.32	1.89	5.42		2.56	0.93	2.94		5.2	2.83	6.34		4.56	1.82	5.25		660	249	823	

Among the different days of leaf clipping, one-third of leaf clipping on 45 DAT (L_3) recorded higher value of the plant height (235cm), maximum number of tillers hill⁻¹ (20.56), longer root length (31.86cm), larger root volume (40.19 cc) and elevated dry matter production (11868 kg ha⁻¹) of bridegroom rice. Non lodging of crop, erectness of leaves, better utilization of sun light, higher photosynthetic activity improved the cell division, cell elongation and enlargement caused tallest plant at one third leaf clipping on 45 DAT [11]. One third of leaf clipping on 45 DAT increased the root length and root volume of rice, therefore, the plant extracted higher amount of applied nutrients, which favourably enhanced growth attributes of rice [12]. Cutting of leaves had no significant effect on tiller number plant⁻¹ in bridegroom rice [13]. Leaf clipping did not affect dry matter production at harvest stage, which significantly increased the dry matter accumulation of rice due to production of more number of tillers hill⁻¹, root length, root volume and number leaves [14]. Weaker culm caused crop lodging which affects the photosynthetic activity and accumulation photosynthates resulted in lesser growth attributes at L_1 (no leaf clipping).

Among the different split dose of N, application of N in 6 equal split doses (N_4) recorded the higher value of plant height (241.23cm), more number of tillers hill⁻¹ (22.32), increased in root length (31.38), root volume (41.26cc) and dry matter production (12044 kg ha⁻¹) of bridegroom rice. This was on par with application of N in 5 equal splits (N_3) which was superior over recommend application of N in 4 equal splits (N_2) which recorded next higher plant height (239.01cm), total number of tillers hill⁻¹ (20.90), root length (29.56cm), root volume (38.793 cc) and dry matter production (11573kg ha⁻¹) in bridegroom rice. The lowest values on growth attributes were observed without nitrogen application (N_1). The plant height was significantly increased with the increase in the availability of nitrogen levels at different growth stages. Higher availability of nitrogen due to split doses enhanced the cell division; cell elongation and enlargement resulted in higher growth parameters in bridegroom rice [15]. Increase in plant height in response to split application of N fertilizers is probably due to higher photo assimilates and there by more dry matter accumulation. Elevated leaf area caused higher photo assimilate production resulted in improved dry matter accumulation [16]. Increased in N splits increased the dry matter accumulation of

bridegroom rice due to increasing vegetative growth resulting from higher photosynthetic activities caused by nitrogen [17]. Split application of N also produced more DMP that could produce additional photosynthates for the development of root system and increased root growth [18] who reported that split application of N significantly increased the dry matter of plant height, root growth and number of leaves as well as total dry matter at tillering, flowering and harvesting stage.

Among the various treatment combinations, one third of leaf clipping on 45 DAT and application of nitrogen in 6 equal splits (L_3N_4) significantly registered higher plant height (243cm), higher number of tillers hill⁻¹ (23.42), more root length (33.56cm), larger root volume (45.19cc) and increased dry matter production (12592 kg ha⁻¹) in bridegroom rice. This was followed by 45 DAT of leaf clipping and application of N in 5 equal splits (L_3N_3) recorded relatively higher plant height (240.89 cm), total number of tillers hill⁻¹ (22.89), root length (32.89 cm), root volume (43.21cc) and dry matter production (12384 kg ha⁻¹) over other treatment combination. Leaf clipping at the early vegetative stage significantly increased plant growth, reduced the lodging due to increased culm diameter and culm wall thickness which improved root growth and involvement of leaves in better photosynthesis resulted in improved growth attributes in lateral stages, similarly, N application improved photosynthetic activity, enhanced the crop growth and increases the number and size of meristematic cells, which leads to formation of new shoot due to the role of split application of nitrogen in improving rice growth, internode elongation, and photosynthesis, vigorous vegetative growth by increased cell division and cell elongation [19]. The least values on growth attributes were recorded with no leaf clipping and no N application (L_1N_1), which recorded the fewer values on plant height (186.98cm), total number of productive tillers hill⁻¹ (10.61), root length (11.78 cm), root volume (20.23cc) and dry matter production (5413 kg ha⁻¹) in bridegroom rice.

3.2 Yield Attributes and Yield

Leaf clipping and application nitrogen in split doses significantly enhanced the yield attributes and yield of bridegroom rice (Table 2). Among the different days of leaf clipping, one third leaf clipping on 45 DAT (L_3) recorded significantly highest values for number of panicle m⁻² (334.38), number of grains panicle⁻¹ (150.11),

Table 2. Effect of leaf clipping and nitrogen split application on yield attributes and yields (grain and straw)of bridegroom rice

F -I	Number of panicle m ⁻²				Number of filled grains panicle ⁻¹				1000 grain weight (g)				Grain yield (Kg ha ⁻¹)				Straw yield (Kg ha ⁻¹)			
	L ₁	L ₂	L ₃	MEAN	L ₁	L ₂	L ₃	MEAN	L ₁	L ₂	L ₃	MEAN	L ₁	L ₂	L ₃	MEAN	L ₁	L ₂	L ₃	MEAN
N₁	169.2	210.3	239.6	206.3	97.6	126.7	139.5	121.2	29.6	32.7	38.9	33.7	1801	2548	2825	2391	3220	5180	5420	4606
N₂	251.7	287.9	310.6	283.4	125.3	137.1	148.4	136.9	35.4	39.4	40.3	38.3	3021	3445	3675	3380	5160	5678	5940	5592
N₃	279.9	317.6	379.6	325.7	136.1	149.6	154.1	146.6	38.2	40.9	41.32	40.1	3195	3610	3865	3556	5505	6085	6350	5980
N₄	319.9	340.2	407.6	355.9	139.3	156.5	158.3	151.4	39.3	41.6	42.34	41.1	3290	3700	3950	3646	5695	6215	6530	6146
MEAN	255.2	289.0	334.3		124.6	142.5	150.1		35.6	38.7	40.72		2826	3325	3578		5250	5789	6060	
	L	N	L·N		L	N	L·N		L	N	L·N		L	N	L·N		L	N	L·N	
SEd	26.1	13.0	34.3		4.31	1.93	5.34		1.44	0.43	1.23		39.4	45.5	78.92		68.14	78.68	136.2	
CD (P=0.05)	52.5	26.1	68.6		8.7	3.91	10.6		2.95	0.85	3.48		81.8	94.5	163.6		141.3	163.2	282.6	

1000 grain weight (40.72g) in bridegroom rice. The same treatment registered significantly higher grain yield (3578 kg ha⁻¹) and straw yield (6060 kg ha⁻¹) of bridegroom rice. The per cent increase in grain yield over one-third of leaf clipping on 30 DAT (L₂) and no leaf clipping L₁ was 7.3 % and 21.7% respectively. It could be due higher photosynthetic activity, production of higher number of spikelets per panicle, increased in accumulation photosynthates in the economic part which caused higher grain and straw yield [20]. One third leaf clipping on 30 DAT (L₂) was next in order and recorded relatively higher values for number of panicle m⁻² (289.05), number of filled grains panicle⁻¹ (142.51), 1000 grain weight (38.7 g), grain (3325 kg ha⁻¹) and straw yield (5789 kg ha⁻¹) over no leaf clipping (L₁). The lesser yield attributes viz., number of panicle m⁻² (255.2), number of filled grains panicle⁻¹ (124.6) and 1000 grain weight (35.6 g) and yields viz., grain yield (2826 kg ha⁻¹) and straw yield (5250 kg ha⁻¹) in bridegroom rice was recorded under L₁ (no leaf clipping).

Among the different split dose of N, application of N in 6 equal splits (N₄) recorded significantly more number of panicle m⁻² (355.96), higher number of filled grains panicle⁻¹ (151.41), increased in 1000 grain weight (41.11) and elevated grain (3646 kg ha⁻¹) and straw (6146 kg ha⁻¹) yield of bridegroom rice which was comparable with application of N in 5 equal split and recorded number of panicle m⁻² of 325.7, number of filled grains panicle⁻¹ of 146.6 and 1000 grain weight of 40.1g. The grain and straw yield at N₃ was 3556 kg ha⁻¹ and 5980 kg ha⁻¹, respectively. Recommend N in 4 equal splits was next in order. Yield attributes were significantly increased with increase in availability of nitrogen level at different growth stages. Increase in grain yield at application of N in 6 equal splits could be due to application nitrogen in split doses enhanced availability and utilization of N by rice crop which improved the number of panicle m⁻², number of grains panicle⁻¹ and 1000 grains weight. Higher number of panicle m⁻² is influenced by the number of tillers that develop during the vegetative stage; higher number of grains panicle⁻¹ and number filled spikelet due to N fertilization might be due to optimum availability of N, resulting in more synthesis of photosynthates and a possible source to sink relationship [21]. Increased N splits significantly enhanced the grain yield due to higher yield attributes coupled with efficient translocation of food material to sink. Application of nitrogenous fertilizer at proper time with a suitable rate

according to the soil conditions can significantly increased grain yield. Split application of nitrogen at critical growth stages like tillering can increase the number of panicles and number of spikelets per spike in rice [22]. The lowest yield attributes and yield of bridegroom rice was observed without N addition (N₁).

Among the treatment combinations, one third of leaf clipping on 45 DAT and application of N in 6 equal splits (L₃N₄) significantly recorded more number of panicle m⁻² (407.6), higher number of filled grains panicle⁻¹ (158.34), increased 1000 grains weight (42.34g) and highest grain (3950 kg ha⁻¹) and straw (6530 kg ha⁻¹) yield in bridegroom rice. This was followed by one third of leaf clipping on 45 DAT and application of N in 5 equal splits (L₃N₃) and recorded relatively more number of panicle m⁻² (379.6), higher number of filled grains panicle⁻¹ (154.1), increased 1000 grains test weight (41.32g) and highest grain (3865 kg ha⁻¹) and straw (6530 kg ha⁻¹) yield in bridegroom rice. The highest yield attributes and yield recorded at one third of leaf clipping on 45 DAT and application of N in 6 equal splits (L₃N₄) could be due to leaf clipping at early vegetative stage significantly increased growth attributed which favoured the yield attributed resulted in higher grain and straw yield. Increased synthesis of carbohydrates might have increased the grain yield due to assimilation of photosynthates, further split application of N enhanced N availability in the soil at all critical growth stages and increased the nutrient uptake by the crop resulted in higher growth and yield attributes resulted in higher grain and straw yield [21].

4. CONCLUSION

The experimental results showed that there was a marked variation in the productivity of rice due to one-third of leaf clipping and split application of nitrogen. In the light of above said facts, it can be concluded that the combination of one third of leaf clipping on 45 DAT along with application of nitrogen in 6 equal splits are significantly enhanced the overall productivity of bridegroom rice. Therefore, this treatment can be recommended to the farming community, especially to traditional rice growers.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Mondal P, Sadhukhan AK, Ganguly A, Gupta P. Optimization of process parameters for bio-enzymatic and enzymatic saccharification of waste broken rice for ethanol production using response surface methodology and artificial neural network genetic algorithm. *Biotech.* 2021; 11: 1-18.
2. Das T, Das AK, A Review of the Prospects of Agro biodiversity Management in the Traditional Rice Fields and Home gardens of the Region. *Springer Socio-economic and Eco-biological Dimensions in Resource use and Conservation: Strategies for Sustainability.* 2020;117-134.
3. Verma DK, Srivastav PP, Mohan M, Nutritional quality evaluation of different rice cultivars. *Agronomic rice practices and postharvest processing.* 2018;331-394.
4. Rondanelli M, Braschi V, Gasparri C, Nichetti M, Faliva MA Peroni G, Perna S. Effectiveness of non-animal chondroitin sulfate supplementation in the treatment of moderate knee osteoarthritis *Nutrients.* 2019;11(9):2027.
5. Rubilah MJ, Nancy M, Parimala K. Traditional Rice Varieties; A storehouse of nutrition. *Center for Indian knowledge System.* 2023;1- 20.
6. Bhat FM, Riar CS, Health benefits of traditional rice varieties of temperate regions. *Medicinal Aromatic Plants.* 2015; 4(198):2167-0412.
7. Georgias NJ, Ruess RW, Mcnaughton SJ, Western D. Ecological conditions that determine when grazing stimulates grass production. *Oecologia.* 1989; 81:316–322.
8. Khalid Usman. Effect of detopping on forage and grain yield of rice under agro-climatic conditions of DI Khan. *Sarhad J Agric.* 2007;23(1):1-4.
9. Subramanian E, Aathithyan C, Raghavendran VB, Vijayakumar S. Optimization of nitrogen fertilization for aerobic rice (*Oryza sativa*). *Indian J Agro.* 2020;65(2):180-184.
10. Gomez KA, Gomez AA. Duncan's Multiple Range Test. *Statistical procedures for Agric Res.* 1984;207- 215.
11. Karmaker Banani, Biswajit Karmakar. Effect of nitrogen rates and leaf clipping on forage and grain yield, and seed quality of transplant Aman rice. *Bangladesh Rice J.* 2019;23(2):49-57.
12. Medhi AK, Dey PC, Chetia SK, Ghose TJ. Effect of foliage pruning on vegetative growth and grain yield of rice under rainfed low land situation. 2015;33(11): 1335-1338.
13. Boonreund C, Marsom K. Effect of rice leaf cutting length on rice yield. *International Conference on Agricultural and Food Industry: Rajamangala Uni Tech.* 2015; 474-481.
14. Khalifa AB, Misra AN, Salem A K, Effect of leaf cutting on physiological traits and yield of two rice cultivars. *African Journal of Plant Sci.* 2008; 2(12):147-150.
15. Paul SK, Rahman MM. Effects of spacing and nitrogen levels on yield and yield contributing characters of maize. *Journal of the Bangladesh Agricultural University.* 2016;14(1):43-48.
16. Kumar U, Nayak AK, Shahid M, Gupta VV, Panneerselvam P, Mohanty S, Panda BB. Continuous application of inorganic and organic fertilizers over 47 years in paddy soil alters the bacterial community structure and its influence on rice production. *Agric Eco and Env.* 2018; 262:65-75.
17. Verma P, Kumar Chatterjee S, Ghosh S, Kumar Jangde M. Study of Soil Nitrogen level and Splitting on Nutrient Uptake around the Industries of Chhattisgarh, India. *Adv Materials Letters.* 2021;12(5):1-5.
18. Haque MA, Haque MM. Growth, yield and nitrogen use efficiency of new rice variety under variable nitrogen rates. *American J Plant Sci.* 2016;7(3):612-622.
19. Metwally TF, Gewaily EE, Naeem SS. Nitrogen response curve and nitrogen use efficiency of Egyptian hybrid rice. *Journal of Agricu Res Kafer El-Sheikh Uni.* 2011;37(1):73-84.
20. Ghosh A, Sharma AR. Effect of foliage pruning on performance of rice under semi-deep water situation. *Int Rice Res..* 1998;22(2):36-37.
21. Islam MD, Hasanuzzaman M, Rokonuzzman M. Effect of split application of nitrogen fertilizer on morphophysiological parameters of rice genotypes. *Int J of Plant Production.* 2009; 3(1):1735-6814.

22. Sathiya K, Ramesh T. Effect of split application of nitrogen on growth and yield of aerobic rice. Asian J Exp Sci.. 2009; 23:303- 306.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/123337>