



Effect of Foliar Application of Nano fertilizers on Soil Properties of Rice (*Oryza sativa* L.) Under Western UP, India

Sandeep Kumar Verma ^{a++*}, Mukesh Kumar ^{a#},
Vivek ^a, P. K. Singh ^{a#}, S. P. Singh ^{b#}, Bhim Singh ^{c#},
Aparna Verma ^{a#} and Shalini Roy ^{a#}

^a Department of Agronomy, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut Uttar Pradesh, 250110, India.

^b Department of Soil Science, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut Uttar Pradesh, 250110, India.

^c Department of Statistics, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut Uttar Pradesh, 250110, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i95048>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124550>

Short Research Article

Received: 24/07/2024
Accepted: 26/09/2024
Published: 30/09/2024

⁺⁺ Research Scholar;

[#] Professor;

^{*}Corresponding author: E-mail: sandeepverma35529@gmail.com;

Cite as: Verma, Sandeep Kumar, Mukesh Kumar, Vivek, P. K. Singh, S. P. Singh, Bhim Singh, Aparna Verma, and Shalini Roy. 2024. "Effect of Foliar Application of Nano Fertilizers on Soil Properties of Rice (*Oryza Sativa* L.) Under Western UP, India". *International Journal of Plant & Soil Science* 36 (9):973-79. <https://doi.org/10.9734/ijpss/2024/v36i95048>.

ABSTRACT

An experiment was conducted during *kharif* 2022 and 2023 at the Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut, Uttar Pradesh, to study the Effect of Foliar Application of Nano on Soil properties of Rice (*Oryza sativa* L.) Under Western UP. The investigation was carried out with fourteen treatments *viz.* Control, RDF (N:P:K-120:60:40 kg ha⁻¹ + 25 kg zinc sulphate), 100% RDF + Nano urea, 100% RDF + Nano potash, 100% RDF + Nano Zn, 100% RDF + Nano urea + Nano potash + Nano Zn, 75% RDF + Nano urea, 75% RDF + Nano potash, 75% RDF + Nano Zn, 75% RDF + Nano urea + Nano potash + Nano Zn, 50% RDF + Nano urea, 50% RDF + Nano potash, 50% RDF + Nano Zn and 50% RDF + Nano urea + Nano potash + Nano Zn. The nano fertilizers were applied at tillering stage (35 DAT). Result revealed that highest soil properties *viz.*, organic matter % (0.48 & 0.52), available nitrogen (225.6 & 228.7 kg ha⁻¹), available phosphorus (19.5 & 20.7 kg ha⁻¹), available potassium (235.6 & 233.8 kg ha⁻¹) and available zinc (0.80 & 0.83 g ha⁻¹) were recorded with application of 100% RDF + Nano urea + Nano potash + Nano Zn spray at tillering stage (35 DAT) respectively, during both the years. The lowest values of these parameters were recorded under control treatment.

Keywords: Nano fertilizers; rice; organic carbon; nutrient uptake; Soil pH.

1. INTRODUCTION

Rice (*Oryza sativa* L.) belongs to poaceae family with chromosome number (2n = 24). It is one of the major staple foods for over half the population worldwide. Rice is a crop of wet tropical climate but also grown in humid regions of subtropics. At globe the area, production and productivity of rice during 2022-23 was 165.70 million hectares (mha), 512.98 million metric tons (mmt) and 46.20 q ha⁻¹ respectively. India ranked first, in terms of area and second position in production under rice cultivation in the world with 47.83 mha area, 135.76 mmt production. This however reflects poor crop productivity being 42.60 q ha⁻¹ as against world's average of 46.20 q ha⁻¹, 82.80 q ha⁻¹ in United States and 70.8 q ha⁻¹ in China [1]. There are improved technologies and various interventions which could be adapted to increase the productivity in the country. In India, Uttar Pradesh ranks first in terms of area (5.70 mha) and second in terms of production (15.27 mt), followed by West Bengal. The productivity (2679 kg ha⁻¹) of rice was found to be sixth position in UP after Punjab (4340 kg ha⁻¹), Tamilnadu (3658 kg ha⁻¹), Andhra Pradesh (3470 kg ha⁻¹), Telangana (3366 kg ha⁻¹) and West Bengal (2996 kg ha⁻¹). E&S Division, DA&FW, [2].

In rice, fertilizers along with soil fertility, irrigation and quality seed are mainly responsible for increase the production as well as productivity. The nutrient requirement of the rice, In general is high and inadequate supply of nutrient often leads to low productivity [3]. The major nutrient element, which is insufficient in most of the

Indian soil, plays appreciably an important role in rice. Continuous and sole applications of inorganic fertilizer induce the soil sickness and disturb the soil environment resulted low productivity and unsustainability. Shortage of arable land, limited water and nutrient resources, necessitates increase in resource use efficiency without sacrificing production through effective use of modern technologies. In the context nanotechnology, nano fertilizers holds promise and can go a long way in sustaining soil health and crop production.

As per the present context, Nanotechnology, a new and rising aspect of science and technology provides a revolutionary scope of research as far as the field of Agriculture and Biotechnology is concerned. Nano fertilizers have the potential to lower soil toxicity, increase the efficiency with which nutrients are used, and mitigate the potential drawbacks of excessive dosing, frequent application, and production costs. Due to its smaller particle size and greater surface area, nano fertilizer is easily absorbed by plants, boosting their dry matter content for photosynthesis and ultimately crop yield [4]. For nourishing the rising population, keeping in view the environmental health, soil health and economic condition of farmers, nano fertilizers may be the best substitute to traditional chemical fertilizers.

2. MATERIALS AND METHODS

An experiment was conducted during *kharif* 2022 and 2023 at the Sardar Vallabhbhai Patel University of Agriculture and Technology,

Meerut, Uttar Pradesh, to study the Effect of Foliar Application of Nano on Soil properties of Rice (*Oryza sativa* L.) Under Western UP. The investigation was carried out on well drained sandy clay loam soil, low in organic carbon (0.45 & 0.46 %) and available nitrogen (217.2 & 220.3 kg ha⁻¹), medium in available phosphorus (17.5 & 18.1 kg ha⁻¹), potassium (231.2 & 234.2 kg ha⁻¹), zinc (0.70 & 0.73 ppm) and slightly alkaline pH (7.7 & 7.6) in randomized block design with 14 treatments during both the years. The area is ploughed twice—once with a moldboard plough and once with a cultivator before transplanting. After that, the plots were marked according to the layout plan and dressed properly with spade. The rice nursery 'Pusa Basmati 1637' was transplanted manually at a distance 20x10 cm. The entire amount of phosphorus, potassium, and zinc, along with 50 % of the nitrogen, were given at the time of transplanting, and the remaining 50% was top-dressed in two equal portions at 25 and 55 DAT. Nano urea and potash (4 ml/litre of water) and nano zinc (2 ml/liter of water), were sprayed at tillering stage by mixing in 500 liters of water/ha. Experimental data were statistically analyzed by using the analysis of variance (ANOVA) as described by Gomez and Gomez [5].

3. RESULTS AND DISCUSSION

3.1 Organic Carbon (%)

The data pertaining to organic carbon have been presented in Table 1.

The highest organic carbon (0.48 & 0.52) was recorded with the application of 100% RDF + Nano urea + Nano potash + Nano Zn which remained statistically at par with 75% RDF + Nano urea + Nano potash + Nano Zn spray at tillering stage (35 DAT) but significantly higher than rest of the treatments during both the years. The lowest organic carbon (0.30 & 0.31) was recorded under control treatment during both the years. The increase in organic carbon due to addition of nano fertilizers along with chemical fertilizers considerably boosted nutrient uptake by rice and improved the organic carbon content, N, P, and K status of soil [6,7].

3.2 Soil pH

The data pertaining to soil reaction after harvest of the paddy crop have been presented in Table 1.

The soil reaction did not significantly influence by foliar application of nano fertilizers, however the maximum (7.9 & 8.0) and minimum (7.3 & 7.4) was found under control and 75 % RDF + Nano urea + Nano potash + Nano Zn, respectively spray at tillering stage (35 DAT) during both the years.

3.3 Electrical Conductivity (dSm⁻¹)

The data pertaining to electrical conductivity after harvest of the paddy crop have been presented in Table 1.

The electrical conductivity did not significantly influence by foliar application of nano fertilizers, however the maximum (0.25 & 0.26) and minimum (0.19 & 0.20) was found with the application 100 % RDF + Nano urea + Nano potash + Nano Zn spray at tillering stage (35 DAT) and under control respectively during both the years.

3.4 Bulk and Particle Density

The data pertaining to bulk and particle density after harvest of the paddy crop have been presented in Table 1.

The bulk and particle density did not significantly influence by foliar application of nano fertilizers, however the maximum bulk density (1.41 & 1.39) and minimum (1.34 & 1.33) was found under control and 100 % RDF + Nano urea + Nano potash + Nano Zn respectively spray at tillering stage (35 DAT) during both the years. Particle density also record same trend with maximum (2.66 & 2.65) and minimum (2.61 & 2.59) was found under control and 100 % RDF + Nano urea + Nano potash + Nano Zn respectively spray at tillering stage (35 DAT) during both the years.

3.5 Available Nitrogen (kg ha⁻¹)

The data pertaining to available nitrogen after harvest of the paddy crop have been presented in Table 2.

The maximum amount of available nitrogen (225.6 & 228.7 kg ha⁻¹) was recorded with the application of 100% RDF + Nano urea + Nano potash + Nano Zn which remained statistically at par with 100% RDF + Nano urea spray at tillering stage (35 DAT) but significantly higher than rest of the treatment respectively, during both the years whereas lowest amount of available nitrogen (203.3 & 205.3 kg ha⁻¹) was recorded

Table 1. Effect of foliar application of nano fertilizers on physical and chemical properties of soil after harvest of paddy crop

S. No.	Treatments	Organic carbon (%)		Soil pH		EC (dSm ⁻¹)		Bulk density (g cc ⁻¹)		Particle density (g cc ⁻¹)	
		2022	2023	2022	2023	2022	2023	2022	2023	2022	2023
T ₁	Control	0.30	0.31	7.9	8.0	0.19	0.20	1.41	1.39	2.66	2.65
T ₂	RDF (N:P:K-120:60:40 kg ha ⁻¹ + 25 kg zinc sulphate)	0.40	0.43	7.5	7.8	0.21	0.21	1.36	1.35	2.64	2.62
T ₃	100% RDF + Nano urea	0.43	0.47	7.4	7.6	0.24	0.23	1.35	1.35	2.63	2.60
T ₄	100% RDF + Nano potash	0.42	0.45	7.5	7.6	0.24	0.24	1.35	1.34	2.64	2.63
T ₅	100% RDF + Nano Zn	0.43	0.46	7.4	7.6	0.23	0.25	1.35	1.34	2.62	2.60
T ₆	100% RDF + Nano urea + Nano potash + Nano Zn	0.48	0.52	7.3	7.5	0.25	0.26	1.34	1.33	2.61	2.59
T ₇	75% RDF + Nano urea	0.40	0.42	7.5	7.7	0.21	0.22	1.36	1.34	2.64	2.61
T ₈	75% RDF + Nano potash	0.39	0.39	7.7	7.6	0.22	0.22	1.36	1.35	2.64	2.63
T ₉	75% RDF + Nano Zn	0.40	0.41	7.5	7.8	0.20	0.21	1.36	1.35	2.64	2.63
T ₁₀	75% RDF + Nano urea + Nano potash + Nano Zn	0.46	0.49	7.3	7.4	0.23	0.24	1.35	1.33	2.62	2.60
T ₁₁	50% RDF + Nano urea	0.38	0.40	7.8	7.8	0.22	0.23	1.38	1.38	2.65	2.65
T ₁₂	50% RDF + Nano potash	0.36	0.37	7.9	7.9	0.22	0.21	1.38	1.37	2.65	2.64
T ₁₃	50% RDF + Nano Zn	0.38	0.37	7.8	7.8	0.21	0.21	1.38	1.36	2.65	2.64
T ₁₄	50% RDF + Nano urea + Nano potash + Nano Zn	0.39	0.43	7.8	8.0	0.23	0.22	1.37	1.36	2.65	2.65
SEm±		0.01	0.01	0.2	0.3	0.07	0.02	0.05	0.04	0.1	0.2
CD (P= 0.05)		0.03	0.04	NS	NS	NS	NS	NS	NS	NS	NS

Table 2. Effect of foliar application of nano fertilizers on available N, P, K and Zn of soil after harvest of paddy crop

Symbol	Treatments	Available Nutrients (kg ha ⁻¹)						Zinc (g ha ⁻¹)	
		Nitrogen		Phosphorus		Potassium		2022	2023
		2022	2023	2022	2023	2022	2023		
T ₁	Control	203.3	205.3	13.2	14.4	213.1	214.4	0.61	0.63
T ₂	RDF (N:P:K-120:60:40 kg ha ⁻¹ + 25 kg ha ⁻¹ zinc sulphate)	220.2	224.5	18.1	18.7	233.0	234.5	0.73	0.74
T ₃	100% RDF + Nano urea	224.8	227.3	19.2	19.9	234.2	236.2	0.75	0.78
T ₄	100% RDF + Nano potash	221.5	225.3	18.7	19.3	234.8	237.1	0.74	0.77
T ₅	100% RDF + Nano Zn	222.7	225.8	17.8	18.4	233.5	235.6	0.78	0.82
T ₆	100% RDF + Nano urea + Nano potash + Nano Zn	225.6	228.7	19.5	20.7	235.6	237.9	0.80	0.83
T ₇	75% RDF + Nano urea	218.4	222.2	16.3	17.1	230.8	231.8	0.71	0.72
T ₈	75% RDF + Nano potash	216.3	219.5	15.7	16.7	231.5	233.2	0.70	0.70
T ₉	75% RDF + Nano Zn	217.8	220.7	15.3	16.4	230.3	231.1	0.73	0.75
T ₁₀	75% RDF + Nano urea + Nano potash + Nano Zn	219.5	222.8	15.8	17.8	232.2	233.8	0.77	0.81
T ₁₁	50% RDF + Nano urea	214.6	216.3	14.8	15.7	227.3	229.1	0.66	0.68
T ₁₂	50% RDF + Nano potash	213.3	214.2	14.5	15.5	228.2	229.5	0.65	0.66
T ₁₃	50% RDF + Nano Zn	213.8	215.8	14.1	15.2	226.6	228.2	0.68	0.69
T ₁₄	50% RDF + Nano urea + Nano potash + Nano Zn	215.7	217.4	15.0	16.1	228.8	229.9	0.69	0.71
	SEm±	0.8	0.9	0.4	0.6	0.4	0.5	0.01	0.01
	CD (P= 0.05)	2.4	2.7	1.2	1.7	1.3	1.5	0.02	0.04

under control treatment respectively, during both the years. The level of accessible nitrogen after crop harvest was sustained by the addition of nitrogen using chemical fertilizers such as urea and DAP. This increase may be related to the nitrogen in the soil becoming mineralized from nano fertilizers. When compared to chemical fertilizer alone, Jhanzab et al. [7] found that in addition of nano fertilizers along with chemical fertilizers considerably boosted nutrient uptake by rice and improved the organic carbon content, N, P, and K status. Similar findings also reported by Liu et al. [8].

3.6 Available Phosphorus (kg ha⁻¹)

The data pertaining to available phosphorus after harvest of the paddy crop have been presented in Table 2.

The maximum amount of available phosphorus (19.5 & 20.7 kg ha⁻¹) was recorded with the application of 100% RDF + Nano urea + Nano potash + Nano Zn which was at par with 100% RDF + Nano urea and 100% RDF + Nano potash spray at tillering stage (35 DAT) but significantly higher than rest of the treatment respectively, during both the years whereas lowest amount of available phosphorus (13.2 & 14.4 kg ha⁻¹) was recorded under control treatment respectively, during both the years. The production of organic acids during decomposition, which then helped in the release of native phosphorus by the solubilizing activity of these acids, could be attributed to the increase in accessible P with the application of fertilizers [7,9].

3.7 Available Potassium (kg ha⁻¹)

The data pertaining to available potassium after harvest of the paddy crop have been presented in Table 2.

The maximum amount of available potassium (235.6 & 237.9 kg ha⁻¹) was recorded with the application of 100% RDF + Nano urea + Nano potash + Nano Zn which remained statistically at par with 100% RDF + Nano potash spray at tillering stage (35 DAT) but significantly higher than rest of the treatment during both the years whereas lowest amount of available potassium (213.1 & 214.5 kg ha⁻¹) was recorded under control treatment respectively, during both the years.. Similar result was also observed by Du et al. [6] and Jhanzab et al. [7].

3.8 Available Zinc (kg ha⁻¹)

The data pertaining to available zinc after harvest of the paddy crop have been presented in Table 2.

The maximum amount of available zinc (0.80 g ha⁻¹) was recorded with the application of 100% RDF + Nano urea + Nano potash + Nano Zn which remained statistically at par with 100% RDF + Nano Zn spray at tillering stage (35 DAT) but significantly higher than rest of the treatment during 2022. Whereas during 2023 The maximum amount of available zinc (0.83 g ha⁻¹) was recorded with the application of 100% RDF + Nano urea + Nano potash + Nano Zn which was at par with 100% RDF + Nano Zn and 75% RDF + Nano urea + Nano potash + Nano Zn but higher than rest of the treatments. The lowest amount of available zinc (0.61 & 0.63 g ha⁻¹) was recorded under control treatment respectively, during both the years. The zinc level in soil increased due to nano zinc because it decreases their susceptibility to adsorption, fixation and precipitation in soil Xiao et al. [10] & Naderi and Shahraki (2013).

4. CONCLUSION

The investigation concluded that among the different foliar application of nano fertilizers, treatment T6 (100 % RDF + nano urea + nano potash + nano Zn) sprays at tillering stage (35 DAT) @ 4 ml per liter of water nano urea and nano potash and 2 ml per liter of water nano zinc was found to be the best treatment in chemical and physical properties of soil during both the years. The lowest value of all the parameters was recorded under control treatment during both the years.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. United State Department of Agriculture, Report. 2022-23:28.

2. Economics, Statistics & Evaluation Division, Department of Agriculture & Farmer Welfare 2021-22:29
3. Mandal S, Ghosh GK. Response of rice (*Oryza sativa* L.) to soil and foliar application of Nano-ZnO and bulk Zn-fertilizer in red acidic soil of West Bengal, India. Egyptian Journal of Soil Science. 2021;61(2):387-10.
4. Pandey G. Challenges and future prospects of agri-nanotechnology for sustainable agriculture in India. Environmental Technology and Innovation. 2018;11:299-307.
5. Gomez KA, Gomez AA. Statistical procedure for agriculture research 2nd edition Jon Wielly and Sons, New York; 1984.
6. Du W, Sun Y, Ji R, Zhu J, Wu J, Guo H. TiO₂ and ZnO nano-particles negatively affect wheat growth and soil enzyme activities in agricultural soil. Journal of Environmental Monitoring. 2011;13:822-828.
7. Jhanzab HM, Razaq A, Jilani G, Rehman, A, Hafeez A, Yasmeen F. Silver nano-particles enhance the growth, yield and nutrient use efficiency of wheat. International Journal of Agronomy and Agricultural Research. 2015;7(1):15-22.
8. Liu R, Lal R. Potentials of engineered nano particles as fertilizers for increasing agronomic productions. Science of the Total Environment. 2015;514: 131-139.
9. Naderi, M. and Shahraki, A. 2013. Nano-fertilizers and their roles in sustainable agriculture. International Journal of Agriculture and Crop Science. 2013;5(19): 2229-2232.
10. Xiao G, Zhao Z, Liang L, Meng F, Wu W. Guo Y. Improving nitrogen and water use efficiency in a wheat-maize rotation system in the North China Plain using optimized farming practices. Agricultural Water Management. 2013;212: 172–180.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/124550>