



# Restoring Degraded Lands in Maroua I Municipality, Cameroon: Application of Veazcomutech Techniques

Haiwa G. <sup>a\*</sup>, Djibrilla M. <sup>b</sup>, Kengne O. C. <sup>c</sup>, Kolah A. <sup>a</sup>,  
Nira G. <sup>a</sup>, Tchobsala <sup>d</sup> and Ngakou A. <sup>e</sup>

<sup>a</sup> Department of Environmental Sciences, National Advanced School of Engineering of Maroua, University of Maroua, P.O Box: 46 Maroua, Cameroon.

<sup>b</sup> Department of Plant Science, Faculty of sciences, University of Buea, P.O Box: 63 Buea, Cameroon.

<sup>c</sup> Department of Life and Earth Sciences, Higher Teachers' Training College, University of Maroua, P.O. Box 45 Maroua, Cameroon.

<sup>d</sup> Department of Biological Sciences, Faculty of Sciences, University of Maroua, P.O Box: 814 Maroua, Cameroun.

<sup>e</sup> Department of Biological Sciences, Faculty of Sciences, University of Ngaoundere, P.O Box: 454 Ngaoundere, Cameroun.

## Authors' contributions

This work was carried out in collaboration among all authors. Author HG conducted the field study and wrote the manuscript. Authors DM, KOC, KA, NG, Tchobsala and NA read and made suggestions and comments in order to improve the scientific quality of the document. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i101554>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124874>

Original Research Article

Received: 14/08/2024

Accepted: 16/10/2024

Published: 23/10/2024

\*Corresponding author: E-mail: [ghaiwa@yahoo.fr](mailto:ghaiwa@yahoo.fr);

**Cite as:** G., Haiwa, Djibrilla M., Kengne O. C., Kolah A., Nira G., Tchobsala, and Ngakou A. 2024. "Restoring Degraded Lands in Maroua I Municipality, Cameroon: Application of Veazcomutech Techniques". *Journal of Advances in Biology & Biotechnology* 27 (10):1341-62. <https://doi.org/10.9734/jabb/2024/v27i101554>.

## ABSTRACT

This study was carried out in the Sudano-Sahelian zone of Cameroon. It aims to contribute to the restoration of degraded land in the localities of Maroua I; to set up a sustainable soil restoration technique based on *Vetiveria zizanioides*- *Azadirachta indica*-eggshell and *Musa paradisiaca* peel; to evaluate the restoration of this land through the crop set up and to assess the state of the art of the "VEACOMUTECH" technique on an area of 103.5 m<sup>2</sup>. The method envisaged in this research is based on the new modern technique "VEAZCOMUTECH): Technique of *V. zizanioides*, *A. indica*, *Musa paradisiaca* and eggshell" with a view to restoring the fertility of the land and diversifying the population's sources of income by using this technique. The study was carried out in the Maroua I municipality on experimental plots at the Technical School of Agriculture. A control plot and a main plot treated with VEAZCOMUTECH were selected. The most vigorous species were chosen to measure growth and yield parameters over a three-week period. The results of this study show that the application of "VEAZCOMUTECH" on the site increases plant growth. In the first week, leaf length ranged from six (06) to fifteen (15) centimetres on average in the control plot. In the main plot where "VEAZCOMUTECH" was applied, leaf length ranged from ten (10) to thirty-two (32) centimetres during the same week. By the third week, leaf length had increased from 15 to 20 centimetres, while in the main plot, leaf length had increased from 32 to 80 centimetres. There was also a clear difference in the number of leaves at week three, with 20 and 87 leaves for the control and main plots respectively. The largest number of flower buds (23 flower buds) was observed on the plant in the main plot, compared with 2 flower buds in the control plot. These results could serve as a reference for the strategy of restoration and fertilization of degraded land in the Far North Region in particular and in the Sudano-Sahelian zone in general.

**Keywords:** Municipality; land restoration; veazcomutech; *Chrysopogon zizanioides*; flower buds.

## 1. INTRODUCTION

Soil degradation worldwide is mainly caused by human activities: intensive agriculture, irrigation, deforestation, overgrazing and industrial pollution [1]. Ranging from slight to severe degradation, this phenomenon affects about 20 million km<sup>2</sup>, or almost 60 % of arable land. Uncultivated soils are already degraded by water or wind erosion, salinization or deforestation [2, 3]. In 2001, at the 2nd International Conference on Land Degradation and Desertification, it was reported that desertification affects 33 % of the total land surface in the world, representing 42 million km<sup>2</sup> [4, 5]. However, it is often assumed that land degradation in Africa has had a very negative impact on agricultural ecosystems and food production, creating a handicap to achieving food security and improving livelihoods. Thus, the cost of action against soil degradation includes investments to restore degraded soils [6, 7]. In dry areas, farmers who have been living there for a long time have gradually accumulated experience and knowledge of the local environment that can serve as a starting point for improving management to increase the sustainable productivity of their soils, plants [8]. When we stop disturbing the soil, it is possible to gradually rebuild the vegetation cover and soil organic matter. Where there is a high demand for

crop residues to feed livestock, the biomass needs to be managed so that it improves productivity over a long period" [9]. In contrast, Cameroon has 12 million hectares of degraded land, including 8 million in its northern part. The incidence of poverty in this Sudano-Sahelian zone is increasing, mainly because of land degradation and the resulting loss of productivity [10]. However, the situation is worrying in the Far North Region of Cameroon, where soil degradation and rainfall variability are compounded by land saturation, low yields and difficulties in marketing certain products. Since 1960, the centre of gravity of the cotton production basin has been in the Far North of the country (Diamare and Mayo-Kani Divisions). However, this Region is heading for increasing drought, and above all an imbalance between uncontrolled population growth and available space. In 2010, Cameroon had a population of 19.5 million, with a demographic rate of 2.6 % per year, above the world rate of 2.6 % per year [11]. The application of appropriate, easy and inexpensive methods such as "VEAZCOMUTECH" to restore degraded soils will increase the yield of agricultural products in the Sudano-Sahelian zone. The main objective of this work is to contribute to the restoration of degraded lands in Maroua I. Specifically, the aims are to: set up a sustainable soil restoration

technique based on *Vetiveria zizanioides* - *Azadirachta indica*- *Musa paradisiaca* and eggshell peel; assess the restoration of these lands through the cultivation set up; take stock of the "VEACOMUTECH" technique on an area of 103.5 m<sup>2</sup>.

## 2. MATERIALS AND METHODS

### 2.1 Presentation of the Study Area

Maroua I is a district of the Sub-division of Cameroon, located in the Far North Region and more precisely in the Diamare Division, between the geographical coordinates 10°36'42"N latitude and 14°22'51"E longitude. It borders to the north by the municipalities of Maroua II and Meri, Mindif and Moutourwa to the south, Maroua III to the east and Gazawa and Ndoukoula to the west (Fig. 1).

The Maroua I Municipality covers an area of 660 km<sup>2</sup> and has an estimated population of 60,000 inhabitants, giving a population density of 91 inhabitants/km<sup>2</sup>. The population of Maroua I is cosmopolitan, made up of the Guiziga (the majority ethnic group), the Peulhs, the Haoussa,

the Bornouans, the Moufou, the Mafa, the Toupouri, the Massa, the Moundang, the Betis, the Bamilekes and the Sarahs. All these ethnic groups live together in perfect harmony [12].

The rainfall gradient in the study area varies between 500 and 1,000 mm. The annual rainfall is mainly concentrated in the 4 to 5 months from June to October [11]. The average annual temperature is around 28°C, with very large thermal differences between extreme values (7.7°C average annual temperature). In addition, the high variability of rainfall in space and time constitutes a major risk [12]. The relief of the Region is characterized by plains and mountains. It is not very hilly. In fact, the territory of the Maroua I Municipality is made up of plains topped by two mountains: Mount Makabaye and Mount Yamdjidjim [13]. The soils of the Maroua I Municipality are very diverse. There are bare, clay-textured soils found in the karals; the hardes; sandy-clay-textured soils and alluvial soils with a sandy-loamy texture, which are found in the plains and especially on the edges of the mayos and which support intense agricultural activity [14].




Fig. 1. Map showing the location of the study area

The Maroua I Municipality is irrigated by three mayos: the mayo Tsanaga which has its source in the mountains of the Mayo-Tsanaga Division, the mayo Boula which has its source in the northern Region and the mayo Kaliao which has its source in the mountains of Meri and the surrounding area. These rivers are irregular and dry for much of the year. Their surface run-off only lasts three to four months: July, August, September and October (rainy season) [15]. They cross the entire Municipality and divide, giving rise to the mayo Ziling. The dominant vegetation is the shrubby savannah characteristic of the Sudano-Sahelian zone. The flora is fairly diversified and includes species such as *Faidherbia albida*, *Zizyphus mauritiana*, *Tamarindus indica*, *Azadirachta indica*, *Acacia seyal*, etc. Some of these species are used in the production of oil and gas. Some of these species are used in traditional pharmacopoeia [16].

## 2.2 Study Materials

- ***Vetiveria zizanioides* (vetiver)**

*Chrysopogon zizanioides* formerly known as *Vetiveria zizanioides* L. Nash (Vetiver) is a perennial graminaceous plant. The vetiver system, based on the application of the vetiver species is the main restorative plant used in the VEAZCOMUTECH technique due to its integration in agroforestry systems, improving the efficiency of reforestation and bioengineering. The vetiver system (VS) is very simple, practical, cheap, extremely effective and requires very little maintenance. It is used for soil and water conservation, sediment control, soil stabilization,

rehabilitation and phytoremediation. As a plant, it is not harmful to the environment [17]. Vetiver has several parts: the leaves are sheathing, growing in clumps up to 1-2 m high, and are long and thin with serrated, sharp edges. The stems are straight and 1 to 3 m high. What's more, vetiver's extremely deep and thick root system, which can reach a depth of 3 to 4 m, attaches itself to the soil, making it very difficult to dislodge by high-speed water flows. The very deep, fast-growing root system also makes vetiver a very drought-tolerant plant, highly suitable for stabilizing steep slopes. This plant was the main material used to restore the soil in this study (Photo 1).

- ✓ The leaves of *Azadirachta indica* A. Juss. (Neem leaves) are evergreen and green in colour. They are broad, alternate and pinnate. They were used as an amendment and insect barrier in this study (Photo 2).
- ✓ The hen's eggshells, which are more or less resistant, mineralized envelopes covering the hen's eggs. They also act as a source of organic almonds and an insect barrier (Photo 3).
- ✓ The peel of *Musa paradisiaca* (Banana peel) is the outer part of the banana, also known as banana waste. It can be any colour, green or yellow, as long as it is ripe. In our study, we used chartreuse green peels, which acted as a rooting hormone, facilitating the attachment of *Vetiveria zizanioides* and providing kernels photo 4.



Photo 1. *Vetiveria zizanioides* (Vetiver grass)



**Photo 2. Leaves of *Azadirachta indica***



**Photo 3. Eggshells**



**Photo 4. *Musa paradisiaca* peel**



**Photo 5. *Solanum nigrum***

- ***Solanum nigrum* L.**

*Solanum nigrum* is one of the most common French domestic plants. It belongs to the Solanaceae family and is an annual herbaceous plant. It has branches bearing dark green, simple alternate leaves with toothed or wavy margins. In this experiment, it is used as a soil restoration indicator (Photo 5).

### 2.3 Methods

In order to achieve our research objectives, we adopted the "VEAZCOMUTECH" technique. This method involves several activities, such as collecting shells, neem leaves, banana peel, transplanting vetiver and growing our crop, *S. nigrum*, and monitoring plant growth parameters.

- **Choice and planning of the site**

Maroua I Municipality was chosen for this study. This choice was motivated by thesevere soil degradation caused by agricultural practices in this part of the Far North Region. After the choice of this Municipality, a prospective visit to the site was carried out in order to get a concrete idea of the realities on the ground before sampling. The aim of this mobilization was to recognize the site and familiarize ourselves with the study area. This gave us a superficial knowledge of the nature of the soil, the previous crops, which were maize and other legumes, and then the dimensions of our experimental site. The second stage consisted of laying out the site, the different plots and planting vetiver around the main plot. Vetiver cultivation began four days after the selection of the

experimental site, followed by the *S. nigrum* nursery two weeks later. The dimensions of the experimental area were measured using a decametre (dm), giving a surface area of 103.5 m<sup>2</sup>, i.e. a length of 11.5 m and a width of 9 m for the vetiver plot. The control plot was 9 m long and 2.5 m wide, with a total surface area of 22.5 m<sup>2</sup>. After this phase, the site was secured with stakes, thorns and used sacks.

- **Preparation of raw material**

This decisive stage consists of choosing our raw material and making the best choice among the different existing varieties, *Vetiveria zizanioides*, which is the best known variety in Haiti and in several countries around the world and used for its environmental benefits and in traditional medicine [18]. We began by collecting vetiver strains from Lake Porhi in the Mayo Kani Division, more precisely in the Porhi Sub-division, capital of Touloum. These strains were then kept in humid conditions until they reached the experimental plot to optimize transplanting (Photo 6).

- **Collecting the eggshells**

This phase involves carefully collecting the clean eggshells because they help to reduce soil acidity and combat parasites and diseases. They are also an organic fertilizer for plant growth and provide calcium for the plants. After the collection phase, we moved on to shredding using a mortar and pestle. Then we put 0.25 kg in each hole. See Plate 1, where A represents the collection stage, B the crushing and C the introduction into the holes.



**Photo 6. Vetiver after sampling**



**A**



**B**



**C**

**Plate 1. Stages in the use of eggshells**



**A**



**B**



**C**

**Plate 2. Stages in the processing of Neem leaves**



A



B

**Plate 3. Collecting the banana peel and introducing it into the holes**



A



B

**Plate 4. Preparing the holes and splitting the vetiver**

- **Collecting the green leaves of *Azadirachta indica***

This stage involves collecting the green leaves of neem because of its bioinsecticidal, bionematicidal, bioacaricidal, bioantifungal and biofertilizing properties, then drying the leaves in the shade, as this technique allows the leaves to retain their natural colour and better preserve the vitamins and mineral salts. After two weeks of drying, we moved on to crushing the leaves using a pestle. A quantity of 0.25 kg was introduced into each hole that would receive the vetiver and *Solanum nigrum*. See Plate 2 with A: collection, B: drying and C: crushing.

- **Obtaining the *Musa paradisiaca* peel**

This stage allowed us to collect banana peels and bananas in poor condition from banana sales outlets at the Maroua artisanal market and then from rubbish bins. The banana peel is used

in our plot because it has many properties and virtues. It is rich in fibre, potassium, magnesium, iron, copper and vitamins (A, B6, B9, B12 and C). We sorted the collected peels and cut them into pieces using a knife, then buried 0.25 kg in the holes as shown in Plate 3 with A: banana peel and B: introduction into the hole.

- **Soil preparation, breaking up the vetiver**

We began by preparing the soil by digging the vetiver holes 40 cm deep using a crowbar at 80 cm intervals to create a loose germination bed with a very wide crown. This is followed by the removal of the mature vetiver clumps. Transplanting involves dividing the tillers from a mother clump, which require care, so that each plant has at least two or three splinters and part of the crown. After separation, the splinters are cut to a length of 20 cm, as shown in Fig. 4, with A: the preparing of the holes and B: the splitting of the vetiver plant, Plate 4.



- **Planting (transplanting) the vetiver**

This consists of introducing the splinters (shoots) into the holes around the vetiver plot. The vetiver chips obtained are introduced into the holes containing the *Musa paradisiaca* peel, the eggshells and the Neem leaves. In this way, the planting points are known, the vetiver is transplanted like rice and the roots are firmly planted. Planting should be done after a good watering because the soil must be well soaked with water, and the vetiver plants should be watered abundantly immediately after planting until the new leaves appear. Watering should be done twice a day during the first week and once in the evening from the second week to minimize evaporation losses during the day. The third week after transplanting is used to weed around the vetiver plot before watering (Photo 7).

- **Cultivation of *Solanum nigrum***

- **Soil preparation and sowing of *Solanum nigrum***

This consists of ploughing the inside of the vetiver and control plots with a hoe, then burying the Neem litter and eggshells. The nursery is watered beforehand, then the plots where the young seedlings will be planted, and proceed to mark out the holes with a 30 cm spacing between the lines and the seedlings. Next, we transplant vigorous, healthy plants with 3 to 4 true leaves, with one plant per hole. Transplanting should be done in the evening to allow the seedlings to take root properly, then watered immediately afterwards.

- **Labelling the plots**

One month after rooting the *Vetiveria zizanioides*, we transplanted the *S. nigrum* and made 21 plates indicating the different plots and the plants to be chosen, then took their

parameters. The system chosen for this purpose was a block containing two (02) plots, illustrated in Plate 5, with A the main plot and B the control plot.

- **Overall process of the VEAZCOMUTECH technique**

The process of implementing the new land restoration technique in the Maroua I Municipality involved the following nine (09) stages (Fig. 3):

- **Monitoring of *Solanum nigrum***

- **Monitoring parameters of *Solanum nigrum***

Data collection in the field was carried out over a period of one (01) month, depending on the growth of *Solanum nigrum*. This stage consisted of selecting 18 plants from the main plot and 3 from the black nightshade control plot on which we recorded the parameters every week. The parameters monitored were: height, number of leaves, flowers and flower buds, and success rate. The analysis of all these parameters allowed us to evaluate the effectiveness of the new modern "VEAZCOMUTECH" technique.

- **Statistical analysis**

All the data collected were statistically tested using STATISTICA version 7.1 software. An analysis of variance (ANOVA) was used to assess the effects of the fertilizer treatments on plant growth parameters and yields. Where there was a significant difference between treatments, the Newman-Keuls multiple comparison test with a threshold of 5% was used to classify them into homogeneous groups. Correlation between plots was performed using principal component analysis of the variables. These tests were carried out using the EXCEL 2016 spreadsheet and XLSTAT 2007 software.



**Photo 7. Vetiver transplanting**




Fig. 2. Experimental design of the growing environment




Fig. 3. Overall land restoration process using the VEAZCOMUTEC technique



Plate 5. Labelling of plots

### 3. RESULTS

#### 3.1 Development of a Sustainable Soil Restoration Technique Based on *Vetiveria zizanioides*, *Azadirachta indica*, Eggshell and *Musa paradisiaca* peel

##### 3.1.1 Plant growth on the experimental site

Plate 6 illustrates vetiver growth up to June 2023 with A: vetiver growth one week after transplanting, B: vetiver growth after two weeks and C: vetiver growth after nine weeks. The results of this study show that the application of "VEAZCOMUTECH" on the site increases plant growth. This development depends on the plant's ability to emerge from its sluggish life. However, it is not uniform throughout the plot. In addition, some plants have lost the leaves of the transplants, while others are still growing.

##### 3.2 Effect of Veazcomutech on the average length of *Solanum nigrum* leaves

The analysis of this parameter for monitoring the length of *Solanum nigrum* leaves using the "VEAZCOMUTECH" technique was carried out over a period of three weeks. In this study, P represents the block plot and M the *S. nigrum* (Fig. 4).

Observation of these two Fig. 5 shows a variation in the average size of each plant, but there is a considerable difference between the main and control plots ( $P < 0.05$ ). We also note that leaf

length in the main plot varies from ten (10) to thirty-two (32) cm in P1B1M4, the shortest leaf, and P1B1M6, the longest leaf. This was also the case for P1B2M10, which was only monitored during the first week. That of the control plot ranged from six (06) to fifteen (15) cm in length, with P0M2 the shortest leaf and P0M1 the longest. This Fig. 5 clearly shows the difference in leaf length, where A was the plant with the longest average leaf length in the main plot and B was the control.

##### 3.3 Effect of VEAZCOMUTECH on the average height of *Solanum nigrum*

Analysis of the average height of *Solanum nigrum* over three weeks using the "VEAZCOMUTECH" technique enables us to assess the height of the plants (Fig. 6).

The average height of the control is shown in Fig. 6.

We can see a great difference between the plants in the main plot ( $P < 0.05$ ), whose height varies between 49 and 80 cm, with P1B1M7 the smallest plant and P1B2M13 the tallest. However, the average height of the control varied between 08 and 20 cm, with P0M1 being the tallest. In addition, the difference in height at the second week was not very great for some plants, such as P1B1M1, because they were affected by the strong wind, and it should be noted that the plants in the control plot are dwarfed. Plate 8 below shows the tallest plants in the two plots, with A the plant from the main plot and B the control.




Plate 6. Vetiver growth





Fig. 4. Changes in average leaf length in the main plot

P1B1M1: Morelle 1 in block 1 of plot 1; P1B1M2: Morelle 2 in block 1 of plot 1; P1B1M3: Morelle 3 in block 1 of plot 1; P1B1M4: Morelle 4 in block 1 of plot 1; P1B1M5: Morelle 5 in block 1 of plot 1; P1B1M6: Morelle 6 in block 1 of plot 1; P1B1M7: Morelle 7 in block 1 of plot 1; P1B1M8: Morelle 8 in block 1 of plot 1; P1B2M9: Morelle 9 in block 2 of plot 1; P1B2M10: Morelle 10 in block 2 of plot 1; P1B2M11: Morelle 11 in block 2 of plot 1; P1B2M12: Morelle 12 in block 2 of plot 1; P1B2M13: Morelle 13 in block 2 of plot 1; P1B2M14: Morelle 14 in block 2 of plot 1; P1B3M15: Morelle 15 in block 3 of plot 1; P1B3M16: Morelle 16 in block 3 of plot 1; P1B3M17: Morelle 17 in block 3 of plot 1; P1B3M18: Morelle 18 in block 3 of plot 1.

### 3.4 Effect of VEACOMUTECH on the average number of leaves of *Solanum nigrum*

The analysis of the data obtained from monitoring the average number of leaves of

*Solanum nigrum* over three weeks is shown in Table 1, which represents the change in the number of leaves of the main plot in which the VEACOMUTECH technique was applied, and Table 2 for the control plot.



**Fig. 5. Changes in leaf length in the control plot**  
*PoM1: Morelle 1 in the control plot (p0), PoM2: Morelle 2 in the control plot (p0)*



**A**




**B**

**Plate 7. Plants with the longest leaves in the two plots**




**Photo 8. Impacted plant P1B2M10**



**Fig. 6. Changes in the evolution plant height in the plots**

*P1B1M1: Morelle 1 in block 1 of plot 1; P1B1M2: Morelle 2 in block 1 of plot 1; P1B1M3: Morelle 3 in block 1 of plot 1; P1B1M4: Morelle 4 in block 1 of plot 1; P1B1M5: Morelle 5 in block 1 of plot 1; P1B1M6: Morelle 6 in block 1 of plot 1; P1B1M7: Morelle 7 in block 1 of plot 1; P1B1M8: Morelle 8 in block 1 of plot 1; P1B2M9: Morelle 9 in block 2 of plot 1; P1B2M10: Morelle 10 in block 2 of plot 1; P1B2M11: Morelle 11 in block 2 of plot 1; P1B2M12: Morelle 12 in block 2 of plot 1; P1B2M13: Morelle 13 in block 2 of plot 1; P1B2M14: Morelle 14 in block 2 of plot 1; P1B3M15: Morelle 15 in block 3 of plot 1; P1B3M16: Morelle 16 in block 3 of plot 1; P1B3M17: Morelle 17 in block 3 of plot 1; P1B3M18: Morelle 18 in block 3 of plot 1.*



**Fig. 7. Changes in the average height of plants in the control plot**

*PoM1: Morelle 1 in the control plot (p0), PoM2: Morelle 2 in the control plot (p0)*



A



B

**Plate 8. Average height of plants in the main and the control plots**

**Table 1. Average change in the number of leaves in the main plot.**

	Week 1	Week 2	Week 3
<b>P1B1M1</b>	11	35	51
<b>P1B1M2</b>	12	31	72
<b>P1B1M3</b>	15	29	59
<b>P1B1M4</b>	13	23	49
<b>P1B1M5</b>	10	20	29
<b>P1B1M6</b>	11	19	42
<b>P1B1M7</b>	11	24	76
<b>P1B1M8</b>	14	33	86
<b>P1B2M9</b>	11	29	60
<b>P1B2M10</b>	8	0	0
<b>P1B2M11</b>	12	27	58
<b>P1B2M12</b>	12	36	87
<b>P1B2M13</b>	11	28	73
<b>P1B2M14</b>	12	28	71
<b>P1B3M15</b>	10	21	63
<b>P1B3M16</b>	10	22	51
<b>P1B3M17</b>	12	21	44
<b>P1B3M18</b>	4	15	39
<b>Average/SD</b>	11.06±2.36	24,5±8.41	56,11±21.28

*P1B1M1: Morelle 1 in block 1 of plot 1; P1B1M2: Morelle 2 in block 1 of plot 1; P1B1M3: Morelle 3 in block 1 of plot 1; P1B1M4: Morelle 4 in block 1 of plot 1; P1B1M5: Morelle 5 in block 1 of plot 1; P1B1M6: Morelle 6 in block 1 of plot 1; P1B1M7: Morelle 7 in block 1 of plot 1; P1B1M8: Morelle 8 in block 1 of plot 1; P1B2M9: Morelle 9 in block 2 of plot 1; P1B2M10: Morelle 10 in block 2 of plot 1; P1B2M11: Morelle 11 in block 2 of plot 1; P1B2M12: Morelle 12 in block 2 of plot 1; P1B2M13: Morelle 13 in block 2 of plot 1; P1B2M14: Morelle 14 in block 2 of plot 1; P1B3M15: Morelle 15 in block 3 of plot 1; P1B3M16: Morelle 16 in block 3 of plot 1; P1B3M17: Morelle 17 in block 3 of plot 1; P1B3M18: Morelle 18 in block 3 of plot 1. SD: Standard deviation.*

**Table 2. Mean numbers of leaves in the control plot**

	Week 1	Week 2	Week 3
<b>PoM1</b>	6	9	15
<b>PoM2</b>	4	6	8

*PoM1: Morelle 1 from the control plot (p0), PoM2: Morelle 2 from the control plot (p0)*



**Plate 9. Difference between the numbers of leaves in the two plots**

The average number of leaves in the control plot is given in Table 2.

In the main plot, the difference in the number of leaves was very small and even constant in plant P1B2M10 during the first week. Moreover, from the second week, we observe that the number of leaves increases exponentially in all the plants, following the example of P1B2M12, which has an average of 87 leaves. As for the control plot, the leaves were dwarfed, but P0M2 had more difficulty in increasing the number of leaves than P0M1. Plate 9 shows this difference with (A) the plant with the highest average number of leaves in the main plot and (B) the plant with the lowest average number of leaves in the control. In the present study, "VEAZCOMUTECH" showed the best results in several growth parameters of black nightshade. This is explained by the fact that the concentrations of nutrients N, P and K were higher than those of the control plot, which were zero.

### 3.5 Effect of VEAZCOMUTECH on the average number of flower buds of *Solanum nigrum*


The analysis of the average number of flower buds parameter on the evolution of *S. nigrum* allows us to evaluate the effectiveness of the "VEAZCOMUTECH" technique on the plants in the main plot and compare it with the control. Fig. 8 illustrates the effectiveness of this technique.

The analysis of the graph for the main plot shows the absence of flower buds in plants P1B2M10 and P1B3M18 during the period when the parameters were taken, whereas P1B1M8 at 40 represents the highest average number of flower buds. In contrast to the control plot, P0M1 only showed the appearance of four small flower buds at the last measurement in the third week. Plate 10 illustrates these graphs with (A) plant 8 from the control plot and (B) plant 1 from P0.




**Plate 10. Difference between the average number of flower buds in the two plots**






**Fig. 8. Average change in the number of flower buds in the main plot**

*P1B1M1* : Morelle 1 in block 1 of plot 1; *P1B1M2* : Morelle 2 in block 1 of plot 1; *P1B1M3* : Morelle 3 in block 1 of plot 1; *P1B1M4* : Morelle 4 in block 1 of plot 1; *P1B1M5* : Morelle 5 in block 1 of plot 1; *P1B1M6* : Morelle 6 in block 1 of plot 1; *P1B1M7* : Morelle 7 in block 1 of plot 1; *P1B1M8* : Morelle 8 in block 1 of plot 1; *P1B2M9* : Morelle 9 in block 2 of plot 1; *P1B2M10* : Morelle 10 in block 2 of plot 1; *P1B2M11* : Morelle 11 in block 2 of plot 1; *P1B2M12* : Morelle 12 in block 2 of plot 1; *P1B2M13* : Morelle 13 in block 2 of plot 1; *P1B2M14* : Morelle 14 in block 2 of plot 1; *P1B3M15* : Morelle 15 in block 3 of plot 1; *P1B3M16* : Morelle 16 in block 3 of plot 1; *P1B3M17* : Morelle 17 in block 3 of plot 1; *P1B3M18* : Morelle 18 in block 3 of plot 1.



**Fig. 9. Average change in the number of flower buds in the control plot**

*PoM1*: Morelle 1 in the control plot ( $p_0$ ), *PoM2*: Morelle 2 in the control plot ( $p_0$ )



**Fig. 10. Average change in the number of flowers in the main plot**

P1B1M1 : Morelle 1 in block 1 of plot 1; P1B1M2 : Morelle 2 in block 1 of plot 1; P1B1M3 : Morelle 3 in block 1 of plot 1; P1B1M4 : Morelle 4 in block 1 of plot 1; P1B1M5 : Morelle 5 in block 1 of plot 1; P1B1M6 : Morelle 6 in block 1 of plot 1; P1B1M7 : Morelle 7 in block 1 of plot 1; P1B1M8 : Morelle 8 in block 1 of plot 1; P1B2M9 : Morelle 9 in block 2 of plot 1; P1B2M10 : Morelle 10 in block 2 of plot 1; P1B2M11 : Morelle 11 in block 2 of plot 1; P1B2M12 : Morelle 12 in block 2 of plot 1; P1B2M13 : Morelle 13 in block 2 of plot 1; P1B2M14 : Morelle 14 in block 2 of plot 1; P1B3M15 : Morelle 15 in block 3 of plot 1; P1B3M16 : Morelle 16 in block 3 of plot 1; P1B3M17 : Morelle 17 in block 3 of plot 1; P1B3M18 : Morelle 18 in block 3 of plot 1.

**Table 3. Evolution of the average number of flowers from P0**

	Week 1	Week 2	Week 3
P0M1	0	0	0
PoM2	0	0	0

PoM1 : Morelle 1 of the control plot (p0), PoM2 : Morelle 2 of the control plot (p0)

**Table 4. VEACOMUTECH Status Report**

Constituents	Quantity (kg)	Unit price in FCFA	Total price in FCFA
Neem leaves	10,25	100	1025
Eggshells	10,25	150	1537,5
Banana peel	10,25	100	1025
The Vetiver	27	50	1350
Totals	57,75	400	4937,5

For 1 hectare we will need 990.33 kg of Neem leaves, 990.33 kg of banana peels, 990.33 kg of eggshells and 2608.69 kg of vetiver grass. So we could spend 477.050 FCFA to restore 10,000 m<sup>2</sup> using the "VEAZCOMUTECH" technique

### 3.6 Effect of VEACOMUTECH on the average number of flowers of *Solanum nigrum*

The analysis of the average number of flowers of *S. nigrum* allows us to assess the effectiveness of this technique by comparing the data of the main plot with those of the control plot. The

following graph shows the evolution of the number of flowers in the main plot (Fig. 10).

The analysis of this parameter shows that from the first week to the second harvest in week 2, only plant P1B2M14 had two flowers. We also found that plants P1B1M1, P1B1M4, P1B2M10, P1B2M12, P1B2M13, P1B3M16 and P1B3M18

had no flowers throughout the period during which the parameters were measured. In addition, P1B1M3 has 19 flowers, which is the highest number of flowers in the entire main plot, while the control plant has no flowers at all.

### 3.7 Overview of the VEAZCOMUTECH technique

It is necessary to briefly describe in great detail all the components that contributed to the development of the "VEAZCOMUTECH" technique. To do this, we determined the cost of restoring 103.5 m<sup>2</sup> of the experimental plot and 1 hectare. To restore 103.5 m<sup>2</sup>, we used 10.25 kg of banana peels, 10.25 kg of eggshells and 3 tufts of vetiver grass, which yielded 55 plants after splitting, including 27 kg. Details of the costs involved in setting up the modern land restoration technique for 103.5 m<sup>2</sup> are given in Table 4.

## 4. DISCUSSION

Agriculture plays an important role in the socio-economic development in the northern Region of Cameroon. As such, its development is a priority for the Cameroonian government. Many efforts have been made to achieve sustainable growth in agricultural production. The favourable effect of organic fertilizers observed on aubergine growth compared with the control plot (P0) could be due to their potential to improve soil structure and aeration, and its water and nutrient retention capacity in particular [18]. This would contribute to good penetration of the roots into the soil during their growth and an increase in water and mineral absorption by these organs. The resulting physiological consequences for aubergines were a rapid proliferation of root biomass, with a consequent increase in height and in the number of leaves and shoots per plant. The main plot that received the "VEAZCOMUTECH" treatment showed considerably better vegetative growth than the control plot. This could be the result of good assimilation by the plants of the nitrogen contained in the chicken droppings, due to the abundance of potassium and phosphorus in the banana peel. In fact, adequate phosphorus and potassium nutrition can increase crop growth response [19]. The organic potash used in this work did not consist solely of potassium, according to the characterisation carried out on banana peel [18]. In addition to potassium, it contained minerals such as phosphorus, magnesium and calcium. This difference in

growth may be due to environmental factors, in particular sunshine, since the temperature was over 40°C and young plants exposed to the sun all day were unable to withstand the heat, whereas those that were able to withstand the very high temperature benefited from the shade of a tree in the neighbouring field. With regard to flower bud production, the highest yield parameters in this experiment were obtained following the "VEAZCOMUTECH" treatment. The P0 control without fertilizer produced the lowest yields. These results corroborate studies carried out on pineapple [20,21]. There appears to be a synergistic effect due to the combination of organic and mineral matter, which favours the mineral and water supply of crops [22]. This in turn contributes to good development of flower buds and increased yields [12,23]. Under these conditions, the additional effect of the high potassium content in plantain peel potash to that of the nitrogen contained in the organic matter would have been decisive for a high production of flower buds in nightshade, as also mentioned in citrus [24, 25]. The lack of flower buds in plant P1B3M18 is probably due to the replacement carried out a week after transplanting, which slowed down its growth. In addition, the excessive increase in the number of flower buds in plant P1B1M8 would be the reason for its location, which is close to the vetiver. In addition, block 1 of the main plot and the entire control benefited from the shade of a tree located next to the plot. The absence of flowers on several plants in the main plot is probably due to their young age, as they were replaced a week after transplanting, unlike the number of flowers on plant P1B2M3, which is thought to be due to old age, as it is over the age of the first cut. As for the control plot, its number of flowers is due to the size of its population, which is dwarfed, and to the difficulty of growing on this land. In the control plot, the absence of flower buds during the first and second cut would be due to adaptation problems in the new, barren environment. This non-uniformity is linked to the trauma suffered by the plants during transport. Cameroon, like other Sahelian countries, is confronted with the phenomena of climate change and variability, which have caused, among other things, severe land degradation [26]. In recent years, however, the sector has been affected by continued degradation and a decline in the fertility of agricultural land [27]. However, the successful implementation of the strategy requires the commitment and participation of everyone in order to reverse soil degradation in a sustainable way, so as to make

it healthy and more productive, with a view to making a better contribution to achieving food sovereignty and the sustainable development aims. To this end, [28] demonstrate that organic fertilizers (goat dung, cow dung and chicken dung) increase the height, leaf length, number of leaves and number of fruits of tomato. This difference in the length of *Solanum nigrum* leaves between the main and control plots may be due to the application of *Vetiveria zizanioides*, Neem leaves, eggshells and *Musa paradisiaca* peels in the main plot, while the control plot was not amended. The absence of P1B2M10 could be due to a waterborne disease and this week's strong winds. Similar results were obtained with Chinese cabbage, whose vegetative growth parameters were improved by the addition of organic fertilizers such as chicken manure compost [29, 30]. Colloff [31] showed that other types of fertilizer, such as termite mound soil alone without any combination, improved the growth parameters (stem height, diameter at the crown and number of leaves produced by the plants) of three black nightshade cultivars, due to the beneficial effect of the presence of nitrogen and phosphorus contained in termite mound soil. The climate-smart objective is to improve food and nutritional security, thereby strengthening peace and community cohesion [32, 33].

## 5. CONCLUSION

In definitive, the main objective was to contribute to the restoration of degraded land in the municipality of Maroua I. In support of our ideas, this sustainable restoration technique based on *Vetiveria zizanioides*, *Azadirachta indica* leaves, eggshells and *Musa paradisiaca* peels not only helps to clean up the environment, but also contributes to the recovery of biodegradable waste - Neem leaves, eggshells and banana peels. It also contributes to phytoremediation and biodiversity conservation by protecting the vetiver species. Secondly, the evaluation of the effectiveness of the restoration of this land through the cultivation of black nightshade shows that the growth of plants in the plot where the "VEAZCOMUTECH" technique was applied is faster than that of the control on all parameters, such as the length of the leaves of the P1B1M6 plants, which is 34 centimetres, while the largest plant in the control plot is 15 centimetres. In addition, plants such as P1B2M13 in the main plot were 90 cm tall, while the tallest plant in the control was 22 cm. In addition, the new, modern "VEAZCOMUTECH" technique effectively restores the soil fertility. This new technique,

based on the use of local biodegradable natural resources at lower cost and in an ecologically sustainable way, could help to reduce farmers' expenses, preserve the environment, restore soil fertility on a long-term basis and guarantee the quality of agricultural produce. The restoration of degraded soils using the "VEAZCOMUTECH" technique remains of great interest to farmers in the Sudano-Sahelian zones.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

## DECLARATION OF DATA AVAILABILITY

All data used in the article and in materials and methods have been referenced in the article.

## ACKNOWLEDGEMENTS

We would like to sincerely thank the administrative and traditional authorities and the local populations of Diamare Division for their active participation in carrying out this work.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Sawadogo J, Coulibaly PJA, Bambara FJ, Sawadogo AC, Compaore E & Legma JB. Effects of organic fertilizers on soil physicochemical parameters and on onion (*Allium cepa* L.) productivity in the centre-west region of Burkina Faso, Africa. *SCIENCE*. 2020;17(6):44 – 57.
2. Kouamé N, Beugré MM, Kouassi NJ & Yatty KJ. Response to organic and mineral fertilisation of two varieties of okra (*Abelmoschus esculentus* (L) moench, malvacea) in Daloa, Ivory Cost. *International Journal of Advanced Research*. 2021;9(06):51 – 60.
3. Ouédraogo W, Sawadogo J, Bernadette M, Coulibaly PJ, Savadogo CA, Kaboré A & Legma JB. Comparative study of the quality of three organo- biological amendments on tomato production at Soala in central-western Burkina Faso. *Science et technique, Sciences naturelles et appliquées*. 2019;38(1):42.

4. Smith A, Snapp SS, Dimes J, Gwenambira C & Chikowo R. Doubled-up legume rotations improve soil fertility and maintain productivity under variable conditions in maizebased cropping systems in Malawi Agricultural Systems. 2016;145:139 – 149.
5. Kitabala MA, Tshala UJ, Kalenda MA, Tshijika IM & Mufind KM. Effects of different doses of compost on the production and profitability of tomato (*Lycopersicon esculentum* mill) in the town of Kolwezi, Lualaba province (DR Congo). Journal of Applied Biosciences. 2016;102(1):9669 – 9679.
6. Alexandratos, N et Bruinsma J. World agriculture towards 2030/2050: The 2012 Revision, ESA Working Paper No.12–03, Rome: FAO. 2012;72-159.
7. Favretto N, Dougill A, Stringer L, Afionis S and Quinn C. Links between climate change mitigation, adaptation and development in land policy and ecosystem restoration projects: Lessons from South Africa. Sustainability. 2018;10(3):779
8. Lahmar R, Bationo BA, Lamso ND, Guéro Y, Titonell P. Tailoring conservation agriculture technologies to West Africa semi-arid zones: building on traditional local practices for soil restoration. Field Crops Res. 2012;132:158-167.
9. Kassam A, Friedrich T, Shaxson F, Reeves T, Pretty J, de Moraes Sâ JC. Production systems for sustainable intensification: integrating productivity with ecosystem services. Technikfolgenabschätzung. (pAO). 2012;38-45.
10. MINEPDEP. Launch workshop of the project entitled 'Land degradation neutrality and mitigation of greenhouse gas emissions in the Afro-ecological zone of Sudano-Sahelian Cameroon. 2021;130.
11. MINADT. Lokanadhan S, Muthukrishnan P. et Jeyaraman S., 2012. Neem products and their agricultural applications. JBiopest. 2010;5:72 – 76.
12. Bayé-Niwah C, Mapongmetsem PM. Farmers' perceptions of *Moringa oleifera* Lam. (Moringaceae) in the Sudano-Sahelian zone of Cameroon. International Journal of Innovation and Scientific Research. 2018;39(1):91-102.
13. Abib FC, Ibrahima A, Ntoupka M, Tapsou JM, Harmand JM, Thaler P, Peltier R, Dreyer E. Inter-provenance valorization of gum arabic production by *Acacia senegal* (L.) Will in relation to tree growth and water use efficiency. Afrique Science. 2016;12(2):135-150.
14. Aboubakar. Effects of amendments on the growth, yield and chemical composition of black nightshade (*Solanum nigrum* L.) in a peri-urban area of Yaoundé, Cameroon. Int. J. Biologie Chemie. Science. 2020;14(6):2134-2146pp. Available:https://dx.doi.org/10.4314/ijbcs.v14i6.16
15. Bidzoni. City/village: Maroua (Far North Region/Department. 2019;5-10.
16. Kouadio EJP, Konan KH, Brou K, Dabonne S, Due AE and Kouame LP. Studies of some growth and nutritional value parameters of *Archachatina marginata* (Swainson) snail varieties reared in the wild. Tropicultura. 2015;33(1):38 – 45.
17. Toundou ON. Evaluation of the chemical and agronomic characteristics of five waste composts and study of their effects on soil chemical properties, physiology and yield of maize (*Zea mays* L. Var. Ikenne) and tomato (*Lycop ersicum esculentum* L. Var. Tropimech) under two water regimes in Togo. PhD, University of Limoges. 2016;214.
18. Alla KT, Bomisso EL, Ouattara G et Dick AE. Effect of fertilisation with plantain peel by-products on the agromorphological parameters of the F1 aubergine variety Kalenda (*Solanum melongena*) in the locality of Bingerville in Ivory Coast. Journal of Animal and plant Sciences. 2018;38(3):6292 – 6306.
19. Gala bi TJ, Camara M, Yao-Kouame A, Keli zJ. Profitability of mineral fertilisers in rainfed upland rice production: The case of the Gagnoa area in west-central Ivory Coast. Journal of Applied Biosciences. 2011;46:3153 – 3162.
20. Kotaix AJA, Angui TKP, Bakayoko S, Kassin KEL, N'goran KE, N Kouame, Koné B et Pierre CZK. Effects of liquid organic fertilizer (NPK 5-9- 18) and mineral fertilizer (NPK 12-11-18) on soil organic matter and tomato yield in southern and central-western Ivory Coast. Journal of Animal & Plant Sciences. 2019;41(3):7055 - 7067
21. Mukalay MJ, Shutcha MN, Tshomba KJ, Mulowayi K, Kamb CF et Ngongo LM, Causes of high plant heterogeneity in a maize field under the soil and climate conditions of Lubumbashi. University Press of Lubumbashi, Annals of Faculty of Agronomic Sciences. 2013;1(2):4 – 11.

22. Kitabala MA, Tshala UJ, Kalenda MA, Tshijika IM, Mufind KM. Effects of different doses of compost on tomato (*Solanum lycopersicum*) production and profitability in the town of Kolwezi, Lualaba Province (DR Congo). Journal of Applied Biosciences. 2016;102:9669 – 9679.
23. Akanza KP et Yao KA. Organomineral fertilisation of cassava (*Manihot esculenta*) and diagnosis of soil deficiencies. Journal of Applied Biosciences. 2011;46:3163 – 3172.
24. Quaggio JA, Junior DM and Boaretto RM. Sources and rates of potassium for sweet orange production. Scientia Agricola (Piracicaba, Braz.). 2011;68(3):369 – 375.
25. Yasin M, Ashraf M, J Yaqub, Akhtar M, Athar K, Alikhan M et Ebert G, Controlling excessive fruit drop and improving yield and juice quality of Kinnow (*Citrus deliciosa* X *Citrus nobilis*) through nutrient management. Pakistan Journal of Botany. 2012;44:259 – 265.
26. Dixon SJ, Sear DA, Odoni NA, Sykes T and Lane SN. The effects of river restoration on catchment scale flood risk and flood hydrology. Earth Surface Processes and Landforms. 2016;41(7): 997–1008.
27. Djoudi H, Brockhaus M and Locatelli B. Once there was a lake: Vulnerability to environmental changes in northern Mali. Regional Environmental Change. 2011; 13(3):493-508.
28. Batamoussi MH, Tovihoudji PG, Tokore OMSBJ, Boulga J, Essegnon MI. Effect of organic fertilizers on the growth and yield of two varieties of tomato (*Lycopersicon esculentum*) in the commune of Parakou (North Benin). Int. J. Sci. 2016;24(1):86-94.
29. Kimuni NL, Mwali KM, Mulembo MT, Lwalaba WLJ, Lubobo KA, Katombe NB, Mpundu MM, Baboy LL. Effects of increasing doses of chicken manure composts on the yield of Chinese cabbage (*Brassica chinensis* L.) planted on an acidic soil of Lubumbashi. J. Appl. Biosci. 2014;(77):6509 – 6522.
30. Kpangba TPMK, Mpika J, Makoundou A, Bita MA and Attibayéba. Effect of termite mound soil on the growth of three cultivars of black nightshade (*Solanum nigrum* L.: Solanaceae) grown in Brazzaville (Congo). Int. J. Biol. Chem. Sci. 2020;14(1):192-203.
31. Colloff MJ, Martín-López B, Lavorel S, Locatelli B, Gorddard R, Longaretti PY, Walters G, Kerkhoff LV, Wyborn C, Coreau A. An integrative research framework for enabling transformative adaptation. Environmental Science & Policy. 2017;68:87–96.
32. Likoko BA, Murefu K, Likoko AG et Posho NB. Effects of woody legume biomass on maize growth and yield in the corridors of a ferralsol in Yangambi, DR Congo, Journal of Applied Biosciences. 2018;131:13382 – 1339.
33. Kitabala MA, Tshala UJ, Kalenda MA, Tshijika IM et Mufind KM. Effects of different doses of compost on the production and profitability of tomato (*Lycopersicon esculentum* Mill) in the town of Kolwezi, Lualaba Province (DR Congo). Journal of Applied Biosciences. 2016; 102:9669 – 9679.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/124874>